arXiv:1107.3266v1 [math.CA] 17 Jul 2011

Dissertationes Math. (Rozprawy Mat.) (to appear)

DACHUN YANG and SIBEI YANG

Weighted local Orlicz-Hardy spaces with applications to pseudo-differential operators

Dachun Yang
School of Mathematical Sciences
Beijing Normal University
Laboratory of Mathematics and Complex Systems
Ministry of Education
Beijing 100875
People's Republic of China
E-mail: dcyang@bnu.edu.cn

Sibei Yang
School of Mathematical Sciences
Beijing Normal University
Laboratory of Mathematics and Complex Systems
Ministry of Education
Beijing 100875
People's Republic of China
E-mail: yangsibei@mail.bnu.edu.cn

Contents

1. Introduction 5
2. Preliminaries 9
    2.1. $A_p^{\rm loc}(\mathbb{R}^n)$ weights 9
    2.2. Orlicz functions 12
3. Weighted local Orlicz-Hardy spaces and their maximal function characterizations 13
4. Calderón-Zygmund decompositions 29
5. Weighted atomic decompositions of $h_{\omega,\,N}^{\Phi}(\mathbb{R}^n)$ 35
6. Finite atomic decompositions 42
7. Dual spaces 50
8. Some applications 60
References 78

Abstract

Let Φ be a concave function on $(0, \infty)$ of strictly lower type $p_\Phi \in (0, 1]$ and $\omega \in A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$ (the class of local weights introduced by V. S. Rychkov). We introduce the weighted local Orlicz-Hardy space $h_\omega^\Phi(\mathbb{R}^n)$ via the local grand maximal function. Let $\rho(t) \equiv t^{-1}/\Phi^{-1}(t^{-1})$ for all $t \in (0, \infty)$. We also introduce the BMO-type space $\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$ and establish the duality between $h_\omega^\Phi(\mathbb{R}^n)$ and $\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$. Characterizations of $h_\omega^\Phi(\mathbb{R}^n)$, including the atomic characterization, the local vertical and the local nontangential maximal function characterizations, are presented. Using the atomic characterization, we prove the existence of finite atomic decompositions achieving the norm in some dense subspaces of $h_\omega^\Phi(\mathbb{R}^n)$, from which, we further deduce that for a given admissible triplet $(\rho,\, q,\, s)_\omega$ and a β-quasi-Banach space $\mathcal{B}_\beta$ with $\beta \in (0, 1]$, if T is a $\mathcal{B}_\beta$-sublinear operator, and maps all $(\rho,\, q,\, s)_\omega$-atoms and $(\rho,\, q)_\omega$-single-atoms with $q < \infty$ (or all continuous $(\rho,\, q,\, s)_\omega$-atoms with $q = \infty$) into uniformly bounded elements of $\mathcal{B}_\beta$, then T uniquely extends to a bounded $\mathcal{B}_\beta$-sublinear operator from $h_\omega^\Phi(\mathbb{R}^n)$ to $\mathcal{B}_\beta$. As applications, we show that the local Riesz transforms are bounded on $h_\omega^\Phi(\mathbb{R}^n)$, the local fractional integrals are bounded from $h_{\omega^p}^p(\mathbb{R}^n)$ to $L_{\omega^q}^q(\mathbb{R}^n)$ when $q > 1$ and from $h_{\omega^p}^p(\mathbb{R}^n)$ to $h_{\omega^q}^q(\mathbb{R}^n)$ when $q \le 1$, and some pseudo-differential operators are also bounded on both $h_\omega^\Phi(\mathbb{R}^n)$. All results for any general Φ even when $\omega \equiv 1$ are new.

Acknowledgements. The first author is supported by the National Natural Science Foundation (Grant No. 10871025) of China and Program for Changjiang Scholars and Innovative Research Team in University of China.

Both authors would like to thank Professor Lin Tang for some helpful discussions on the subject of this paper and they would also like to thank the referee and the copy editor for their valuable remarks which made this article more readable.

2010 *Mathematics Subject Classification*: Primary 46E30; Secondary 42B35, 42B30, 42B25, 42B20, 35S05, 47G30, 47B06.

Key words and phrases: local weight, local Orlicz-Hardy space, atom, local grand maximal function, quasi-Banach space, BMO-type space, duality, local Riesz transform, local fractional integral, pseudo-differential operator.

1. Introduction

It is well known that the theory of the classical local Hardy spaces, originally introduced by Goldberg [18], plays an important role in partial differential equations and harmonic analysis; see, for example, [18, 6, 43, 51, 52, 53] and their references. In particular, pseudo-differential operators are bounded on local Hardy spaces $h^p(\mathbb{R}^n)$ with $p \in (0,1]$, but they are not bounded on Hardy spaces $H^p(\mathbb{R}^n)$ with $p \in (0,1]$; see [18] (also [52, 53]). In [6], Bui studied the weighted version $h^p_\omega(\mathbb{R}^n)$ of the local Hardy space $h^p(\mathbb{R}^n)$ with $\omega \in A_\infty(\mathbb{R}^n)$, where and in what follows, $A_q(\mathbb{R}^n)$ for $q \in [1,\infty]$ denotes the *class of Muckenhoup's weights*; see, for example, [17] for their definitions and properties.

Rychkov [43] introduced and studied a class of local weights, denoted by $A^{\mathrm{loc}}_\infty(\mathbb{R}^n)$ (see also Definition 2.1 below), and the weighted Besov-Lipschitz spaces and Triebel-Lizorkin spaces with weights belonging to $A^{\mathrm{loc}}_\infty(\mathbb{R}^n)$, which contains $A_\infty(\mathbb{R}^n)$ weights and exponential weights introduced by Schott [44] as special cases. In particular, Rychkov [43] generalized some of the theory of weighted local Hardy spaces developed by Bui [6] to $A^{\mathrm{loc}}_\infty(\mathbb{R}^n)$ weights. In fact, Rychkov established the local vertical and the local nontangential maximal function characterizations of weighted local Hardy spaces with $A^{\mathrm{loc}}_\infty(\mathbb{R}^n)$ weights. Very recently, Tang [49] established the weighted atomic decomposition characterization of the weighted local Hardy space $h^p_\omega(\mathbb{R}^n)$ with $\omega \in A^{\mathrm{loc}}_\infty(\mathbb{R}^n)$ via the local grand maximal function. Motivated by [5], Tang also established some criterions for boundedness of $\mathcal{B}_\beta$-sublinear operators on $h^p_\omega(\mathbb{R}^n)$ (see Section 6 for the notion of $\mathcal{B}_\beta$-sublinear operators, which was first introduced in [56]). As applications, Tang [49, 50] proved that some strongly singular integrals, pseudo-differential operators and their commutators are bounded on $h^p_\omega(\mathbb{R}^n)$. It is worth pointing out that in recent years, many papers are focused on criterions for boundedness of (sub)linear operators on various Hardy spaces with different underlying spaces (see, for example, [4, 35, 57, 5, 20, 56, 42, 49]), and on entropy and approximation numbers of embeddings of function spaces with Muckenhoupt weight (see, for example, [21, 22, 23, 24]).

It is also well known that the classical BMO space (the *space of functions with bounded mean oscillation*) originally introduced by John and Nirenberg [29] and the classical Morrey space originally introduced by Morrey [37] play an important role in the study of partial differential equations and harmonic analysis; see, for example, [15, 11, 14, 38]. In particular, Fefferman and Stein [15] proved that $\mathrm{BMO}(\mathbb{R}^n)$ is the dual space of the Hardy space $H^1(\mathbb{R}^n)$. Moveover, Goldberg [18] introduced the space $\mathrm{bmo}(\mathbb{R}^n)$ and proved that $\mathrm{bmo}(\mathbb{R}^n)$ is the dual space of the local Hardy space $h^1(\mathbb{R}^n)$.

On the other hand, as the generalization of $L^p(\mathbb{R}^n)$, the Orlicz space was introduced by Birnbaum-Orlicz in [2] and Orlicz in [39], since then, the theory of the Orlicz spaces themselves has been well developed and these spaces have been widely used in probability, statistics, potential theory, partial differential equations, as well as harmonic analysis and some other fields of analysis; see, for example, [40, 41, 7, 34, 25]. Moreover, Orlicz-Hardy spaces are also suitable substitutes of the Orlicz spaces in dealing with many problems of analysis; see, for example, [26, 55, 47, 27]. Recall that Orlicz-Hardy spaces and their dual spaces were studied by Janson [26] on $\mathbb{R}^n$ and Viviani [55] on spaces of homogeneous type in the sense of Coifman and Weiss [10]. Recently, Orlicz-Hardy spaces associated with some differential operators and their dual spaces were introduced and studied in [28, 27].

Let $\omega \in A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$, Φ be a concave function on $(0, \infty)$ of strictly lower type $p_\Phi \in (0, 1]$ (see (2.6) below for the definition) and

$$\rho(t) \equiv t^{-1}/\Phi^{-1}(t^{-1})$$

for all $t \in (0, \infty)$, where Φ^{-1} is the inverse function of Φ. A typical example of such Orlicz functions is $\Phi(t) \equiv t^p$ for all $t \in (0, \infty)$ and $p \in (0, 1]$. Motivated by [43, 49, 18, 28, 27, 5], in this paper, we introduce the weighted local Orlicz-Hardy space $h_\omega^\Phi(\mathbb{R}^n)$ via the local grand maximal function. We also introduce the BMO-type space $\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$ and establish the duality between $h_\omega^\Phi(\mathbb{R}^n)$ and $\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$. Characterizations of $h_\omega^\Phi(\mathbb{R}^n)$, including the atomic characterization, the local vertical and the local nontangential maximal function characterizations, are presented. Using the atomic characterization, we prove the existence of finite atomic decompositions achieving the norm in some dense subspaces of $h_\omega^\Phi(\mathbb{R}^n)$, from which, we further deduce that for a given admissible triplet $(\rho,\, q,\, s)_\omega$ and a β-quasi-Banach space $\mathcal{B}_\beta$ with $\beta \in (0, 1]$, if T is a $\mathcal{B}_\beta$-sublinear operator, and maps all $(\rho,\, q,\, s)_\omega$-atoms and $(\rho,\, q)_\omega$-single-atoms with $q < \infty$ (or all continuous $(\rho,\, q,\, s)_\omega$-atoms with $q = \infty$) into uniformly bounded elements of $\mathcal{B}_\beta$, then T uniquely extends to a bounded $\mathcal{B}_\beta$-sublinear operator from $h_\omega^\Phi(\mathbb{R}^n)$ to $\mathcal{B}_\beta$. As applications, we show that the local Riesz transforms are bounded on $h_\omega^\Phi(\mathbb{R}^n)$, the local fractional integrals are bounded from $h_{\omega^p}^p(\mathbb{R}^n)$ to $L_{\omega^q}^q(\mathbb{R}^n)$ when $q > 1$ and from $h_{\omega^p}^p(\mathbb{R}^n)$ to $h_{\omega^q}^q(\mathbb{R}^n)$ when $q \le 1$, and some pseudo-differential operators are also bounded on both $h_\omega^\Phi(\mathbb{R}^n)$ and $\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$. We point out that the Orlicz-Hardy spaces $h_\omega^\Phi(\mathbb{R}^n)$ include the classical local Hardy spaces of Goldberg [18], the weighted local Hardy spaces of Bui [6] and the weighted local Hardy spaces of Tang [49] as special cases. Moreover, the method of obtaining atomic decompositions used in this paper (see the proof of Theorem 5.6 below) is different from the classical methods in [18, 6]. Indeed, following Bownik [3] (see also [5, 49]), we give a direct proof for the weighted atomic characterization of $h_\omega^\Phi(\mathbb{R}^n)$, without invoking the atomic characterization of $H_\omega^\Phi(\mathbb{R}^n)$. All results of this paper for any general Φ even when $\omega \equiv 1$ are new.

Precisely, this paper is organized as follows. In Section 2, we first recall some definitions and notation concerning local weights introduced in [43, 49], then describe some basic assumptions and present some properties of Orlicz functions considered in this paper.

In Section 3, we first introduce the weighted local Orlicz-Hardy space $h_{\omega,\,N}^\Phi(\mathbb{R}^n)$ via the local grand maximal function, and then the weighted atomic local Orlicz-Hardy space $h_\omega^{\rho,\,q,\,s}(\mathbb{R}^n)$ for any admissible triplet $(\rho,\, q,\, s)_\omega$ (see Definition 3.4 below). We point out that when $\Phi(t) \equiv t^p$ for all $t \in (0, \infty)$ and $p \in (0, 1]$, the weighted local Orlicz-Hardy space $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$ is just the weighted local Hardy space $h^{p}_{\omega,\,N}(\mathbb{R}^n)$ introduced by Tang in [49]. Next, we establish the local vertical and the local nontangential maximal function characterizations of $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$ via a local Calderón reproducing formula and some useful estimates established by Rychkov [43], which generalizes [43, Theorem 2.24] by taking $\Phi(t) \equiv t^p$ for all $t \in (0,\infty)$ and $p \in (0,1]$; see Theorems 3.12 and 3.14 and Remark 3.13 below. Finally, we present some properties of these weighted local Orlicz-Hardy spaces $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$ and weighted atomic local Orlicz-Hardy spaces $h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$.

Throughout the whole paper, as usual, $\mathcal{D}(\mathbb{R}^n)$ denotes the *set of all* $C^\infty(\mathbb{R}^n)$ *functions on* $\mathbb{R}^n$ *with compact support*, endowed with the inductive limit topology, and $\mathcal{D}'(\mathbb{R}^n)$ its *topological dual space*, endowed with the weak $*$-topology. Let $\lfloor r \rfloor$ for any $r \in \mathbb{R}$ denote the *maximal integer not more than* r. In Section 4, for any given $f \in \mathcal{D}'(\mathbb{R}^n)$, integer

$$s \geq \lfloor n(q_\omega/p_\Phi - 1)\rfloor$$

and $\lambda > \inf_{x\in\mathbb{R}^n} \mathcal{G}_{N,\,\widetilde{R}}(f)(x)$, where q_ω, p_Φ and $\mathcal{G}_{N,\,\widetilde{R}}(f)$ are respectively as in (2.4), (2.6) and (3.2) below, and $\widetilde{R} = 2^{3(10+n)}$, following [46, 3, 5, 49], via a Whitney decomposition of Ω_λ in (4.1), we obtain the Calderón-Zygmund decomposition $f \equiv g + \sum_i b_i$ in $\mathcal{D}'(\mathbb{R}^n)$ of degree s and height λ associated with the local grand maximal function $\mathcal{G}_{N,\,\widetilde{R}}(f)$. The main task of Section 4 is to establish some subtle estimates for g and $\{b_i\}_i$. Precisely, Lemmas 4.2 through 4.5 are estimates on $\{b_i\}_i$, the bad part of f, while Lemmas 4.6 and 4.7 on g, the good part of f. As an application of these estimates, we obtain the density of $L^q_\omega(\mathbb{R}^n) \cap h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$ in $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$, where $q \in (q_\omega,\infty)$ (see Corollary 4.8 below). Differently from the proof of [49, Lemma 4.9], via an application of the boundedness of the local vector-valued Hardy-Littlewood maximal operator obtained by Rychkov [43] (see also Lemma 3.10 below), our Lemma 4.7 below improves [49, Lemma 4.9] even when $\Phi(t) \equiv t^p$ for all $t \in (0,\infty)$ and $p \in (0,1]$, which is necessary for Corollary 4.8.

In Section 5, we prove that for any given admissible triplet $(\rho,\,q,\,s)_\omega$,

$$h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n) = h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$$

with equivalent norms when positive integer $N \geq N_{\Phi,\,\omega}$ (see (3.25) below for the definition of $N_{\Phi,\,\omega}$), by using the Calderón-Zygmund decomposition and some subtle estimates obtained in Section 4, which completely covers [49, Theorem 5.1] by taking $\Phi(t) \equiv t^p$ for all $p \in (0,1]$ and $t \in (0,\infty)$; see Theorem 5.6 and Remark 5.7 below. It is worth pointing out that we show Theorem 5.6 by a way different from the methods in [18, 6], but close to those in [3, 5, 49]. For simplicity, in the rest of this introduction, we denote by $h^{\Phi}_{\omega}(\mathbb{R}^n)$ the *weighted local Orlicz-Hardy space* $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$ *with* $N \geq N_{\Phi,\,\omega}$.

Assume that $(\rho,\,q,\,s)_\omega$ is an admissible triplet. Let $h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$ be the *space of all finite linear combinations of* $(\rho,\,q,\,s)_\omega$*-atoms and* $(\rho,\,q)_\omega$*-single-atoms* (see Definition 6.1 below), and $h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin},\,c}(\mathbb{R}^n)$ the *space of all* $f \in h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$ *with compact support.* In Section 6, we prove that $\|\cdot\|_{h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)}$ and $\|\cdot\|_{h^{\Phi}_{\omega}(\mathbb{R}^n)}$ are equivalent quasi-norms on $h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$ when $q < \infty$, and $\|\cdot\|_{h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)}$ and $\|\cdot\|_{h^{\Phi}_{\omega}(\mathbb{R}^n)}$ are equivalent quasi-norms on $h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin},\,c}(\mathbb{R}^n) \cap C(\mathbb{R}^n)$ when $q = \infty$ (see Theorem 6.2 below). As an application, we prove that for a given admissible triplet $(\rho,\,q,\,s)_\omega$ and a β-quasi-Banach space $\mathcal{B}_\beta$ with $\beta \in (0,1]$, if Φ has a upper type $\widetilde{p}$ satisfying $0 < \widetilde{p} \leq \beta$, and T is a $\mathcal{B}_\beta$-sublinear operator mapping all $(\rho,\,q,\,s)_\omega$-atoms and $(\rho,\,q)_\omega$-single-atoms with $q \in (q_\omega,\infty)$ (or all continuous

$(\rho,\, q,\, s)_\omega$-atoms with $q=\infty$) into uniformly bounded elements of $\mathcal{B}_\beta$, then T uniquely extends to a bounded $\mathcal{B}_\beta$-sublinear operator from $h^\Phi_\omega(\mathbb{R}^n)$ to $\mathcal{B}_\beta$ which coincides with T on these $(\rho,\, q,\, s)_\omega$-atoms and $(\rho,\, q)_\omega$-single-atoms; see Theorem 6.4 below. We remark that this extends both the results of Meda-Sjögren-Vallarino [35] and Yang-Zhou [57] to the setting of weighted local Orlicz-Hardy spaces. We also point out that Theorem 6.2(i) and Theorem 6.4(i) below completely cover [49, Theorem 6.1] and [49, Theorem 6.2], respectively, by taking $\Phi(t)\equiv t^p$ for all $t\in(0,\infty)$ and $p\in(0,1]$; and Theorem 6.2(ii) and Theorem 6.4(ii) below are new even for $\Phi(t)\equiv t^p$ with $t\in(0,\infty)$ and $p\in(0,1]$; see Remark 6.5 below.

Let $(\rho,\, q,\, s)_\omega$ be an admissible triplet, q' the dual exponent of q and q_ω the critical index of ω. In Section 7, we introduce the BMO-type space $\mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$ and prove that

$$[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*=\mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n),$$

where $[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*$ denotes the dual space of $h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)$; see Theorem 7.5 below. Denote $\mathrm{bmo}^{1}_{\rho,\,\omega}(\mathbb{R}^n)$ simply by $\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$. As applications of Theorems 5.6 and 7.5, we see that for $q\in[1,\frac{q_\omega}{q_\omega-1})$, $\mathrm{bmo}^{q}_{\rho,\,\omega}(\mathbb{R}^n)=\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$ with equivalent norms and

$$\left[h^\Phi_\omega(\mathbb{R}^n)\right]^*=\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n);$$

see Corollaries 7.6 and 7.7 below.

In Section 8, as applications of Theorem 6.2, we obtain the boundedness of some operators from $h^\Phi_\omega(\mathbb{R}^n)$ to some β-quasi-Banach space $\mathcal{B}_\beta$ with $\beta\in(0,1]$. First, we prove that the local Riesz transforms are bounded on $h^\Phi_\omega(\mathbb{R}^n)$ if $p_\Phi=p^+_\Phi$ and (2.5) holds for p^+_Φ with $t\in[1,\infty)$ (see Section 2 below for the definitions of p^+_Φ), which completely covers [49, Lemma 8.3] by taking $\Phi(t)\equiv t$ for all $t\in(0,\infty)$; see Theorem 8.2 and Remark 8.4 below. Then we introduce the local fractional integral operator I^{loc}_α and show that I^{loc}_α is bounded from $h^p_{\omega^p}(\mathbb{R}^n)$ to $L^q_{\omega^q}(\mathbb{R}^n)$ when $\alpha\in(0,n)$, $p\in[\frac{n}{n+\alpha},1]$, $\frac{1}{q}=\frac{1}{p}-\frac{\alpha}{n}$, and $\omega^{\frac{nr}{nr-n-r\alpha}}\in A^{\mathrm{loc}}_1(\mathbb{R}^n)$ for some $r\in(\frac{n}{n-\alpha},\infty)$ and $\int_{\mathbb{R}^n}[\omega(x)]^p\,dx=\infty$ (see Theorem 8.10 below); furthermore, when $\alpha\in(0,1)$, $p\in(0,\frac{n}{n+\alpha}]$, $\frac{1}{q}=\frac{1}{p}-\frac{\alpha}{n}$, and ω satisfies the same conditions, we prove that I^{loc}_α is bounded from $h^p_{\omega^p}(\mathbb{R}^n)$ to $h^q_{\omega^q}(\mathbb{R}^n)$ (see Theorem 8.11 below). To the best of our knowledge, Theorems 8.10 and 8.11 are new even when $\omega\equiv1$. Finally, let $\omega\in A_\infty(\phi)$ which was introduced by Tang [50] (see also Definition 8.13 below) and T be an $S^0_{1,\,0}(\mathbb{R}^n)$ pseudo-differential operator. We prove that T is bounded on $h^\Phi_\omega(\mathbb{R}^n)$ if $p_\Phi=p^+_\Phi$ and (2.5) holds for p^+_Φ with $t\in[1,\infty)$; see Theorem 8.18 below. We point out that $A_\infty(\phi)\subset A^{\mathrm{loc}}_\infty(\mathbb{R}^n)$ but $A_\infty(\phi)\supset A_\infty(\mathbb{R}^n)$. We also remark that Theorem 8.18 below extends [18, Theorem 4] to the setting of weighted local Orlicz-Hardy spaces, and completely covers [49, Theorem 7.3] by taking $\Phi(t)\equiv t^p$ for all $t\in(0,\infty)$ and $p\in(0,1]$ and also [32, Theorem 2] by taking $\Phi(t)\equiv t$ for all $t\in(0,\infty)$ and $\omega\in A_1(\mathbb{R}^n)$; see Remark 8.19 below. As an application of Theorems 7.5 and 8.18, we also obtain that T is bounded on $\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$; see Corollary 8.20 below.

A main motivation of this paper is to pave the way for the study of weighted Orlicz-Hardy spaces associated with divergence operators on strongly Lipschitz domains of $\mathbb{R}^n$. The corresponding Hardy spaces associated with divergence operators on strongly Lipschitz domains of $\mathbb{R}^n$ were originally studied by Auscher and Russ [1], where the atomic characterization of the classical Hardy spaces plays a key role. Earlier works on Hardy spaces on domains can be found, for example, in [31, 36, 9, 8, 54]. It was shown in these papers that the theory of Hardy spaces on domains plays an important role in partial differential equations and harmonic analysis.

Finally we make some conventions on notation. Throughout the whole paper, we denote by C *a positive constant* which is independent of the main parameters, but it may vary from line to line. We also use $C(\gamma, \beta, \cdots)$ to denote *a positive constant depending on the indicated parameters* γ, β, $\cdots$. The *symbol* $A \lesssim B$ means that $A \le CB$. If $A \lesssim B$ and $B \lesssim A$, then we write $A \sim B$. The *symbol* $\lfloor s \rfloor$ for $s \in \mathbb{R}$ denotes the maximal integer not more than s. For any given normed spaces $\mathcal{A}$ and $\mathcal{B}$ with the corresponding norms $\|\cdot\|_{\mathcal{A}}$ and $\|\cdot\|_{\mathcal{B}}$, the *symbol* $\mathcal{A} \subset \mathcal{B}$ means that for all $f \in \mathcal{A}$, then $f \in \mathcal{B}$ and $\|f\|_{\mathcal{B}} \lesssim \|f\|_{\mathcal{A}}$. For any subset G of $\mathbb{R}^n$, we denote by $G^{\complement}$ the *set* $\mathbb{R}^n \setminus G$; for a measurable set E, denote by χ_E the *characteristic function of* E. We also set $\mathbb{N} \equiv \{1, 2, \cdots\}$ and $\mathbb{Z}_+ \equiv \mathbb{N} \cup \{0\}$. For any $\theta \equiv (\theta_1, \dots, \theta_n) \in \mathbb{Z}_+^n$, let $|\theta| \equiv \theta_1 + \cdots + \theta_n$ and $\partial_x^\theta \equiv \frac{\partial^{|\theta|}}{\partial x_1^{\theta_1} \cdots \partial x_n^{\theta_n}}$. Given a function g on $\mathbb{R}^n$, if $\int_{\mathbb{R}^n} g(x)\,dx \ne 0$, let $L_g \equiv -1$; otherwise, let $L_g \in \mathbb{Z}_+$ be the *maximal integer* such that g has vanishing moments up to order L_g, namely, $\int_{\mathbb{R}^n} g(x) x^\alpha\,dx = 0$ for all multi-indices α with $|\alpha| \le L_g$.

2. Preliminaries

In this section, we first recall some notions and notation concerning local weights introduced in [43, 49], then describe some basic assumptions and present some properties of Orlicz functions considered in this paper.

2.1. $A_p^{\mathrm{loc}}(\mathbb{R}^n)$ weights. In this subsection, we recall some notions and properties of the local weights. Let Q be a cube in $\mathbb{R}^n$ and we denote its Lebesgue measure by $|Q|$. Throughout the whole paper, *all cubes are assumed to be closed and their sides parallel to the coordinate axes.*

Definition 2.1. Let $p \in (1, \infty)$. The *weight class* $A_p^{\mathrm{loc}}(\mathbb{R}^n)$ is defined to be the set of all nonnegative locally integrable functions ω on $\mathbb{R}^n$ such that

$$A_p^{\mathrm{loc}}(\omega) \equiv \sup_{|Q|\le 1} \frac{1}{|Q|^p} \int_Q \omega(x)\,dx \left(\int_Q [\omega(y)]^{-p'/p}\,dy \right)^{p/p'} < \infty, \tag{2.1}$$

where the supremum is taken over all cubes $Q \subset \mathbb{R}^n$ with $|Q| \le 1$ and $\frac{1}{p} + \frac{1}{p'} = 1$.

When $p = 1$, the *weight class* $A_1^{\mathrm{loc}}(\mathbb{R}^n)$ is defined to be the set of all nonnegative locally integrable functions ω on $\mathbb{R}^n$ such that

$$A_1^{\mathrm{loc}}(\omega) \equiv \sup_{|Q|\le 1} \frac{1}{|Q|} \int_Q \omega(x)\,dx \left(\operatorname*{esssup}_{y \in Q} [\omega(y)]^{-1} \right) < \infty, \tag{2.2}$$

where the supremum is taken over all cubes $Q \subset \mathbb{R}^n$ with $|Q| \le 1$.

When $p = \infty$, the *weight class* $A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$ is defined to be the set of all nonnegative locally integrable functions ω on $\mathbb{R}^n$ such that for any $\alpha \in (0,1)$,

$$A_{\infty}^{\mathrm{loc}}(\omega;\,\alpha) \equiv \sup_{|Q|\le 1}\left[\sup_{F\subset Q,|F|\ge\alpha|Q|}\frac{\omega(Q)}{\omega(F)}\right] < \infty, \tag{2.3}$$

where F runs through all measurable sets in $\mathbb{R}^n$ with the indicated properties, the supremum is taken over all cubes $Q \subset \mathbb{R}^n$ with $|Q| \le 1$ and $\omega(Q) \equiv \int_Q \omega(x)\,dx$.

REMARK 2.2. (i) We point out that the weight class $A_p^{\mathrm{loc}}(\mathbb{R}^n)$ for $p \in (1,\infty]$ was introduced by Rychkov [43] and $A_1^{\mathrm{loc}}(\mathbb{R}^n)$ by Tang [49]. By Hölder's inequality, we see that $A_{p_1}^{\mathrm{loc}}(\mathbb{R}^n) \subset A_{p_2}^{\mathrm{loc}}(\mathbb{R}^n) \subset A_{\infty}^{\mathrm{loc}}(\mathbb{R}^n)$, if $1 \le p_1 < p_2 < \infty$. Conversely, it was proved in [43, Lemma 1.3] that if $\omega \in A_{\infty}^{\mathrm{loc}}(\mathbb{R}^n)$, then $\omega \in A_p^{\mathrm{loc}}(\mathbb{R}^n)$ for some $p \in (1,\infty)$. Thus, we have $A_{\infty}^{\mathrm{loc}}(\mathbb{R}^n) = \cup_{1\le p<\infty} A_p^{\mathrm{loc}}(\mathbb{R}^n)$.

(ii) For any constant $\widetilde{C} \in (0,\infty)$, the condition $|Q| \le 1$ can be replaced by $|Q| \le \widetilde{C}$ in (2.1), (2.2) and (2.3); see [43, Remark 1.5]. In this case, $A_p^{\mathrm{loc}}(\omega)$ with $p \in [1,\infty)$ and $A_{\infty}^{\mathrm{loc}}(\omega,\,\alpha)$ depend on $\widetilde{C}$.

In what follows, $Q(x,t)$ denotes the *closed cube centered at* x *and of the sidelength* t. Similarly, given $Q = Q(x,t)$ and $\lambda \in (0,\infty)$, we write λQ for the λ-*dilated cube*, which is the cube with the same center x and with sidelength λt. Given a Lebesgue measurable set E and a weight $\omega \in A_{\infty}^{\mathrm{loc}}(\mathbb{R}^n)$, let $\omega(E) \equiv \int_E \omega(x)\,dx$. For any $\omega \in A_{\infty}^{\mathrm{loc}}(\mathbb{R}^n)$, the *space* $L_\omega^p(\mathbb{R}^n)$ with $p \in (0,\infty)$ denotes the set of all measurable functions f such that

$$\|f\|_{L_\omega^p(\mathbb{R}^n)} \equiv \left\{\int_{\mathbb{R}^n} |f(x)|^p \omega(x)\,dx\right\}^{1/p} < \infty,$$

and $L_\omega^\infty(\mathbb{R}^n) \equiv L^\infty(\mathbb{R}^n)$. The *space* $L_\omega^{1,\,\infty}(\mathbb{R}^n)$ denotes the set of all measurable functions f such that

$$\|f\|_{L_\omega^{1,\,\infty}(\mathbb{R}^n)} \equiv \sup_{\lambda>0}\{\lambda\omega(\{x \in \mathbb{R}^n : |f(x)| > \lambda\})\} < \infty.$$

For a positive constant $\widetilde{C}$, any locally integrable function f and $x \in \mathbb{R}^n$, the *local Hardy-Littlewood maximal function* $M_{\widetilde{C}}^{\mathrm{loc}}(f)$ is defined by

$$M_{\widetilde{C}}^{\mathrm{loc}}(f)(x) \equiv \sup_{Q\ni x,\,|Q|\le\widetilde{C}} \frac{1}{|Q|}\int_Q |f(y)|\,dy,$$

where the supremum is taken over all cubes $Q \subset \mathbb{R}^n$ such that $Q \ni x$ and $|Q| \le \widetilde{C}$. If $\widetilde{C} = 1$, we denote $M_{\widetilde{C}}^{\mathrm{loc}}(f)$ simply by $M^{\mathrm{loc}}(f)$.

Next, we recall some properties for weights in $A_{\infty}^{\mathrm{loc}}(\mathbb{R}^n)$ and $A_p(\mathbb{R}^n)$, where and in what follows, $A_p(\mathbb{R}^n)$ for $p \in [1,\infty)$ denotes the classical *Muckenhoupt weights*; see [17, 46] for their definitions.

LEMMA 2.3. (i) *Let* $p \in [1,\infty)$, $\omega \in A_p^{\mathrm{loc}}(\mathbb{R}^n)$, *and* Q *be a unit cube, namely,* $l(Q) = 1$. *Then there exist an* $\overline{\omega} \in A_p(\mathbb{R}^n)$ *such that* $\overline{\omega} = \omega$ *on* Q, *and a positive constant* C *independent of* Q *such that* $A_p(\overline{\omega}) \le C A_p^{\mathrm{loc}}(\omega)$, *where* $A_p(\overline{\omega})$ *denotes the weight constant of* $\overline{\omega}$, *which is as in* (2.1) *and* (2.2) *by removing the restriction* $l(Q) \le 1$.

(ii) *If* $\omega \in A_p^{\mathrm{loc}}(\mathbb{R}^n)$ *with* $p \in (1,\infty)$, *then there exist* $\eta_1, \eta_2 \in (0,\infty)$ *such that* $\omega \in A_{p-\eta_1}^{\mathrm{loc}}(\mathbb{R}^n)$ *with* $p - \eta_1 \in (1,\infty)$, *and* $\omega^{1+\eta_2} \in A_p^{\mathrm{loc}}(\mathbb{R}^n)$.

(iii) *If* $1 \le p_1 < p_2 < \infty$, *then* $A_{p_1}^{\mathrm{loc}}(\mathbb{R}^n) \subset A_{p_2}^{\mathrm{loc}}(\mathbb{R}^n)$.

(iv) *For* $p \in (1, \infty)$, $\omega \in A_p^{\mathrm{loc}}(\mathbb{R}^n)$ *if and only if* $\omega^{-1/(p-1)} \in A_{p'}^{\mathrm{loc}}(\mathbb{R}^n)$, *where*

$$1/p + 1/p' = 1.$$

(v) *For* $\omega \in A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$ *and* $Q = Q(x_0, l(Q))$, *there exists a positive constant* C *such that* $\omega(2Q) \le C\omega(Q)$ *when* $l(Q) < 1$, *and* $\omega(Q(x_0, l(Q) + 1)) \le C\omega(Q)$ *when* $l(Q) \ge 1$.

(vi) *If* $p \in (1, \infty)$ *and* $\omega \in A_p^{\mathrm{loc}}(\mathbb{R}^n)$, *then the local Hardy-Littlewood maximal operator* M^{loc} *is bounded on* $L^p_\omega(\mathbb{R}^n)$.

(vii) *If* $\omega \in A_1^{\mathrm{loc}}(\mathbb{R}^n)$, *then* M^{loc} *is bounded from* $L^1_\omega(\mathbb{R}^n)$ *to* $L^{1,\infty}_\omega(\mathbb{R}^n)$.

(viii) *If* $\omega \in A_p(\mathbb{R}^n)$ *with* $p \in [1, \infty)$, *then there exists a positive constant* C *such that for all cubes* Q_1, $Q_2 \subset \mathbb{R}^n$ *with* $Q_1 \subset Q_2$,

$$\frac{\omega(Q_2)}{\omega(Q_1)} \le C \left(\frac{|Q_2|}{|Q_1|} \right)^p .$$

Lemma 2.3(i) is just [43, Lemma 1.1]. The statements (ii) through (vii) of Lemma 2.3 are just Lemma 2.1 and Corollary 2.1 in [49], which are deduced from Lemma 2.3(i) and the properties of $A_p(\mathbb{R}^n)$; see the proofs of [49, Lemma 2.1, Corollary 2.1]. Lemma 2.3(viii) is included, for example, in [16, 17, 46].

REMARK 2.4. Let $\widetilde{C}$ be a positive constant. It was pointed out in [43, Remark 1.5] and [49] that (i) through (vii) of Lemma 2.3 are also true, if $l(Q) = 1$, $l(Q) \ge 1$, $l(Q) < 1$, $Q(x_0, l(Q) + 1)$ and M^{loc} are respectively replaced by $l(Q) = \widetilde{C}$, $l(Q) \ge \widetilde{C}$, $l(Q) < \widetilde{C}$, $Q(x_0, l(Q) + \widetilde{C})$ and $M^{\mathrm{loc}}_{\widetilde{C}}$. In this case, the constants appearing in (i), (vi) and (vii) of Lemma 2.3 depend on $\widetilde{C}$.

For any given $\omega \in A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$, define the *critical index of* ω by

$$q_\omega \equiv \inf \left\{ p \in [1, \infty) : \omega \in A_p^{\mathrm{loc}}(\mathbb{R}^n) \right\}. \tag{2.4}$$

REMARK 2.5. Obviously, $q_\omega \in [1, \infty)$. If $q_\omega \in (1, \infty)$, by Lemma 2.2(ii), it is easy to know that $\omega \notin A_{q_\omega}^{\mathrm{loc}}(\mathbb{R}^n)$. Moreover, there exists an $\omega \notin A_1^{\mathrm{loc}}(\mathbb{R}^n)$ such that $q_\omega = 1$. Indeed, for $t \in \mathbb{R} \setminus \{0\}$, let $\omega(t) \equiv [\ln(\frac{1}{|t|})]^{-1}$. Johnson and Neugebauer [30, p. 254, Remark] showed that $\omega \in (\cap_{p>1} A_p(\mathbb{R}^n)) \setminus A_1(\mathbb{R}^n)$. By the fact that $A_p(\mathbb{R}^n) \subset A_p^{\mathrm{loc}}(\mathbb{R}^n)$ for all $p \in [1, \infty)$, which is obvious by their definitions, we see that $\omega \in \cap_{p>1} A_p^{\mathrm{loc}}(\mathbb{R}^n)$. We claim that $\omega \notin A_1^{\mathrm{loc}}(\mathbb{R}^n)$. In fact, taking $x \in (0, 1/2)$, then we have

$$M^{\mathrm{loc}}(\omega)(x) \ge \frac{1}{2} \int_{x-1}^{x+1} \omega(t)\, dt \ge \int_0^{1/2} \left[\ln\left(\frac{1}{t}\right) \right]^{-1} dt \equiv \infty.$$

Moreover, it is easy to see that $\omega(x) \to 0$ as $x \to 0$. Thus, by (2.2), we know that $\omega \notin A_1^{\mathrm{loc}}(\mathbb{R}^n)$.

For $\mathcal{D}(\mathbb{R}^n)$, $\mathcal{D}'(\mathbb{R}^n)$ and $L^q_\omega(\mathbb{R}^n)$, we have the following conclusions.

LEMMA 2.6. *Let* $\omega \in A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$, q_ω *be as in* (2.4) *and* $p \in (q_\omega, \infty]$.

(i) *If* $\frac{1}{p} + \frac{1}{p'} = 1$, *then* $\mathcal{D}(\mathbb{R}^n) \subset L^{p'}_{\omega^{-1/(p-1)}}(\mathbb{R}^n)$.

(ii) $L^p_\omega(\mathbb{R}^n) \subset \mathcal{D}'(\mathbb{R}^n)$ *and the inclusion is continuous.*

(iii) *Let* $\phi \in \mathcal{D}(\mathbb{R}^n)$ *and* $\int_{\mathbb{R}^n} \phi(x)\, dx = 1$. *If* $q \in (q_\omega, \infty)$, *then for any* $f \in L^q_\omega(\mathbb{R}^n)$, $f * \phi_t \to f$ *in* $L^q_\omega(\mathbb{R}^n)$ *as* $t \to 0$, *where and in what follows,* $\phi_t(x) \equiv \frac{1}{t^n}\phi(\frac{x}{t})$ *for all* $t \in (0, \infty)$ *and* $x \in \mathbb{R}^n$.

We remark that (i) and (ii) of Lemma 2.6, and Lemma 2.6(iii) are, respectively, Lemma 2.2 and Proposition 2.1 in [49].

2.2. Orlicz functions. Let Φ be a positive function on $\mathbb{R}_+ \equiv (0, \infty)$. The function Φ is said to be of *upper type p (resp. lower type p)* for some $p \in [0, \infty)$, if there exists a positive constant C such that for all $t \in [1, \infty)$ (resp. $t \in (0, 1]$) and $s \in (0, \infty)$,

$$\Phi(st) \le C t^p \Phi(s). \tag{2.5}$$

Obviously, if Φ is of lower type p for some $p \in (0, \infty)$, then $\lim_{t \to 0_+} \Phi(t) = 0$. So for the sake of convenience, if it is necessary, we may assume that $\Phi(0) = 0$. If Φ is of both upper type p_1 and lower type p_0, then Φ is said to be of *type* (p_0, p_1). Let

$$\begin{aligned} p_\Phi^+ \equiv \inf\{p \in (0, \infty) :\ &\text{there exists } C \in (0, \infty) \\ &\text{such that (2.5) holds for all } t \in [1, \infty) \text{ and } s \in (0, \infty)\}, \end{aligned}$$

and

$$\begin{aligned} p_\Phi^- \equiv \sup\{p \in (0, \infty) :\ &\text{there exists } C \in (0, \infty) \\ &\text{such that (2.5) holds for all } t \in (0, 1] \text{ and } s \in (0, \infty)\}. \end{aligned}$$

The function Φ is said to be of *strictly lower type p* if for all $t \in (0, 1)$ and $s \in (0, \infty)$, $\Phi(st) \le t^p \Phi(s)$, and we define

$$p_\Phi \equiv \sup\{p \in (0, \infty) :\ \Phi(st) \le t^p \Phi(s) \text{ holds for all } t \in (0, 1) \text{ and } s \in (0, \infty)\}. \tag{2.6}$$

It is easy to see that $p_\Phi \le p_\Phi^- \le p_\Phi^+$ for all Φ. In what follows, p_Φ, p_Φ^- and p_Φ^+ are respectively called the *strictly critical lower type index*, the *critical lower type index* and the *critical upper type index* of Φ. We point out that if p_Φ is defined as in (2.6), then Φ is also of strictly critical lower type p_Φ; see [27] for the proof.

Throughout the whole paper, we always assume that Φ satisfies the following assumption.

Assumption (A). Let Φ be a positive function defined on $\mathbb{R}_+$, which is of strictly lower type and its strictly critical lower type index $p_\Phi \in (0, 1]$. Also assume that Φ is continuous, strictly increasing, subadditive and concave.

Notice that if Φ satisfies Assumption (A), then $\Phi(0) = 0$ and Φ is obviously of upper type 1. For any concave and positive function $\widetilde{\Phi}$ of strictly lower type p, if we set $\Phi(t) \equiv \int_0^t \frac{\Phi(s)}{s}\, ds$ for $t \in [0, \infty)$, then by [55, Proposition 3.1], Φ is equivalent to $\widetilde{\Phi}$, namely, there exists a positive constant C such that $C^{-1}\widetilde{\Phi}(t) \le \Phi(t) \le C\widetilde{\Phi}(t)$ for all $t \in [0, \infty)$; moreover, Φ is strictly increasing, concave, subadditive and continuous function of strictly lower type p. Notice that all our results are invariant on equivalent functions satisfying Assumption (A). From this, we deduce that all results in this paper with Φ as in Assumption (A) also holds for all concave and positive functions $\widetilde{\Phi}$ of the same strictly critical lower type p_Φ as Φ.

Let Φ satisfy Assumption (A) and $\omega \in A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$. A measurable function f on $\mathbb{R}^n$ is called to belong to the *space* $L_\omega^\Phi(\mathbb{R}^n)$ if $\int_{\mathbb{R}^n} \Phi(|f(x)|)\omega(x)\, dx < \infty$. Moreover, for any

$f \in L^{\Phi}_{\omega}(\mathbb{R}^n)$, define

$$\|f\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)} \equiv \inf\left\{\lambda \in (0,\infty) : \int_{\mathbb{R}^n} \Phi\left(\frac{|f(x)|}{\lambda}\right) \omega(x)\,dx \le 1\right\}.$$

Since Φ is strictly increasing, we define the *function* ρ on $\mathbb{R}_+$ by setting, for all $t \in (0,\infty)$,

$$\rho(t) \equiv \frac{t^{-1}}{\Phi^{-1}(t^{-1})}, \tag{2.7}$$

where Φ^{-1} is the inverse function of Φ. Then the types of Φ and ρ have the following relation. Let $0 < p_0 \le p_1 \le 1$ and Φ be an increasing function, then Φ is of type (p_0, p_1) if and only if ρ is of type $(p_1^{-1} - 1, p_0^{-1} - 1)$; see [55] for its proof. Moreover, it is easy to see that for all $t \in (0,\infty)$,

$$t\Phi\left(\frac{1}{t\rho(t)}\right) = 1, \tag{2.8}$$

which is used in what follows.

3. Weighted local Orlicz-Hardy spaces and their maximal function characterizations

In this section, we introduce the weighted local Orlicz-Hardy space $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$ via the local grand maximal function and establish its local vertical and nontangential maximal function characterizations via a local Calderón reproducing formula and some useful estimates obtained by Rychkov [43]. We also introduce the weighted atomic local Orlicz-Hardy space $h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$ and give some basic properties of these spaces.

First, we introduce some local maximal functions. For $N \in \mathbb{Z}_+$ and $R \in (0,\infty)$, let

$$\mathcal{D}_{N,\,R}(\mathbb{R}^n) \equiv \left\{ \psi \in \mathcal{D}(\mathbb{R}^n) :\ \operatorname{supp}(\psi) \subset B(0,R), \right.$$
$$\left. \|\psi\|_{\mathcal{D}_N(\mathbb{R}^n)} \equiv \sup_{x\in\mathbb{R}^n} \sup_{\alpha\in\mathbb{Z}^n_+,\,|\alpha|\le N} |\partial^\alpha \psi(x)| \le 1 \right\}.$$

Definition 3.1. Let $N \in \mathbb{Z}_+$ and $R \in (0,\infty)$. For any $f \in \mathcal{D}'(\mathbb{R}^n)$, the *local nontangential grand maximal function* $\widetilde{\mathcal{G}}_{N,\,R}(f)$ of f is defined by setting, for all $x \in \mathbb{R}^n$,

$$\widetilde{\mathcal{G}}_{N,\,R}(f)(x) \equiv \sup\left\{|\psi_t * f(z)| :\ |x-z| < t < 1,\ \psi \in \mathcal{D}_{N,\,R}(\mathbb{R}^n)\right\}, \tag{3.1}$$

and the *local vertical grand maximal function* $\mathcal{G}_{N,\,R}(f)$ of f is defined by setting, for all $x \in \mathbb{R}^n$,

$$\mathcal{G}_{N,\,R}(f)(x) \equiv \sup\left\{|\psi_t * f(x)| :\ t \in (0,1),\ \psi \in \mathcal{D}_{N,\,R}(\mathbb{R}^n)\right\}. \tag{3.2}$$

For convenience's sake, when $R = 1$, we denote $\mathcal{D}_{N,\,R}(\mathbb{R}^n)$, $\widetilde{\mathcal{G}}_{N,\,R}(f)$ and $\mathcal{G}_{N,\,R}(f)$ simply by $\mathcal{D}^0_N(\mathbb{R}^n)$, $\widetilde{\mathcal{G}}^0_N(f)$ and $\mathcal{G}^0_N(f)$, respectively; when $R = 2^{3(10+n)}$, we denote $\mathcal{D}_{N,\,R}(\mathbb{R}^n)$, $\widetilde{\mathcal{G}}_{N,\,R}(f)$ and $\mathcal{G}_{N,\,R}(f)$ simply by $\mathcal{D}_N(\mathbb{R}^n)$, $\widetilde{\mathcal{G}}_N(f)$ and $\mathcal{G}_N(f)$, respectively. For any $N \in \mathbb{Z}_+$ and $x \in \mathbb{R}^n$, obviously,

$$\mathcal{G}^0_N(f)(x) \le \mathcal{G}_N(f)(x) \le \widetilde{\mathcal{G}}_N(f)(x).$$

For the local grand maximal function $\mathcal{G}_N^0(f)$, we have the following Proposition 3.2, which is just [49, Proposition 2.2].

Proposition 3.2. *Let* $N \geq 2$.

(i) *Then there exists a positive constant* C *such that for all* $f \in (L^1_{\mathrm{loc}}(\mathbb{R}^n) \cap \mathcal{D}'(\mathbb{R}^n))$ *and almost every* $x \in \mathbb{R}^n$,

$$|f(x)| \leq \mathcal{G}_N^0(f)(x) \leq M^{\mathrm{loc}}(f)(x).$$

(ii) *If* $\omega \in A_p^{\mathrm{loc}}(\mathbb{R}^n)$ *with* $p \in (1, \infty)$, *then* $f \in L^p_\omega(\mathbb{R}^n)$ *if and only if* $f \in \mathcal{D}'(\mathbb{R}^n)$ *and* $\mathcal{G}_N^0(f) \in L^p_\omega(\mathbb{R}^n)$; *moreover,*

$$\|f\|_{L^p_\omega(\mathbb{R}^n)} \sim \|\mathcal{G}_N^0(f)\|_{L^p_\omega(\mathbb{R}^n)}.$$

(iii) *If* $\omega \in A_1^{\mathrm{loc}}(\mathbb{R}^n)$, *then* $\mathcal{G}_N^0$ *is bounded from* $L^1_\omega(\mathbb{R}^n)$ *to* $L^{1,\infty}_\omega(\mathbb{R}^n)$.

Now we introduce the weighted local Orlicz-Hardy space via the local grand maximal function as follows.

Definition 3.3. Let Φ satisfy Assumption (A), $\omega \in A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$, q_ω and p_Φ be respectively as in (2.4) and (2.6), and $\widetilde{N}_{\Phi,\,\omega} \equiv \lfloor n(\frac{q_\omega}{p_\Phi} - 1)\rfloor + 2$. For each $N \in \mathbb{N}$ with $N \geq \widetilde{N}_{\Phi,\,\omega}$, the *weighted local Orlicz-Hardy space* is defined by

$$h^\Phi_{\omega,\,N}(\mathbb{R}^n) \equiv \left\{f \in \mathcal{D}'(\mathbb{R}^n):\ \mathcal{G}_N(f) \in L^\Phi_\omega(\mathbb{R}^n)\right\}.$$

Moreover, let $\|f\|_{h^\Phi_{\omega,\,N}(\mathbb{R}^n)} \equiv \|\mathcal{G}_N(f)\|_{L^\Phi_\omega(\mathbb{R}^n)}$.

We remark that when $\Phi(t) \equiv t^p$ for all $t \in (0, \infty)$ and $p \in (0, 1]$, $h^\Phi_{\omega,\,N}(\mathbb{R}^n)$ above is the weighted local Hardy space $h^p_{\omega,\,N}(\mathbb{R}^n)$ introduced by Tang [49]. Obviously, for any integers N_1 and N_2 with $N_1 \geq N_2 \geq \widetilde{N}_{\Phi,\,\omega}$,

$$h^\Phi_{\omega,\,\widetilde{N}_{\Phi,\,\omega}}(\mathbb{R}^n) \subset h^\Phi_{\omega,\,N_2}(\mathbb{R}^n) \subset h^\Phi_{\omega,\,N_1}(\mathbb{R}^n),$$

and the inclusions are continuous. We also point out that Theorem 3.14 below further implies that

$$\|\mathcal{G}_N^0(f)\|_{L^\Phi_\omega(\mathbb{R}^n)} \sim \|\widetilde{\mathcal{G}}_N^0(f)\|_{L^\Phi_\omega(\mathbb{R}^n)} \sim \|\mathcal{G}_N(f)\|_{L^\Phi_\omega(\mathbb{R}^n)} \sim \|\widetilde{\mathcal{G}}_N(f)\|_{L^\Phi_\omega(\mathbb{R}^n)}$$

for all $N \in \mathbb{N}$ with $N \geq N_{\Phi,\,\omega}$ (see (3.25) for the definition of $N_{\Phi,\,\omega}$).

Next, we introduce the weighted local atoms, via which, we introduce the weighted atomic local Orlicz-Hardy space.

Definition 3.4. Let Φ satisfy Assumption (A), $\omega \in A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$ and q_ω, ρ be respectively as in (2.4) and (2.7). A triplet $(\rho,\, q,\, s)_\omega$ is called *admissible* if $q \in (q_\omega, \infty]$, $s \in \mathbb{Z}_+$ and $s \geq \lfloor n(\frac{q_\omega}{p_\Phi} - 1)\rfloor$. A function a on $\mathbb{R}^n$ is called a $(\rho,\, q,\, s)_\omega$*-atom* if there exists a cube $Q \subset \mathbb{R}^n$ such that

(i) $\mathrm{supp}\,(a) \subset Q$;

(ii) $\|a\|_{L^q_\omega(\mathbb{R}^n)} \leq [\omega(Q)]^{\frac{1}{q}-1}[\rho(\omega(Q))]^{-1}$;

(iii) $\int_{\mathbb{R}^n} a(x)x^\alpha\, dx = 0$ for all $\alpha \in \mathbb{Z}^n_+$ with $|\alpha| \leq s$, when $l(Q) < 1$.

Moreover, a function a on $\mathbb{R}^n$ is called a $(\rho,\, q)_\omega$*-single-atom* with $q \in (q_\omega, \infty]$, if

$$\|a\|_{L^q_\omega(\mathbb{R}^n)} \leq [\omega(\mathbb{R}^n)]^{\frac{1}{q}-1}[\rho(\omega(\mathbb{R}^n))]^{-1}.$$

We point out that when $\Phi(t) \equiv t^p$ for $t \in (0,\infty)$ and $p \in (0,1]$, $(\rho,\, q,\, s)_\omega$-atoms and $(\rho,\, q)_\omega$-single-atoms are respectively $(p,\, q,\, s)_\omega$-atoms and $(p,\, q)_\omega$-single-atoms, introduced by Tang [49].

Definition 3.5. Let Φ satisfy Assumption (A), $\omega \in A_\infty^{\rm loc}(\mathbb{R}^n)$, q_ω and ρ be respectively as in (2.4) and (2.7), and $(\rho,\, q,\, s)_\omega$ be admissible. The *weighted atomic local Orlicz-Hardy space* $h_\omega^{\rho,\, q,\, s}(\mathbb{R}^n)$ is defined be the set of all $f \in \mathcal{D}'(\mathbb{R}^n)$ satisfying that

$$f = \sum_{i=0}^{\infty} \lambda_i a_i$$

in $\mathcal{D}'(\mathbb{R}^n)$, where $\{a_i\}_{i\in\mathbb{N}}$ are $(\rho,\, q,\, s)_\omega$-atoms with $\mathrm{supp}\,(a_i) \subset Q_i$, a_0 is a $(\rho,\, q)_\omega$-single-atom, $\{\lambda_i\}_{i\in\mathbb{Z}_+} \subset \mathbb{C}$, and

$$\sum_{i=1}^{\infty} \omega(Q_i)\Phi\left(\frac{|\lambda_i|}{\omega(Q_i)\rho(\omega(Q_i))}\right) + \omega(\mathbb{R}^n)\Phi\left(\frac{|\lambda_0|}{\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))}\right) < \infty.$$

Moreover, letting

$$\Lambda(\{\lambda_i a_i\}_i) \equiv \inf\left\{\lambda > 0 :\ \sum_{i=1}^{\infty} \omega(Q_i)\Phi\left(\frac{|\lambda_i|}{\lambda\omega(Q_i)\rho(\omega(Q_i))}\right)\right.$$
$$\left. +\omega(\mathbb{R}^n)\Phi\left(\frac{|\lambda_0|}{\lambda\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))}\right) \le 1\right\},$$

the *quasi-norm* of $f \in h_\omega^{\rho,\, q,\, s}(\mathbb{R}^n)$ is defined by

$$\|f\|_{h_\omega^{\rho,\, q,\, s}(\mathbb{R}^n)} \equiv \inf\left\{\Lambda(\{\lambda_i a_i\}_{i\in\mathbb{Z}_+})\right\},$$

where the infimum is taken over all the decompositions of f as above.

Remark 3.6. (i) Notice that the definition of $\Lambda(\{\lambda_i a_i\}_{i\in\mathbb{Z}_+})$ above is different from [55]. In fact, if $p \in (0,1]$ and $\Phi(t) \equiv t^p$ for all $t \in (0,\infty)$, then $\Lambda(\{\lambda_i a_i\}_{i\in\mathbb{Z}_+})$ coincides with $(\sum_{i\in\mathbb{Z}_+} |\lambda_i|^p)^{1/p}$.

(ii) Let $\{\lambda_i^k\}_{i,\,k}$ and $\{a_i^k\}_{i,\,k}$ satisfy $\Lambda(\{\lambda_i^k a_i^k\}_{i\in\mathbb{Z}_+}) < \infty$, where $k = 1,\, 2$. By the fact that Φ is subadditive and of strictly lower type p_Φ, we have

$$\left[\Lambda(\{\lambda_i^1 a_i^1,\ \lambda_i^2 a_i^2\}_{i\in\mathbb{Z}_+})\right]^{p_\Phi} \le \sum_{k=1}^{2} \left[\Lambda(\{\lambda_i^k a_i^k\}_{i\in\mathbb{Z}_+})\right]^{p_\Phi}.$$

(iii) Since Φ is concave, it is of upper type 1. Thus, for any $f \in h_\omega^{\rho,\, q,\, s}(\mathbb{R}^n)$, there exist $\{a_i\}_{i\in\mathbb{Z}_+}$ and $\{\lambda_i\}_{i\in\mathbb{Z}_+}$ as in Definition 3.5 such that

$$\sum_{i\in\mathbb{Z}_+} |\lambda_i| \lesssim \Lambda(\{\lambda_i a_i\}_{i\in\mathbb{Z}_+}) \lesssim \|f\|_{h_\omega^{\rho,\, q,\, s}(\mathbb{R}^n)}.$$

Next, we introduce some local vertical, tangential and nontangential maximal functions, and then establish the characterizations of the weighted local Orlicz-Hardy space $h_{\omega,\, N}^{\Phi}(\mathbb{R}^n)$ on these local maximal functions.

Definition 3.7. Let

$$\psi_0 \in \mathcal{D}(\mathbb{R}^n) \text{ with } \int_{\mathbb{R}^n} \psi_0(x)\, dx \neq 0. \tag{3.3}$$

For $j \in \mathbb{Z}_+$, $A,\, B \in [0,\infty)$ and $y \in \mathbb{R}^n$, let $m_{j,\,A,\,B}(y) \equiv (1+2^j|y|)^A 2^{B|y|}$. The *local vertical maximal function* $\psi_0^+(f)$ of f associated to ψ_0 is defined by setting, for all $x \in \mathbb{R}^n$,

$$\psi_0^+(f)(x) \equiv \sup_{j\in\mathbb{Z}_+} |(\psi_0)_j * f(x)|, \tag{3.4}$$

the *local tangential Peetre-type maximal function* $\psi_{0,\,A,\,B}^{**}(f)$ of f associated to ψ_0 is defined by setting, for all $x \in \mathbb{R}^n$,

$$\psi_{0,\,A,\,B}^{**}(f)(x) \equiv \sup_{j\in\mathbb{Z}_+,\, y\in\mathbb{R}^n} \frac{|(\psi_0)_j * f(x-y)|}{m_{j,\,A,\,B}(y)} \tag{3.5}$$

and the *local nontangential maximal function* $(\psi_0)_\triangledown^*(f)$ of f associated to ψ_0 is defined by setting, for all $x \in \mathbb{R}^n$,

$$(\psi_0)_\triangledown^*(f)(x) \equiv \sup_{|x-y|<t<1} |(\psi_0)_t * f(y)|, \tag{3.6}$$

where and in what follows, for all $x \in \mathbb{R}^n$, $(\psi_0)_j(x) \equiv 2^{jn}\psi_0(2^j x)$ for all $j \in \mathbb{Z}_+$ and $(\psi_0)_t(x) \equiv \frac{1}{t^n}\psi_0(\frac{x}{t})$ for all $t \in (0,\infty)$.

Obviously, for any $x \in \mathbb{R}^n$, we have

$$\psi_0^+(f)(x) \le (\psi_0)_\triangledown^*(f)(x) \lesssim \psi_{0,\,A,\,B}^{**}(f)(x).$$

We remark that the local tangential Peetre-type maximal function $\psi_{0,\,A,\,B}^{**}(f)$ was introduced by Rychkov [43].

In order to establish the local vertical and the local nontangential maximal function characterizations of $h_{\omega,\,N}^\Phi(\mathbb{R}^n)$, we first establish some relations in the norm of $L_\omega^\Phi(\mathbb{R}^n)$ of the local maximal functions $\psi_{0,\,A,\,B}^{**}(f)$, $\psi_0^+(f)$ and $\widetilde{\mathcal{G}}_{N,\,R}(f)$, which further imply the desired characterizations. We begin with some technical lemmas.

LEMMA 3.8. *Let ψ_0 be as in* (3.3) *and $\psi(x) \equiv \psi_0(x) - \frac{1}{2^n}\psi_0(\frac{x}{2})$ for all $x \in \mathbb{R}^n$. Then for any given integer $L \in \mathbb{Z}_+$, there exist $\eta_0,\, \eta \in \mathcal{D}(\mathbb{R}^n)$ such that $L_\eta \ge L$ and*

$$f = \eta_0 * \psi_0 * f + \sum_{j=1}^\infty \eta_j * \psi_j * f$$

in $\mathcal{D}'(\mathbb{R}^n)$ for all $f \in \mathcal{D}'(\mathbb{R}^n)$.

Lemma 3.8 is just [43, Theorem 1.6].

REMARK 3.9. Let ψ_0, ψ, η_0 and η be as in Lemma 3.8. From the proof of [43, Theorem 1.6], it is easy to deduce that for any $j \in \mathbb{Z}_+$ and $f \in \mathcal{D}'(\mathbb{R}^n)$,

$$f = (\eta_0)_j * (\psi_0)_j * f + \sum_{k=j+1}^\infty \eta_k * \psi_k * f$$

in $\mathcal{D}'(\mathbb{R}^n)$ (see also [43, (2.11)]). We omit the details.

For $f \in L_{\rm loc}^1(\mathbb{R}^n)$, $B \in [0,\infty)$ and $x \in \mathbb{R}^n$, let

$$K_B f(x) \equiv \int_{\mathbb{R}^n} |f(y)| 2^{-B|x-y|}\, dy, \tag{3.7}$$

where and in what follows, the *space* $L_{\rm loc}^1(\mathbb{R}^n)$ denotes the set of all locally integrable functions on $\mathbb{R}^n$.

Lemma 3.10. *Let* $p\in(1,\infty)$, $q\in(1,\infty]$, *and* $\omega\in A_p^{\mathrm{loc}}(\mathbb{R}^n)$. *Then there exists a positive constant* C *such that for any sequence* $\{f^j\}_j$ *of measurable functions,*

$$\left\|\{M^{\mathrm{loc}}(f^j)\}_j\right\|_{L^p_\omega(l_q)}\le C\left\|\{f^j\}_j\right\|_{L^p_\omega(l_q)},\tag{3.8}$$

where and in what follows,

$$\left\|\{f^j\}_j\right\|_{L^p_\omega(l_q)}\equiv\left\|\left\{\sum_j|f^j|^q\right\}^{1/q}\right\|_{L^p_\omega(\mathbb{R}^n)}.$$

Also, there exists positive constants C *and* $B_0\equiv B_0(\omega,n)$ *such that for all* $B\ge B_0/p$,

$$\left\|\{K_B(f^j)\}_j\right\|_{L^p_\omega(l_q)}\le C\left\|\{f^j\}_j\right\|_{L^p_\omega(l_q)}.\tag{3.9}$$

Lemma 3.10 is just [43, Lemma 2.11]. Moreover, from the proof of [43, Lemma 2.11], it is easy to deduce that (3.8) also holds for $M^{\mathrm{loc}}_{\widetilde{C}}$ with any given positive constant $\widetilde{C}$. In this case, the positive constant C in Lemma 3.10 depends on $\widetilde{C}$.

Lemma 3.11. *Let* ψ_0 *be as in* (3.3) *and* $r\in(0,\infty)$. *Then there exists a positive constant* A_0 *depending only on the support of* ψ_0 *such that for any* $A\in(\max\{A_0,\frac{n}{r}\},\infty)$ *and* $B\in[0,\infty)$, *there exists a positive constant* C, *depending only on* n, r, ψ_0, A *and* B, *satisfying that for all* $f\in\mathcal{D}'(\mathbb{R}^n)$, $x\in\mathbb{R}^n$ *and* $j\in\mathbb{Z}_+$,

$$\begin{aligned}\left[(\psi_0)^*_{j,A,B}(f)(x)\right]^r\le C\sum_{k=j}^{\infty}2^{(j-k)(Ar-n)}&\left\{M^{\mathrm{loc}}(|(\psi_0)_k*f|^r)(x)\right.\\&\left.+K_{Br}(|(\psi_0)_k*f|^r)(x)\right\},\end{aligned}$$

where

$$(\psi_0)^*_{j,A,B}(f)(x)\equiv\sup_{y\in\mathbb{R}^n}\frac{|(\psi_0)_j*f(x-y)|}{m_{j,A,B}(y)}$$

for all $x\in\mathbb{R}^n$.

Proof. Lemma 3.11 is a modified version of [43, Lemma 2.10], and was essentially obtained by Rychkov in the proof of [43, Theorem 2.24]. Let ψ be as in Lemma 3.8. Indeed, Rychkov [43] showed Lemma 3.11 under the assumption that $f\in\mathcal{S}'_e$, namely, there exist positive constant A_f and nonnegative integer N_f such that for all $\gamma\in\mathcal{D}(\mathbb{R}^n)$,

$$|\langle f,\gamma\rangle|\le A_f\sup\left\{|\partial^\alpha\gamma(x)|e^{N_f|x|}:\ x\in\mathbb{R}^n,\ \alpha\in\mathbb{Z}^n_+\text{ and }|\alpha|\le N_f\right\},$$

which guarantees that for all $x\in\mathbb{R}^n$ and $j\in\mathbb{Z}_+$,

$$M_{A,B}(x,j)\equiv\sup_{k\ge j,\,y\in\mathbb{R}^n}2^{(j-k)A}\frac{|\psi_k*f(x-y)|}{m_{j,A,B}(y)}<\infty.$$

By [19, Proposition 2.3.4(a)], we have that for any $f\in\mathcal{D}'(\mathbb{R}^n)$, $M_{A,B}(x,j)<\infty$ for all $x\in\mathbb{R}^n$ and $j\in\mathbb{Z}_+$, provided $A>A_0$, where A_0 is a positive constant depending only on the support of ψ_0. Thus, Lemma 3.11 is true for all $f\in\mathcal{D}'(\mathbb{R}^n)$. This finishes the proof of Lemma 3.11. ■

Theorem 3.12. *Let* Φ *satisfy Assumption* (A), $\omega\in A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$, $R\in(0,\infty)$, ψ_0, q_ω *and* p_Φ *be respectively as in* (3.3) , (2.4) *and* (2.6), $\psi_0^+(f)$, $\psi^{**}_{0,A,B}(f)$, *and* $\widetilde{\mathcal{G}}_{N,R}(f)$ *be respectively* *as in* (3.4), (3.5) *and* (3.1). *Let*

$$A_1 \equiv \max\{A_0,\, nq_\omega/p_\Phi\},$$

$B_1 \equiv B_0/p_\Phi$ *and integer* $N_0 \equiv \lfloor 2A_1 \rfloor + 1$, *where* A_0 *and* B_0 *are respectively as in Lemmas 3.3 and 3.2. Then for any* $A \in (A_1, \infty)$, $B \in (B_1, \infty)$ *and integer* $N \geq N_0$, *there exists a positive constant* C, *depending only on* A, B, N, R, ψ_0, Φ, ω *and* n, *such that for all* $f \in \mathcal{D}'(\mathbb{R}^n)$,

$$\left\|\psi_{0,\,A,\,B}^{**}(f)\right\|_{L_\omega^\Phi(\mathbb{R}^n)} \leq C\left\|\psi_0^+(f)\right\|_{L_\omega^\Phi(\mathbb{R}^n)}, \tag{3.10}$$

and

$$\left\|\widetilde{\mathcal{G}}_{N,\,R}(f)\right\|_{L_\omega^\Phi(\mathbb{R}^n)} \leq C\left\|\psi_0^+(f)\right\|_{L_\omega^\Phi(\mathbb{R}^n)}. \tag{3.11}$$

Proof. Let $f \in \mathcal{D}'(\mathbb{R}^n)$. First, we prove (3.10). Let $A \in (A_1, \infty)$ and $B \in (B_1, \infty)$. By $A_1 \equiv \max\{A_0,\, nq_\omega/p_\Phi\}$ and $B_1 \equiv B_0/p_\Phi$, we know that there exists $r_0 \in (0, \frac{p_\Phi}{q_\omega})$ such that $A > \frac{n}{r_0}$ and $Br_0 > \frac{B_0}{q_\omega}$, where A_0 and B_0 are respectively as in Lemmas 3.3 and 3.10. Thus, by Lemma 3.11, for all $x \in \mathbb{R}^n$, we have

$$\begin{aligned}\left[(\psi_0)_{j,\,A,\,B}^*(f)(x)\right]^{r_0} \lesssim \sum_{k=j}^\infty 2^{(j-k)(Ar_0-n)}\Big\{&M^{\mathrm{loc}}\left(|(\psi_0)_k * f|^{r_0}\right)(x)\\ &+K_{Br_0}\left(|(\psi_0)_k * f|^{r_0}\right)(x)\Big\}.\end{aligned} \tag{3.12}$$

Let $\psi_0^+(f)$ and $\psi_{0,\,A,\,B}^{**}(f)$ be respectively as in (3.4) and (3.5). We notice that for any $x \in \mathbb{R}^n$ and $k \in \mathbb{Z}_+$,

$$|(\psi_0)_k * f(x)| \leq \psi_0^+(f)(x),$$

which together with (3.12) implies that for all $x \in \mathbb{R}^n$,

$$\left[\psi_{0,\,A,\,B}^{**}(f)(x)\right]^{r_0} \lesssim M^{\mathrm{loc}}\left([\psi_0^+(f)]^{r_0}\right)(x) + K_{Br_0}\left([\psi_0^+(f)]^{r_0}\right)(x). \tag{3.13}$$

By (3.12) and the subadditivity of Φ, we have

$$\begin{aligned}&\int_{\mathbb{R}^n} \Phi\left(\psi_{0,\,A,\,B}^{**}(f)(x)\right)\omega(x)\,dx\\ &\quad\lesssim \int_{\mathbb{R}^n} \Phi\left(\left\{M^{\mathrm{loc}}\left([\psi_0^+(f)]^{r_0}\right)(x)\right\}^{1/r_0}\right)\omega(x)\,dx\\ &\quad\quad+\int_{\mathbb{R}^n} \Phi\left(\left\{K_{Br_0}\left([\psi_0^+(f)]^{r_0}\right)(x)\right\}^{1/r_0}\right)\omega(x)\,dx \equiv \mathrm{I}_1 + \mathrm{I}_2.\end{aligned} \tag{3.14}$$

First, we estimate I_1. By $r_0 < \frac{p_\Phi}{q_\omega}$, we know that there exists $q \in (q_\omega, \infty)$ such that $r_0 q < p_\Phi$ and $\omega \in A_q^{\mathrm{loc}}(\mathbb{R}^n)$. For any $\alpha \in (0, \infty)$ and $g \in L_{\mathrm{loc}}^1(\mathbb{R}^n)$, let

$$g = g\chi_{\{x\in\mathbb{R}^n:\ |g(x)|\leq\alpha\}} + g\chi_{\{x\in\mathbb{R}^n:\ |g(x)|>\alpha\}} \equiv g_1 + g_2.$$

It is easy to see that

$$\left\{x \in \mathbb{R}^n :\, M^{\mathrm{loc}}(g)(x) > 2\alpha\right\} \subset \left\{x \in \mathbb{R}^n :\, M^{\mathrm{loc}}(g_2)(x) > \alpha\right\},$$

which together with Lemma 2.3(vi) implies that

$$\omega\left(\left\{x \in \mathbb{R}^n :\, M^{\mathrm{loc}}(g)(x) > 2\alpha\right\}\right)$$

$$\begin{aligned}&\le \omega\left(\left\{x\in\mathbb{R}^n:\, M^{\mathrm{loc}}(g_2)(x)>\alpha\right\}\right)\le \frac{1}{\alpha^q}\int_{\mathbb{R}^n}\left[M^{\mathrm{loc}}(g_2)(x)\right]^q\omega(x)\,dx\\ &\lesssim \frac{1}{\alpha^q}\int_{\mathbb{R}^n}|g_2(x)|^q\omega(x)\,dx\sim\frac{1}{\alpha^q}\int_{\{x\in\mathbb{R}^n:\,|g(x)|>\alpha\}}|g(x)|^q\omega(x)\,dx. \qquad (3.15)\end{aligned}$$

Thus, for any $\alpha\in(0,\infty)$, by (3.15), we have

$$\begin{aligned}&\omega\left(\left\{x\in\mathbb{R}^n:\,\left[M^{\mathrm{loc}}\left([\psi_0^+(f)]^{r_0}\right)(x)\right]^{1/r_0}>\alpha\right\}\right)\\ &\quad\lesssim\frac{1}{\alpha^{r_0q}}\int_{\{x\in\mathbb{R}^n:\,[\psi_0^+(f)(x)]^{r_0}>\frac{\alpha^{r_0}}{2}\}}[\psi_0^+(f)(x)]^{r_0q}\omega(x)\,dx\\ &\quad\sim\sigma_{\psi_0^+(f)}\left(\frac{\alpha}{2^{1/r_0}}\right)+\frac{1}{\alpha^{r_0q}}\int_{\frac{\alpha}{2^{1/r_0}}}^{\infty}r_0qs^{r_0q-1}\sigma_{\psi_0^+(f)}(s)\,ds, \qquad (3.16)\end{aligned}$$

where and in what follows,

$$\sigma_{\psi_0^+(f)}(s)\equiv\omega(\{x\in\mathbb{R}^n:\,\psi_0^+(f)(x)>s\}).$$

From the fact that Φ is concave and of lower type p_Φ, we infer that $\Phi(t)\sim\int_0^t\frac{\Phi(s)}{s}\,ds$ for all $t\in(0,\infty)$. By this, (3.16) and the lower type p_Φ property of Φ, the fact $r_0q<p_\Phi$ and Fubini's theorem, we have

$$\begin{aligned}\mathrm{I}_1&\sim\int_{\mathbb{R}^n}\left\{\int_0^{\left\{M^{\mathrm{loc}}([\psi_0^+(f)]^{r_0})(x)\right\}^{1/r_0}}\frac{\Phi(t)}{t}\,dt\right\}\omega(x)\,dx\\ &\sim\int_0^\infty\frac{\Phi(t)}{t}\sigma_{\left\{M^{\mathrm{loc}}([\psi_0^+(f)]^{r_0})\right\}^{1/r_0}}(t)\,dt\\ &\lesssim\int_0^\infty\frac{\Phi(t)}{t}\left\{\sigma_{\psi_0^+(f)}\left(\frac{t}{2^{1/r_0}}\right)+\frac{1}{t^{r_0q}}\int_{\frac{t}{2^{1/r_0}}}^\infty r_0qs^{r_0q-1}\sigma_{\psi_0^+(f)}(s)\,ds\right\}dt\\ &\lesssim\mathrm{J}_f+\int_0^\infty r_0qs^{r_0q-1}\sigma_{\psi_0^+(f)}(s)\left\{\int_0^{2^{\frac{1}{r_0}}s}\frac{\Phi(t)}{t}\frac{1}{t^{r_0q}}\,dt\right\}ds\\ &\sim\mathrm{J}_f+\int_0^\infty r_0qs^{r_0q-1}\sigma_{\psi_0^+(f)}(s)\Phi(2^{\frac{1}{r_0}}s)\left\{\int_0^{2^{\frac{1}{r_0}}s}\left(\frac{t}{2^{\frac{1}{r_0}}s}\right)^{p_\Phi}\frac{1}{t^{r_0q+1}}\,dt\right\}ds\\ &\sim\mathrm{J}_f\sim\int_{\mathbb{R}^n}\Phi\left(\psi_0^+(f)(x)\right)\omega(x)\,dx, \qquad (3.17)\end{aligned}$$

where $\mathrm{J}_f\equiv\int_0^\infty\frac{\Phi(t)}{t}\sigma_{\psi_0^+(f)}(t)\,dt$.

Next, we estimate I_2. For any $\alpha\in(0,\infty)$ and $g\in L^1_{\mathrm{loc}}(\mathbb{R}^n)$, let g_1 and g_2 be as above. For $H\in[\frac{B_0}{q},\infty)$, let $\int_{\mathbb{R}^n}2^{-H|x-y|}\,dy\equiv c_H$. It is easy to see that for all $x\in\mathbb{R}^n$, $K_H(g_1)(x)\le c_H\alpha$, which implies that

$$\{x\in\mathbb{R}^n:\,K_H(g)(x)>(c_H+1)\alpha\}\subset\{x\in\mathbb{R}^n:\,K_H(g_2)(x)>\alpha\},$$

where K_H is as in (3.7). Thus, by Lemma 3.10, we have

$$\omega\left(\{x\in\mathbb{R}^n:\,K_Hg(x)>(c_H+1)\alpha\}\right)\le\omega\left(\{x\in\mathbb{R}^n:\,K_Hg_2(x)>\alpha\}\right)$$

$$\lesssim \frac{1}{\alpha^q}\int_{\{x\in\mathbb{R}^n:\,|g(x)|>\alpha\}}|g(x)|^q\omega(x)\,dx.$$

Similarly to (3.16), from the above estimate, $Br_0>\frac{B_0}{q}$ and Lemma 3.2, we also deduce that

$$\begin{aligned}&\omega\left(\left\{x\in\mathbb{R}^n:\ \left[K_{Br_0}([\psi_0^+(f)]^{r_0})(x)\right]^{1/r_0}>\alpha\right\}\right)\\&\quad\lesssim \sigma_{\psi_0^+(f)}\left(\frac{\alpha}{(c_{Br_0}+1)^{1/r_0}}\right)+\frac{1}{\alpha^{r_0q}}\int_{\frac{\alpha}{(c_{Br_0}+1)^{1/r_0}}}^{\infty}r_0qs^{r_0q-1}\sigma_{\psi_0^+(f)}(s)\,ds.\end{aligned}$$

By this, similarly to the estimate of I_1, we also have

$$\mathrm{I}_2\lesssim\int_{\mathbb{R}^n}\Phi\left(\psi_0^+(f)(x)\right)\omega(x)\,dx. \tag{3.18}$$

Thus, we deduce from (3.14), (3.17) and (3.18) that

$$\int_{\mathbb{R}^n}\Phi\left(\psi_{0,A,B}^{**}(f)(x)\right)\omega(x)\,dx\lesssim\int_{\mathbb{R}^n}\Phi\left(\psi_0^+(f)(x)\right)\omega(x)\,dx.$$

Replacing f by f/λ with $\lambda\in(0,\infty)$ in the above inequality, and noticing that

$$\Phi(\psi_{0,A,B}^{**}(f/\lambda))=\Phi(\psi_{0,A,B}^{**}(f)/\lambda)$$

and $\Phi(\psi_0^+(f/\lambda))=\Phi(\psi_0^+(f)/\lambda)$, we have

$$\int_{\mathbb{R}^n}\Phi\left(\psi_{0,A,B}^{**}(f)(x)/\lambda\right)\omega(x)\,dx\lesssim\int_{\mathbb{R}^n}\Phi\left(\psi_0^+(f)(x)/\lambda\right)\omega(x)\,dx, \tag{3.19}$$

which together with the arbitrariness of $\lambda\in(0,\infty)$ implies (3.10).

Now, we prove (3.11). By $N_0\equiv\lfloor 2A_1\rfloor+1$, we know that there exists $A\in(A_1,\infty)$ such that $2A<N_0$. In the rest of this proof, we fix $A\in(A_1,\infty)$ satisfying $2A<N_0$ and $B\in(B_1,\infty)$. Let integer $N\geq N_0$ and $R\in(0,\infty)$. For any $\gamma\in\mathcal{D}_{N,R}(\mathbb{R}^n)$, $t\in(0,1)$ and $j\in\mathbb{Z}_+$, from Lemma 3.8 and Remark 3.9, it follows that

$$\gamma_t*f=\gamma_t*(\eta_0)_j*(\psi_0)_j*f+\sum_{k=j+1}^{\infty}\gamma_t*\eta_k*\psi_k*f, \tag{3.20}$$

where η_0, $\eta\in\mathcal{D}(\mathbb{R}^n)$ with $L_\eta\geq N$ and ψ is as in Lemma 3.8.

For any given $t\in(0,1)$ and $x\in\mathbb{R}^n$, let $2^{-j_0-1}\leq t<2^{-j_0}$ for some $j_0\in\mathbb{Z}_+$, and $z\in\mathbb{R}^n$ satisfy $|z-x|<t$. Then, by (3.20), we have

$$\begin{aligned}|\gamma_t*f(z)|&\leq|\gamma_t*(\eta_0)_{j_0}*(\psi_0)_{j_0}*f(z)|+\sum_{k=j_0+1}^{\infty}|\gamma_t*\eta_k*\psi_k*f(z)|\\&\leq\int_{\mathbb{R}^n}|\gamma_t*(\eta_0)_{j_0}(y)|\,|(\psi_0)_{j_0}*f(z-y)|\,dy\\&\quad+\sum_{k=j_0+1}^{\infty}\int_{\mathbb{R}^n}|\gamma_t*\eta_k(y)|\,|\psi_k*f(z-y)|\,dy\equiv\mathrm{I}_3+\mathrm{I}_4.\end{aligned} \tag{3.21}$$

To estimate I_3, from

$$\psi_{0,A,B}^{**}(f)(x)=\sup_{j\in\mathbb{Z}_+,\,y\in\mathbb{R}^n}\frac{|(\psi_0)_j*f(x-y)|}{m_{j,A,B}(y)}$$

$$= \sup_{j\in\mathbb{Z}_+,\, y\in\mathbb{R}^n} \frac{|(\psi_0)_j * f(x-(y+x-z))|}{m_{j,\,A,\,B}(y+x-z)}$$
$$= \sup_{j\in\mathbb{Z}_+,\, y\in\mathbb{R}^n} \frac{|(\psi_0)_j * f(z-y)|}{m_{j,\,A,\,B}(y+x-z)},$$

we infer that

$$|(\psi_0)_{j_0} * f(z-y)| \le \psi^{**}_{0,\,A,\,B}(f)(x) m_{j_0,\,A,\,B}(y+x-z),$$

which, together with the facts that

$$m_{j_0,\,A,\,B}(y+x-z) \le m_{j_0,\,A,\,B}(x-z) m_{j_0,\,A,\,B}(y)$$

and $m_{j_0,\,A,\,B}(x-z) \lesssim 2^A$, implies that

$$|(\psi_0)_{j_0} * f(z-y)| \lesssim 2^A \psi^{**}_{0,\,A,\,B}(f)(x) m_{j_0,\,A,\,B}(y).$$

Thus, we have

$$\mathrm{I}_3 \lesssim 2^A \left\{ \int_{\mathbb{R}^n} |\gamma_t * (\eta_0)_{j_0}(y)| m_{j_0,\,A,\,B}(y)\, dy \right\} \psi^{**}_{0,\,A,\,B}(f)(x).$$

To estimate I_4, by the definition of ψ, it is easy to know that for any $k \in \mathbb{N}$,

$$|\psi_k * f(z-y)| \le |(\psi_0)_k * f(z-y)| + |(\psi_0)_{k-1} * f(z-y)|.$$

By the definition of $\psi^{**}_{0,\,A,\,B}(f)$ and the facts that

$$m_{k,\,A,\,B}(y+x-z) \le m_{k,\,A,\,B}(x-z) m_{k,\,A,\,B}(y)$$

for any $k \in \mathbb{N}$ and $m_{k,\,A,\,B}(x-z) \lesssim 2^{(k-j_0)A}$, we conclude that

$$\begin{aligned}|(\psi_0)_k * f(z-y)| &\le \psi^{**}_{0,\,A,\,B}(f)(x) m_{k,\,A,\,B}(y+x-z)\\ &\le \psi^{**}_{0,\,A,\,B}(f)(x) m_{k,\,A,\,B}(x-z) m_{k,\,A,\,B}(y)\\ &\lesssim 2^{(k-j_0)A} m_{k,\,A,\,B}(y) \psi^{**}_{0,\,A,\,B}(f)(x).\end{aligned}$$

Similarly, we also have

$$|(\psi_0)_{k-1} * f(z-y)| \lesssim 2^{(k-j_0)A} m_{k,\,A,\,B}(y) \psi^{**}_{0,\,A,\,B}(f)(x).$$

Thus,

$$\mathrm{I}_4 \lesssim \sum_{k=j_0+1}^{\infty} 2^{(k-j_0)A} \left\{ \int_{\mathbb{R}^n} |\gamma_t * \eta_k(y)| m_{k,\,A,\,B}(y)\, dy \right\} \psi^{**}_{0,\,A,\,B}(f)(x).$$

From (3.21) and the above estimates of I_3 and I_4, it follows that

$$\begin{aligned}|\gamma_t * f(z)| \lesssim &\left\{ \int_{\mathbb{R}^n} |\gamma_t * (\eta_0)_{j_0}(y)| m_{j_0,\,A,\,B}(y)\, dy \right.\\ &\left. + \sum_{k=j_0+1}^{\infty} 2^{(k-j_0)A} \int_{\mathbb{R}^n} |\gamma_t * \eta_k(y)| m_{k,\,A,\,B}(y)\, dy \right\} \psi^{**}_{0,\,A,\,B}(f)(x). \quad (3.22)\end{aligned}$$

Assume that $\operatorname{supp}(\eta_0) \subset B(0, R_0)$. Then $\operatorname{supp}((\eta_0)_j) \subset B(0, 2^{-j}R_0)$ for all $j \in \mathbb{Z}_+$. Moreover, by $\operatorname{supp}(\gamma) \subset B(0, R)$ and $2^{-j_0-1} \le t < 2^{-j_0}$, we see that

$$\operatorname{supp}(\gamma_t) \subset B(0, 2^{-j_0}R).$$

From this, we further deduce that $\operatorname{supp}(\gamma_t * (\eta_0)_{j_0}) \subset B(0, 2^{-j_0}(R_0+R))$ and

$$|\gamma_t * (\eta_0)_{j_0}(y)| \lesssim \int_{\mathbb{R}^n} |\gamma_t(s)||(\psi_0)_{j_0}(y-s)|\,ds \lesssim 2^{j_0 n}\int_{\mathbb{R}^n} |\gamma_t(s)|\,ds \sim 2^{j_0 n},$$

which implies that

$$\begin{aligned}\int_{\mathbb{R}^n} &|\gamma_t * (\eta_0)_{j_0}(y)| m_{j_0,\,A,\,B}(y)\,dy\\ &\lesssim 2^{j_0 n}\int_{B(0,2^{-j_0}(R_0+R))} (1+2^{j_0}|y|)^A 2^{B|y|}\,dy \lesssim 1.\end{aligned} \tag{3.23}$$

Moreover, since η has vanishing moments up to order N, it was proved in [43, (2.13)] that

$$\|\gamma_t * \eta_k\|_{L^\infty(\mathbb{R}^n)} \lesssim 2^{(j_0-k)N}2^{j_0 n}$$

for all $k \in \mathbb{N}$ with $k \ge j_0+1$, which, together with the facts that $N > 2A$ and

$$\operatorname{supp}(\gamma_t * \eta_k) \subset B(0, 2^{-j_0}R_0 + 2^{-k}R),$$

implies that

$$\begin{aligned}\sum_{k=j_0+1}^{\infty} &2^{(k-j_0)A}\int_{\mathbb{R}^n} |\gamma_t * \eta_k(y)| m_{k,\,A,\,B}(y)\,dy\\ &\lesssim \sum_{k=j_0+1}^{\infty} 2^{(k-j_0)A}2^{(j_0-k)N}2^{j_0 n}(2^{-j_0}R_0+2^{-k}R)^n\\ &\quad\times \left[1+2^k(2^{-j_0}R_0+2^{-k}R)\right]^A 2^{(2^{-j_0}R_0+2^{-k}R)B}\\ &\lesssim \sum_{k=j_0+1}^{\infty} 2^{(j_0-k)(N-2A)} \lesssim 1.\end{aligned} \tag{3.24}$$

Thus, from (3.22), (3.23) and (3.24), we deduce that $|\gamma_t * f(z)| \lesssim \psi_{0,\,A,\,B}^{**}(f)(x)$. Then, by the arbitrariness of $t \in (0,1)$ and $z \in B(x,t)$, we know that

$$\widetilde{\mathcal{G}}_{N,R}(f)(x) \lesssim \psi_{0,\,A,\,B}^{**}(f)(x),$$

which together with (3.19) implies that for any $\lambda \in (0,\infty)$,

$$\int_{\mathbb{R}^n} \Phi\left(\widetilde{\mathcal{G}}_{N,\,R}(f)(x)/\lambda\right)\omega(x)\,dx \lesssim \int_{\mathbb{R}^n} \Phi\left(\psi_0^+(f)(x)/\lambda\right)\omega(x)\,dx.$$

From this, we infer that (3.11) holds, which completes the proof of Theorem 3.12. ■

REMARK 3.13. Let $p \in (0,1]$. We point out that Theorem 3.12 when $R \equiv 1$ and $\Phi(t) \equiv t^p$ for all $t \in (0,\infty)$ was obtained by Rychkov [43, Theorem 2.24].

As a corollary of Theorem 3.1, we immediately obtain that the local vertical and the local nontangential maximal function characterizations of $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$ with $N \ge N_{\Phi,\,\omega}$ as follows. Here and in what follows,

$$N_{\Phi,\,\omega} \equiv \max\left\{\widetilde{N}_{\Phi,\,\omega},\, N_0\right\}, \tag{3.25}$$

where $\widetilde{N}_{\Phi,\,\omega}$ and N_0 are respectively as in Definition 3.3 and Theorem 3.12.

THEOREM 3.14. *Let* Φ *satisfy Assumption* (A), $\omega \in A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$, ψ_0 *and* $N_{\Phi,\,\omega}$ *be respectively as in* (3.3) *and* (3.25). *Then for any integer* $N \ge N_{\Phi,\,\omega}$, *the following are equivalent:*

(i) $f \in h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$;
(ii) $f \in \mathcal{D}'(\mathbb{R}^n)$ *and* $\psi_0^+(f) \in L^{\Phi}_{\omega}(\mathbb{R}^n)$;
(iii) $f \in \mathcal{D}'(\mathbb{R}^n)$ *and* $(\psi_0)^*_{\triangledown}(f) \in L^{\Phi}_{\omega}(\mathbb{R}^n)$;
(iv) $f \in \mathcal{D}'(\mathbb{R}^n)$ *and* $\widetilde{\mathcal{G}}_N(f) \in L^{\Phi}_{\omega}(\mathbb{R}^n)$;
(v) $f \in \mathcal{D}'(\mathbb{R}^n)$ *and* $\widetilde{\mathcal{G}}^0_N(f) \in L^{\Phi}_{\omega}(\mathbb{R}^n)$;
(vi) $f \in \mathcal{D}'(\mathbb{R}^n)$ *and* $\mathcal{G}^0_N(f) \in L^{\Phi}_{\omega}(\mathbb{R}^n)$.

Moreover,

$$\begin{aligned}\|f\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)} &\sim \left\|\psi_0^+(f)\right\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)} \sim \|(\psi_0)^*_{\triangledown}(f)\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)}\\ &\sim \left\|\widetilde{\mathcal{G}}_N(f)\right\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)} \sim \left\|\widetilde{\mathcal{G}}^0_N(f)\right\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)} \sim \left\|\mathcal{G}^0_N(f)\right\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)}, \qquad (3.26)\end{aligned}$$

where the implicit constants are independent of f.

Proof. (i) ⇒ (ii). Let integer $N \ge N_{\Phi,\,\omega}$ and $f \in h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$. Let $\widetilde{\psi}_0$ satisfy (3.3) and $\widetilde{\psi}_0 \in \mathcal{D}_N(\mathbb{R}^n)$. Then from the definition of $\mathcal{G}_N(f)$, we infer that $\widetilde{\psi}_0^+(f) \le \mathcal{G}_N(f)$ and hence $\widetilde{\psi}_0^+(f) \in L^{\Phi}_{\omega}(\mathbb{R}^n)$. For any ψ_0 satisfying (3.3), assume that $\operatorname{supp}(\psi_0) \subset B(0,R)$. Then, by (3.11) and the above argument, we have

$$\left\|\widetilde{\mathcal{G}}_{N,\,R}(f)\right\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)} \lesssim \left\|\widetilde{\psi}_0^+(f)\right\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)} \lesssim \|f\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)},$$

which together with $\psi_0^+(f) \lesssim \widetilde{\mathcal{G}}_{N,\,R}(f)$ implies that $\psi_0^+(f) \in L^{\Phi}_{\omega}(\mathbb{R}^n)$ and

$$\left\|\psi_0^+(f)\right\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)} \lesssim \|f\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}.$$

(ii) ⇒ (iii). Let $f \in \mathcal{D}'(\mathbb{R}^n)$ satisfy $\psi_0^+(f) \in L^{\Phi}_{\omega}(\mathbb{R}^n)$, where ψ_0 is as in (3.3). Then from the fact that

$$\psi_0^+(f) \le (\psi_0)^*_{\triangledown}(f) \lesssim \psi^{**}_{0,\,A,\,B}(f)$$

and (3.10), we deduce that $(\psi_0)^*_{\triangledown}(f) \in L^{\Phi}_{\omega}(\mathbb{R}^n)$ and

$$\|(\psi_0)^*_{\triangledown}(f)\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)} \lesssim \|\psi_0^+(f)\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)}.$$

(iii) ⇒ (iv). Let $f \in \mathcal{D}'(\mathbb{R}^n)$ satisfy $(\psi_0)^*_{\triangledown}(f) \in L^{\Phi}_{\omega}(\mathbb{R}^n)$, where ψ_0 is as in (3.3). By (3.11),

$$\|\widetilde{\mathcal{G}}_N(f)\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)} \lesssim \|\psi_0^+(f)\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)},$$

which together with the fact that

$$\psi_0^+(f) \le (\psi_0)^*_{\triangledown}(f)$$

and the assumption that $(\psi_0)^*_{\triangledown}(f) \in L^{\Phi}_{\omega}(\mathbb{R}^n)$ implies $\widetilde{\mathcal{G}}_N(f) \in L^{\Phi}_{\omega}(\mathbb{R}^n)$ and

$$\left\|\widetilde{\mathcal{G}}_N(f)\right\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)} \lesssim \|(\psi_0)^*_{\triangledown}(f)\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)}.$$

(iv) ⇒ (v) ⇒ (vi). By the facts that $\mathcal{G}^0_N(f) \le \widetilde{\mathcal{G}}^0_N(f) \le \widetilde{\mathcal{G}}_N(f)$ for any $f \in \mathcal{D}'(\mathbb{R}^n)$ and Φ is increasing, we see that all the conclusions hold. Moreover, it is obvious that

$$\|\mathcal{G}^0_N(f)\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)} \le \left\|\widetilde{\mathcal{G}}^0_N(f)\right\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)} \le \left\|\widetilde{\mathcal{G}}_N(f)\right\|_{L^{\Phi}_{\omega}(\mathbb{R}^n)}.$$

(vi) ⇒ (i). Let $f\in\mathcal{D}'(\mathbb{R}^n)$ satisfy

$$\mathcal{G}^0_N(f)\in L^\Phi_\omega(\mathbb{R}^n).$$

Let ψ_1 satisfy (3.3) and $\psi_1\in\mathcal{D}^0_N(\mathbb{R}^n)$. Then by (3.10), we have that

$$\left\|\widetilde{\mathcal{G}}_N(f)\right\|_{L^\Phi_\omega(\mathbb{R}^n)}\lesssim\|\psi^+_1(f)\|_{L^\Phi_\omega(\mathbb{R}^n)},$$

which together with the facts that $\psi^+_1(f)\le\mathcal{G}^0_N(f)$ and $\mathcal{G}_N(f)\le\widetilde{\mathcal{G}}_N(f)$ implies that

$$\|\mathcal{G}_N(f)\|_{L^\Phi_\omega(\mathbb{R}^n)}\lesssim\left\|\mathcal{G}^0_N(f)\right\|_{L^\Phi_\omega(\mathbb{R}^n)}.$$

Thus, by the definition of $h^\Phi_{\omega,\,N}(\mathbb{R}^n)$, we know that $f\in h^\Phi_{\omega,\,N}(\mathbb{R}^n)$ and

$$\|f\|_{h^\Phi_{\omega,\,N}(\mathbb{R}^n)}\lesssim\|\mathcal{G}^0_N(f)\|_{L^\Phi_\omega(\mathbb{R}^n)},$$

which completes the proof of Theorem 3.14. ■

As a corollary of Theorems 3.12 and 3.14, we have the following local tangential maximal function characterization of $h^\Phi_{\omega,\,N}(\mathbb{R}^n)$. We omit the details.

Corollary 3.15. *Let Φ satisfy Assumption* (A)*, ψ_0 be as in* (3.3)*, $\omega\in A^{\rm loc}_\infty(\mathbb{R}^n)$, $N_{\Phi,\,\omega}$ be as in* (3.25)*, A and B be as in Theorem 3.12. Then for integer $N\ge N_{\Phi,\,\omega}$,*

$$f\in h^\Phi_{\omega,\,N}(\mathbb{R}^n)$$

if and only if $f\in\mathcal{D}'(\mathbb{R}^n)$ and $\psi^{**}_{0,\,A,\,B}(f)\in L^\Phi_\omega(\mathbb{R}^n)$; moreover,

$$\|f\|_{h^\Phi_{\omega,\,N}(\mathbb{R}^n)}\sim\|\psi^{**}_{0,\,A,\,B}(f)\|_{L^\Phi_\omega(\mathbb{R}^n)}.$$

Next, we give some basic properties concerning $h^\Phi_{\omega,\,N}(\mathbb{R}^n)$ and $h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)$.

Proposition 3.16. *Let Φ satisfy Assumption* (A)*, $\omega\in A^{\rm loc}_\infty(\mathbb{R}^n)$ and $N_{\Phi,\,\omega}$ be as in* (3.25)*. If integer $N\ge N_{\Phi,\,\omega}$, then the inclusion $h^\Phi_{\omega,\,N}(\mathbb{R}^n)\hookrightarrow\mathcal{D}'(\mathbb{R}^n)$ is continuous.*

Proof. Let $f\in h^\Phi_{\omega,\,N}(\mathbb{R}^n)$. For any given $\phi\in\mathcal{D}(\mathbb{R}^n)$, assume that $\operatorname{supp}(\phi)\subset B(0,R)$ with $R\in(0,\infty)$. Then we have

$$|\langle f,\phi\rangle|=\left|f*\widetilde{\phi}(0)\right|\le\left\|\widetilde{\phi}\right\|_{\mathcal{D}_{N,\,R}(\mathbb{R}^n)}\inf_{x\in B(0,1)}\widetilde{\mathcal{G}}_{N,\,R}(f)(x),\tag{3.27}$$

where $\widetilde{\mathcal{G}}_{N,\,R}(f)$ is as in (3.2) and $\widetilde{\phi}(x)\equiv\phi(-x)$ for all $x\in\mathbb{R}^n$. Now, to prove Proposition 3.16, we consider the following two cases for $\|f\|_{h^\Phi_{\omega,\,N}(\mathbb{R}^n)}$.

Case (i) $\|f\|_{h^\Phi_{\omega,\,N}(\mathbb{R}^n)}\ge1$. In this case, by the upper type 1 property of Φ and Theorems 3.12 and 3.14, we obtain

$$\begin{aligned}&\int_{\mathbb{R}^n}\Phi\left(\widetilde{\mathcal{G}}_{N,\,R}(f)(x)\right)\omega(x)\,dx\\&\quad\lesssim\|f\|_{h^\Phi_{\omega,\,N}(\mathbb{R}^n)}\int_{\mathbb{R}^n}\Phi\left(\frac{\widetilde{\mathcal{G}}_{N,\,R}(f)(x)}{\|f\|_{h^\Phi_{\omega,\,N}(\mathbb{R}^n)}}\right)\omega(x)\,dx\lesssim\|f\|_{h^\Phi_{\omega,\,N}(\mathbb{R}^n)}.\end{aligned}\tag{3.28}$$

Notice that the upper type 1 property of Φ implies that for $t\in(0,1]$,

$$\Phi(1)=\Phi\left(t\frac{1}{t}\right)\lesssim\frac{1}{t}\Phi(t)$$

and hence $\Phi(t) \gtrsim t$. Thus, when $\inf_{x\in B(0,1)} \widetilde{\mathcal{G}}_{N,\,R}(f)(x) \le 1$, from (3.27) and (3.28), we deduce that

$$\begin{aligned}|\langle f,\phi\rangle| &\lesssim \|\phi\|_{\mathcal{D}_{N,\,R}(\mathbb{R}^n)}\Phi\left(\inf_{x\in B(0,1)} \widetilde{\mathcal{G}}_{N,\,R}(f)(x)\right)\\ &\lesssim \|\phi\|_{\mathcal{D}_{N,\,R}(\mathbb{R}^n)}\frac{1}{\omega(B(0,1))}\int_{B(0,1)} \Phi\left(\widetilde{\mathcal{G}}_{N,\,R}(f)(y)\right)\omega(y)\,dy\\ &\lesssim \|\phi\|_{\mathcal{D}_{N,\,R}(\mathbb{R}^n)}\frac{1}{\omega(B(0,1))}\|f\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}. \end{aligned}\tag{3.29}$$

Let p_Φ be as in (2.6). Since Φ is lower type p_Φ, then for $t\in(1,\infty)$,

$$\Phi(1) = \Phi\left(t\frac{1}{t}\right) \lesssim \frac{1}{t^{p_\Phi}}\Phi(t)$$

and hence $t \lesssim [\Phi(t)]^{1/p_\Phi}$. Thus, when $\inf_{x\in B(0,1)} \widetilde{\mathcal{G}}_{N,\,R}(f)(x) > 1$, by (3.26) and (3.27), we conclude that

$$\begin{aligned}|\langle f,\phi\rangle| &\lesssim \|\phi\|_{\mathcal{D}_{N,\,R}(\mathbb{R}^n)}\left\{\Phi\left(\inf_{x\in B(0,1)} \widetilde{\mathcal{G}}_{N,\,R}(f)(x)\right)\right\}^{1/p_\Phi}\\ &\lesssim \|\phi\|_{\mathcal{D}_{N,\,R}(\mathbb{R}^n)}[\omega(B(0,1))]^{-1/p_\Phi}\\ &\quad\times\left\{\int_{B(0,1)} \Phi\left(\widetilde{\mathcal{G}}_{N,\,R}(f)(y)\right)\omega(y)\,dy\right\}^{1/p_\Phi}\\ &\lesssim \|\phi\|_{\mathcal{D}_{N,\,R}(\mathbb{R}^n)}[\omega(B(0,1))]^{-1/p_\Phi}\|f\|^{1/p_\Phi}_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}. \end{aligned}\tag{3.30}$$

Case (ii) $\|f\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)} < 1$. In this case, by the lower type p_Φ property of Φ and Theorems 3.12 and 3.14, we see that

$$\begin{aligned}&\int_{\mathbb{R}^n} \Phi\left(\widetilde{\mathcal{G}}_{N,\,R}(f)(x)\right)\omega(x)\,dx\\ &\quad\lesssim \|f\|^{p_\Phi}_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}\int_{\mathbb{R}^n}\Phi\left(\frac{\widetilde{\mathcal{G}}_{N,\,R}(f)(x)}{\|f\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}}\right)\omega(x)\,dx \lesssim \|f\|^{p_\Phi}_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}.\end{aligned}$$

Thus, from this fact and (3.27), similarly to the proof of (3.29) and (3.30), we infer that if $\inf_{x\in B(0,1)} \widetilde{\mathcal{G}}_{N,\,R}(f)(x) \le 1$, then

$$|\langle f,\phi\rangle| \lesssim \|\phi\|_{\mathcal{D}_{N,\,R}(\mathbb{R}^n)}\frac{1}{\omega(B(0,1))}\|f\|^{p_\Phi}_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)},$$

and if $\inf_{x\in B(0,1)} \widetilde{\mathcal{G}}_{N,\,R}(f)(x) > 1$, then

$$|\langle f,\phi\rangle| \lesssim \|\phi\|_{\mathcal{D}_{N,\,R}(\mathbb{R}^n)}[\omega(B(0,1))]^{-1/p_\Phi}\|f\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}.$$

Thus, $f\in\mathcal{D}'(\mathbb{R}^n)$ and the inclusion is continuous, which completes the proof of Proposition 3.16. ∎

PROPOSITION 3.17. *Let Φ satisfy Assumption* (A), *$\omega\in A^{\rm loc}_\infty(\mathbb{R}^n)$ and $N_{\Phi,\,\omega}$ be as in* (3.25). *If integer $N \ge N_{\Phi,\,\omega}$, then the space $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$ is complete.*

Proof. For any $\psi\in\mathcal{D}_N(\mathbb{R}^n)$ and $\{f_i\}_{i\in\mathbb{N}}\subset\mathcal{D}'(\mathbb{R}^n)$ such that $\{\sum_{i=1}^j f_i\}_{j\in\mathbb{N}}$ converges in $\mathcal{D}'(\mathbb{R}^n)$ to a distribution f as $j\to\infty$, the series $\{\sum_{i=1}^j f_i*\psi\}_{j\in\mathbb{N}}$ converges to $f*\psi$

also pointwise as $j \to \infty$. By Assumption (A), we know that Φ is strictly increasing and subadditive, which together with the continuity of Φ implies that for all $x \in \mathbb{R}^n$,

$$\Phi\left(\mathcal{G}_N(f)(x)\right) \le \Phi\left(\sum_{i=1}^{\infty} \mathcal{G}_N(f_i)(x)\right) \le \sum_{i=1}^{\infty} \Phi\left(\mathcal{G}_N(f_i)(x)\right).$$

If $\sum_{i=1}^{\infty} \|f_i\|^{p_\Phi}_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)} < \infty$, letting $\lambda_i = \|f_i\|^{p_\Phi}_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}$, then by the strictly lower type p_Φ property of Φ and the levi lemma, we know that

$$\begin{aligned}
&\int_{\mathbb{R}^n} \Phi\left(\frac{\mathcal{G}_N(f)(x)}{(\sum_{j=1}^{\infty}\lambda_j)^{1/p_\Phi}}\right)\omega(x)\,dx \\
&\quad\le \sum_{i=1}^{\infty} \int_{\mathbb{R}^n} \Phi\left(\frac{\mathcal{G}_N(f_i)(x)}{(\sum_{j=1}^{\infty}\lambda_j)^{1/p_\Phi}}\right)\omega(x)\,dx \\
&\quad\le \sum_{i=1}^{\infty} \frac{\lambda_i}{\sum_{j=1}^{\infty}\lambda_j} \int_{\mathbb{R}^n} \Phi\left(\frac{\mathcal{G}_N(f_i)(x)}{\lambda_i^{1/p_\Phi}}\right)\omega(x)\,dx \le \sum_{i=1}^{\infty} \frac{\lambda_i}{\sum_{j=1}^{\infty}\lambda_j} = 1,
\end{aligned}$$

which further implies that

$$\|f\|^{p_\Phi}_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)} \le \sum_{i=1}^{\infty} \|f_i\|^{p_\Phi}_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}. \tag{3.31}$$

To prove that $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$ is complete, it suffices to show that for every sequence $\{f_j\}_{j\in\mathbb{N}}$ with $\|f_j\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)} < 2^{-j}$ for any $j \in \mathbb{N}$, the series $\{f_j\}_{j\in\mathbb{N}}$ converges in $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$. Since $\{\sum_{i=1}^{j} f_i\}_{j\in\mathbb{N}}$ is a Cauchy sequence in $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$, by Proposition 3.16 and the completeness of $\mathcal{D}'(\mathbb{R}^n)$, $\{\sum_{i=1}^{j} f_i\}_{j\in\mathbb{N}}$ is also a Cauchy sequence in $\mathcal{D}'(\mathbb{R}^n)$ and thus converges to some $f \in \mathcal{D}'(\mathbb{R}^n)$. Therefore, by (3.31),

$$\left\|f - \sum_{i=1}^{j} f_i\right\|^{p_\Phi}_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)} = \left\|\sum_{i=j+1}^{\infty} f_i\right\|^{p_\Phi}_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)} \le \sum_{i=j+1}^{\infty} 2^{-ip_\Phi} \to 0$$

as $j \to \infty$, which completes the proof of Proposition 3.17. ■

THEOREM 3.18. *Let Φ satisfy Assumption* (A), *$\omega \in A^{\mathrm{loc}}_{\infty}(\mathbb{R}^n)$ and $N_{\Phi,\,\omega}$ be as in* (3.25). *If $(\rho,\,q,\,s)_\omega$ is admissible (see Definition 3.4) and integer $N \ge N_{\Phi,\,\omega}$, then*

$$h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n) \subset h^{\Phi}_{\omega,\,N_{\Phi,\,\omega}}(\mathbb{R}^n) \subset h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$$

and, moreover, there exists a positive constant C such that for all $f \in h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$,

$$\|f\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)} \le \|f\|_{h^{\Phi}_{\omega,\,N_{\Phi,\,\omega}}(\mathbb{R}^n)} \le C\|f\|_{h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)}.$$

Proof. Obviously, by Definition 3.3, we only need prove that $h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n) \subset h^{\Phi}_{\omega,\,N_{\Phi,\,\omega}}(\mathbb{R}^n)$, and for all $f \in h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$,

$$\|f\|_{h^{\Phi}_{\omega,\,N_{\Phi,\,\omega}}(\mathbb{R}^n)} \lesssim \|f\|_{h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)}.$$

To this end, by Theorem 3.14 and Definition 3.5, it suffices to prove that for any $(\rho,\,q)_\omega$-single-atom a and $\lambda \in \mathbb{C}$,

$$\int_{\mathbb{R}^n} \Phi\left(\mathcal{G}^0_{N_{\Phi,\,\omega}}(\lambda a)(x)\right)\omega(x)\,dx \lesssim \omega(\mathbb{R}^n)\Phi\left(\frac{|\lambda|}{\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))}\right), \tag{3.32}$$

and for any $(\rho,\, q,\, s)_\omega$-atom a supported in the cube Q and $\lambda \in \mathbb{C}$,

$$\int_{\mathbb{R}^n} \Phi\left(\mathcal{G}^0_{N_\Phi,\,\omega}(\lambda a)(x)\right)\omega(x)\,dx \lesssim \omega(Q)\Phi\left(\frac{|\lambda|}{\omega(Q)\rho(\omega(Q))}\right). \tag{3.33}$$

Indeed, for any $f \in h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)$,

$$f = \sum_{i=0}^\infty \lambda_i a_i$$

in $\mathcal{D}'(\mathbb{R}^n)$, where $\{\lambda_i\}_{i=0}^\infty \subset \mathbb{C}$, a_0 is a $(\rho,\, q)_\omega$-single-atom and for any $i \in \mathbb{N}$, a_i is a $(\rho,\, q,\, s)_\omega$-atom supported in the cube Q_i. Then, for any $\lambda \in (0,\infty)$, by the facts that $\mathcal{G}^0_{N_\Phi,\,\omega}(f/\lambda) = \mathcal{G}^0_{N_\Phi,\,\omega}(f)/\lambda$ and Φ is strictly increasing, subadditive and continuous, (3.32) and (3.33), we have

$$\begin{aligned}
&\int_{\mathbb{R}^n} \Phi\left(\frac{\mathcal{G}^0_{N_\Phi,\,\omega}(f)(x)}{\lambda}\right)\omega(x)\,dx\\
&\quad= \int_{\mathbb{R}^n} \Phi\left(\mathcal{G}^0_{N_\Phi,\,\omega}\left(\frac{f}{\lambda}\right)(x)\right)\omega(x)\,dx \le \sum_{i=0}^\infty \int_{\mathbb{R}^n} \Phi\left(\mathcal{G}^0_{N_\Phi,\,\omega}\left(\frac{\lambda_i a_i}{\lambda}\right)(x)\right)\omega(x)\,dx\\
&\quad\lesssim \omega(\mathbb{R}^n)\Phi\left(\frac{|\lambda_0|}{\lambda\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))}\right) + \sum_{i=1}^\infty \omega(Q_i)\Phi\left(\frac{|\lambda_i|}{\lambda\omega(Q_i)\rho(\omega(Q_i))}\right),
\end{aligned}$$

which together with Theorem 3.14 implies that $\|f\|_{h^\Phi_{\omega,\,N_\Phi,\,\omega}(\mathbb{R}^n)} \lesssim \|f\|_{h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)}$.

We now prove (3.32). Since $q \in (q_\omega, \infty]$, by the definition of q_ω, we have $\omega \in A^{\rm loc}_q(\mathbb{R}^n)$. Let a be a $(\rho,\, q)_\omega$-single-atom and $\lambda \in \mathbb{C}$. When $\omega(\mathbb{R}^n) = \infty$, by the definition of the single atom, we know that $a = 0$ for almost every $x \in \mathbb{R}^n$. In this case, it is easy to see that (3.32) holds. When $\omega(\mathbb{R}^n) < \infty$, since Φ is concave, from Jensen's inequality, Hölder's inequality and Proposition 3.2(ii), we deduce that

$$\begin{aligned}
&\int_{\mathbb{R}^n} \Phi\left(\mathcal{G}^0_{N_\Phi,\,\omega}(\lambda a)(x)\right)\omega(x)\,dx\\
&\quad\le \omega(\mathbb{R}^n)\Phi\left(\frac{1}{\omega(\mathbb{R}^n)}\int_{\mathbb{R}^n} \mathcal{G}^0_{N_\Phi,\,\omega}(\lambda a)(x)\omega(x)\,dx\right)\\
&\quad\le \omega(\mathbb{R}^n)\Phi\left(\frac{1}{[\omega(\mathbb{R}^n)]^{1/q}}\left\{\int_{\mathbb{R}^n}\left[\mathcal{G}^0_{N_\Phi,\,\omega}(\lambda a)(x)\right]^q\omega(x)\,dx\right\}^{1/q}\right)\\
&\quad\lesssim \omega(\mathbb{R}^n)\Phi\left(\frac{1}{[\omega(\mathbb{R}^n)]^{1/q}}|\lambda|\|a\|_{L^q_\omega(\mathbb{R}^n)}\right) \lesssim \omega(\mathbb{R}^n)\Phi\left(\frac{|\lambda|}{\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))}\right).
\end{aligned}$$

That is, (3.32) holds.

Next, we prove (3.33). Let a be a $(\rho,\, q,\, s)_\omega$-atom supported in the cube $Q \equiv Q(x_0, r)$, and $\lambda \in \mathbb{C}$. We consider the following two cases for Q.

Case 1) $|Q| < 1$. In this case, letting $\widetilde{Q} \equiv 2\sqrt{n}Q$, then we have

$$\begin{aligned}
&\int_{\mathbb{R}^n} \Phi\left(\mathcal{G}^0_{N_\Phi,\,\omega}(\lambda a)(x)\right)\omega(x)\,dx\\
&\quad= \int_{\widetilde{Q}} \Phi\left(\mathcal{G}^0_{N_\Phi,\,\omega}(\lambda a)(x)\right)\omega(x)\,dx + \int_{\widetilde{Q}^\complement} \cdots \equiv {\rm I}_1 + {\rm I}_2.
\end{aligned} \tag{3.34}$$

For I_1, by Jensen's inequality, Hölder's inequality, Lemma 2.3(v) and Proposition 3.2(ii), we have

$$\begin{aligned}\mathrm{I}_1 &\le \omega(\widetilde{Q})\Phi\left(\frac{1}{\omega(\widetilde{Q})}\int_{\widetilde{Q}}\mathcal{G}^0_{N_\Phi,\,\omega}(\lambda a)(x)\omega(x)\,dx\right)\\ &\le \omega(\widetilde{Q})\Phi\left(\frac{1}{[\omega(\widetilde{Q})]^{1/q}}\left\{\int_{\widetilde{Q}}\left[\mathcal{G}^0_{N_\Phi,\,\omega}(\lambda a)(x)\right]^q\omega(x)\,dx\right\}^{1/q}\right)\\ &\lesssim \omega(\widetilde{Q})\Phi\left(\frac{1}{[\omega(\widetilde{Q})]^{1/q}}|\lambda|\|a\|_{L^q_\omega(\mathbb{R}^n)}\right)\\ &\lesssim \omega(\widetilde{Q})\Phi\left(\frac{|\lambda|}{\omega(Q)\rho(\omega(Q))}\right)\lesssim \omega(Q)\Phi\left(\frac{|\lambda|}{\omega(Q)\rho(\omega(Q))}\right), \qquad (3.35)\end{aligned}$$

which is the desired estimate for I_1.

To estimate I_2, we claim that for all $x\in\widetilde{Q}^{\complement}$,

$$\begin{aligned}\mathcal{G}^0_{N_\Phi,\,\omega}(\lambda a)(x) &\lesssim |\lambda||Q|^{\frac{s_0+n+1}{n}}[\omega(Q)\rho(\omega(Q))]^{-1}\\ &\quad\times|x-x_0|^{-(s_0+n+1)}\chi_{B(x_0,2\sqrt{n})}(x), \qquad (3.36)\end{aligned}$$

where $s_0\equiv\lfloor n(\frac{q_\omega}{p_\Phi}-1)\rfloor$. Indeed, for any $\psi\in\mathcal{D}^0_N(\mathbb{R}^n)$ and $t\in(0,1)$, let P be the Taylor expansion of ψ about $(x-x_0)/t$ with degree s_0. By Taylor's remainder theorem, for any $y\in\mathbb{R}^n$, we have

$$\begin{aligned}&\left|\psi\left(\frac{x-y}{t}\right)-P\left(\frac{x-y}{t}\right)\right|\\ &\quad\lesssim \sum_{\substack{\alpha\in\mathbb{Z}^n_+\\ |\alpha|=s_0+1}}\left|(\partial^\alpha\psi)\left(\frac{\theta(x-y)+(1-\theta)(x-x_0)}{t}\right)\right|\left|\frac{x_0-y}{t}\right|^{s_0+1},\end{aligned}$$

where $\theta\in(0,1)$. By $t\in(0,1)$ and $x\in\widetilde{Q}^{\complement}$, we see that $\mathrm{supp}\,(a*\psi_t)\subset B(x_0,2\sqrt{n})$ and that $a*\psi_t(x)\neq0$ implies that $t>\frac{|x-x_0|}{2}$. Thus, from the above facts, Definition 3.4 and (2.1), it follows that for all $x\in\widetilde{Q}^{\complement}$,

$$\begin{aligned}|a*\psi_t(x)| &\le \frac{1}{t^n}\left\{\int_Q|a(y)|\left|\psi\left(\frac{x-y}{t}\right)-P\left(\frac{x-y}{t}\right)\right|dy\right\}\chi_{B(x_0,2\sqrt{n})}(x)\\ &\lesssim |x-x_0|^{-(s_0+n+1)}\left\{\int_Q|a(y)||x_0-y|^{s_0+1}\,dy\right\}\chi_{B(x_0,2\sqrt{n})}(x)\\ &\lesssim |Q|^{\frac{s_0+1}{n}}\|a\|_{L^q_\omega(\mathbb{R}^n)}\left(\int_Q[\omega(y)]^{-q'/q}\,dy\right)^{1/q'}|x-x_0|^{-(s_0+n+1)}\chi_{B(x_0,2\sqrt{n})}(x)\\ &\lesssim |Q|^{\frac{s_0+n+1}{n}}[\omega(Q)\rho(\omega(Q))]^{-1}|x-x_0|^{-(s_0+n+1)}\chi_{B(x_0,2\sqrt{n})}(x),\end{aligned}$$

which together with the arbitrariness of $\psi\in\mathcal{D}^0_N(\mathbb{R}^n)$ implies (3.36). Thus, the claim holds.

Let $Q_k\equiv2^k\sqrt{n}Q$ for all $k\in\mathbb{N}$ and $k_0\in\mathbb{N}$ satisfy $2^{k_0}r\le4<2^{k_0+1}r$. By

$$s_0=\left\lfloor n\left(\frac{q_\omega}{p_\Phi}-1\right)\right\rfloor,$$

we know that there exists $q_0 \in (q_\omega, \infty)$ such that $p_\Phi(s_0+n+1) > nq_0$. From Lemma 2.3, it follows that there exists an $\overline{\omega} \in A_{p_0}(\mathbb{R}^n)$ such that $\omega = \overline{\omega}$ on $Q(x_0, 8\sqrt{n})$. By this fact, (3.36), the lower type p_Φ property of Φ and Lemma 2.3(viii), we conclude that

$$\begin{aligned}
\mathrm{I}_2 &\le \int_{\sqrt{n}r\le|x-x_0|<2\sqrt{n}} \Phi\left(\mathcal{G}^0_{N_{\Phi,\,\omega}}(\lambda a)(x)\right)\omega(x)\,dx \\
&\lesssim \int_{\sqrt{n}r\le|x-x_0|<2\sqrt{n}} \Phi\left(|\lambda||Q|^{\frac{s_0+n+1}{n}}\left[\omega(Q)\rho(\omega(Q))\right]^{-1}|x-x_0|^{-(s_0+n+1)}\right)\overline{\omega}(x)\,dx \\
&\lesssim \sum_{k=1}^{k_0}\int_{Q_{k+1}\setminus Q_k} \Phi\left(|\lambda|2^{-k(s_0+n+1)}\left[\omega(Q)\rho(\omega(Q))\right]^{-1}\right)\overline{\omega}(x)\,dx \\
&\lesssim \sum_{k=1}^{k_0} 2^{-k(s_0+n+1)p_\Phi}\overline{\omega}(Q_{k+1})\Phi\left(\frac{|\lambda|}{\omega(Q)\rho(\omega(Q))}\right) \\
&\lesssim \sum_{k=1}^{k_0} 2^{-k[(s_0+n+1)p_\Phi-nq_0]}\omega(Q)\Phi\left(\frac{|\lambda|}{\omega(Q)\rho(\omega(Q))}\right) \\
&\lesssim \omega(Q)\Phi\left(\frac{|\lambda|}{\omega(Q)\rho(\omega(Q))}\right),
\end{aligned}$$

which together with (3.34) and (3.35) implies (3.33) in Case 1).

Case 2) $|Q| \ge 1$. In this case, let $Q^* \equiv Q(x_0, r+2)$. Thus, from

$$\mathrm{supp}\left(\mathcal{G}^0_{N_{\Phi,\,\omega}}(\lambda a)\right) \subset Q^*,$$

Jensen's inequality, Hölder's inequality, Lemma 2.3(v), and Proposition 3.2(ii), we deduce that

$$\begin{aligned}
&\int_{\mathbb{R}^n} \Phi\left(\mathcal{G}^0_{N_{\Phi,\,\omega}}(\lambda a)(x)\right)\omega(x)\,dx \\
&\quad= \int_{Q^*} \Phi\left(\mathcal{G}^0_{N_{\Phi,\,\omega}}(\lambda a)(x)\right)\omega(x)\,dx \le \omega(Q^*)\Phi\left(\frac{1}{\omega(Q^*)}\int_{Q^*}\mathcal{G}^0_{N_{\Phi,\,\omega}}(\lambda a)(x)\omega(x)\,dx\right) \\
&\quad\le \omega(Q^*)\Phi\left(\frac{1}{[\omega(Q^*)]^{1/q}}\left\{\int_{Q^*}\left[\mathcal{G}^0_{N_{\Phi,\,\omega}}(\lambda a)(x)\right]^q\omega(x)\,dx\right\}^{1/q}\right) \\
&\quad\lesssim \omega(Q^*)\Phi\left(\frac{|\lambda|}{[\omega(Q^*)]^{1/q}}\|a\|_{L^q_\omega(\mathbb{R}^n)}\right) \lesssim \omega(Q^*)\Phi\left(\frac{|\lambda|}{\omega(Q)\rho(\omega(Q))}\right) \\
&\quad\lesssim \omega(Q)\Phi\left(\frac{|\lambda|}{\omega(Q)\rho(\omega(Q))}\right),
\end{aligned}$$

which proves (3.33) in Case 2). This finishes the proof of Theorem 3.18. ■

4. Calderón-Zygmund decompositions

In this section, we establish some subtle estimates for the Calderón-Zygmund decomposition associated with local grand maximal functions on the weighted Euclidean space

$\mathbb{R}^n$ given in [49]. Notice that the construction of the Calderón-Zygmund decomposition in [49] is similar to those in [46, 3, 5].

Let Φ be a positive function on $\mathbb{R}_+$ satisfying Assumption (A), $\omega \in A_\infty^{\rm loc}(\mathbb{R}^n)$ and q_ω be as in (2.4). Let integer $N \geq 2$, $\mathcal{G}_N(f)$ and $\mathcal{G}_N^0(f)$ be as in (3.2).

Throughout this section, let $f \in \mathcal{D}'(\mathbb{R}^n)$ satisfy that for all $\lambda \in (0, \infty)$,

$$\omega\left(\{x \in \mathbb{R}^n : \mathcal{G}_N(f)(x) > \lambda\}\right) < \infty.$$

For a given $\lambda > \inf_{x \in \mathbb{R}^n} \mathcal{G}_N(f)(x)$, we set

$$\Omega_\lambda \equiv \{x \in \mathbb{R}^n : \mathcal{G}_N(f)(x) > \lambda\}. \tag{4.1}$$

It is obvious that Ω_λ is a proper open subset of $\mathbb{R}^n$. First, we recall the usual Whitney decomposition of Ω_λ given in [49] (see also [46, 3, 5]). We can find closed cubes $\{Q_i\}_i$ such that

$$\Omega_\lambda = \bigcup_i Q_i, \tag{4.2}$$

their interiors are away from $\Omega_\lambda^\complement$ and

$$\mathrm{diam}\,(Q_i) \leq 2^{-(6+n)}\,\mathrm{dist}\,(Q_i, \Omega_\lambda^\complement) \leq 4\,\mathrm{diam}\,(Q_i).$$

In what follows, fix $a \equiv 1 + 2^{-(11+n)}$ and denote aQ_i by Q_i^* for all i. Then we have $Q_i \subset Q_i^*$. Moveover, $\Omega_\lambda = \cup_i Q_i^*$, and $\{Q_i^*\}_i$ have the bounded interior property, namely, every point in Ω_λ is contained in at most a fixed number of $\{Q_i^*\}_i$.

Now we take a function $\xi \in \mathcal{D}(\mathbb{R}^n)$ such that $0 \leq \xi \leq 1$, $\mathrm{supp}\,(\xi) \subset aQ(0,1)$ and $\xi \equiv 1$ on $Q(0,1)$. For $x \in \mathbb{R}^n$, set $\xi_i(x) \equiv \xi((x - x_k)/l_i)$, where and in what follows, x_i is the *center* of the cube Q_i and l_i its *sidelength.* Obviously, by the construction of $\{Q_i^*\}_i$ and $\{\xi_i\}_i$, for any $x \in \mathbb{R}^n$, we have $1 \leq \sum_i \xi_i(x) \leq L$, where L is a fixed positive integer independent of x. Let

$$\zeta_i \equiv \frac{\xi_i}{\sum_j \xi_j}. \tag{4.3}$$

Then $\{\zeta_i\}_i$ form a smooth partition of unity for Ω_λ subordinate to the locally finite cover $\{Q_i^*\}_i$ of Ω_λ, namely, $\chi_{\Omega_\lambda} = \sum_k \zeta_k$ with each $\zeta_i \in \mathcal{D}(\mathbb{R}^n)$ supported in Q_i^*.

Let $s \in \mathbb{Z}_+$ be some fixed integer and $\mathcal{P}_s(\mathbb{R}^n)$ denote the linear space of polynomials in n variables of degrees no more than s. For each $i \in \mathbb{N}$ and $P \in \mathcal{P}_s(\mathbb{R}^n)$, set

$$\|P\|_i \equiv \left[\frac{1}{\int_{\mathbb{R}^n} \zeta_i(y)\,dy} \int_{\mathbb{R}^n} |P(x)|^2 \zeta_i(x)\,dx\right]^{1/2}. \tag{4.4}$$

Then it is easy to know that $(\mathcal{P}_s(\mathbb{R}^n), \|\cdot\|_i)$ is a finite dimensional Hilbert space. Let $f \in \mathcal{D}'(\mathbb{R}^n)$. Since f induces a linear functional on $\mathcal{P}_s(\mathbb{R}^n)$ via

$$P \mapsto \frac{1}{\int_{\mathbb{R}^n} \zeta_i(y)\,dy} \langle f, P\zeta_i \rangle,$$

by the Riesz represent theorem, there exists a unique polynomial

$$P_i \in \mathcal{P}_s(\mathbb{R}^n) \tag{4.5}$$

for each i such that for all $P \in \mathcal{P}_s(\mathbb{R}^n)$, $\langle f, P\zeta_i\rangle = \langle P_i, P\zeta_i\rangle$. For each i, define the distribution

$$b_i \equiv (f - P_i)\zeta_i \text{ when } l_i \in (0,1), \text{ and } b_i \equiv f\zeta_i \text{ when } l_i \in [1,\infty). \tag{4.6}$$

We show that for suitable choices of s and N, the series $\sum_i b_i$ converge in $\mathcal{D}'(\mathbb{R}^n)$, and in this case, we let $g \equiv f - \sum_i b_i$ in $\mathcal{D}'(\mathbb{R}^n)$. We point out that the represent

$$f = g + \sum_i b_i, \tag{4.7}$$

where g and b_i are as above, is called a *Calderón-Zygmund decomposition* of f of degree s and height λ associated with $\mathcal{G}_N(f)$.

The rest of this section consists of a series of lemmas. Lemma 4.1 gives a property of the smooth partition of unity $\{\zeta_i\}_i$, Lemmas 4.2 through 4.5 are devoted to some estimates for the bad parts $\{b_i\}_i$, and Lemmas 4.6 and 4.7 give some controls over the good part g. Finally, Corollary 4.8 shows that the density of $L^q_\omega(\mathbb{R}^n) \cap h^\Phi_{\omega,N}(\mathbb{R}^n)$ in $h^\Phi_{\omega,N}(\mathbb{R}^n)$, where $q \in (q_\omega, \infty)$. The following Lemmas 4.1 through 4.3, and Lemmas 4.5 and 4.6 are respectively Lemmas 4.2 through 4.5, and Lemmas 4.7 and 4.8 in [49].

LEMMA 4.1. *There exists a positive constant C_1 such that for all $f \in \mathcal{D}'(\mathbb{R}^n)$,*

$$\lambda > \inf_{x\in\mathbb{R}^n} \mathcal{G}_N(f)(x)$$

and $l_i \in (0,1)$,

$$\sup_{y\in\mathbb{R}^n} |P_i(y)\zeta_i(y)| \le C_1\lambda.$$

LEMMA 4.2. *There exists a positive constant C_2 such that for all $i \in \mathbb{N}$ and $x \in Q_i^*$,*

$$\mathcal{G}^0_N(b_i)(x) \le C_2\mathcal{G}_N(f)(x). \tag{4.8}$$

LEMMA 4.3. *Assume that integers s and N satisfy $0 \le s < N$ and $N \ge 2$. Then there exist positive constants C, C_3 and C_4 such that for all $i \in \mathbb{N}$ and $x \in (Q_i^*)^\complement$,*

$$\mathcal{G}^0_N(b_i)(x) \le C\frac{\lambda l_i^{n+s+1}}{(l_i + |x - x_i|)^{n+s+1}}\chi_{B(x_i, C_3)}(x), \tag{4.9}$$

where x_i is the center of the cube Q_i. Moreover, if $x \in (Q_i^*)^\complement$ and $l_i \in [C_4, \infty)$, then $\mathcal{G}^0_N(b_i)(x) = 0$.

LEMMA 4.4. *Let Φ satisfy Assumption* (A), *$\omega \in A^{\rm loc}_\infty(\mathbb{R}^n)$, q_ω and p_Φ be respectively as in* (2.4) *and* (2.6). *If integers $s \ge \lfloor n(q_\omega/p_\Phi - 1)\rfloor$, $N > s$ and $N \ge N_{\Phi,\omega}$, where $N_{\Phi,\omega}$ is as in* (3.25). *Then there exists a positive constant C_5 such that for all $f \in h^\Phi_{\omega,N}(\mathbb{R}^n)$, $\lambda > \inf_{x\in\mathbb{R}^n} \mathcal{G}_N(f)(x)$ and $i \in \mathbb{N}$,*

$$\int_{\mathbb{R}^n} \Phi\left(\mathcal{G}^0_N(b_i)(x)\right)\omega(x)\,dx \le C_5\int_{Q_i^*} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx. \tag{4.10}$$

Moreover, the series $\sum_i b_i$ converges in $h^\Phi_{\omega,N}(\mathbb{R}^n)$ and

$$\int_{\mathbb{R}^n} \Phi\left(\mathcal{G}^0_N\left(\sum_i b_i\right)(x)\right)\omega(x)\,dx \le C_5\int_{\Omega_\lambda} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx. \tag{4.11}$$

Proof. By Lemmas 4.2 and 4.3, we have

$$\int_{\mathbb{R}^n} \Phi\left(\mathcal{G}_N^0(b_i)(x)\right)\omega(x)\,dx \lesssim \int_{Q_i^*} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx + \int_{(2C_3Q_i^0)\setminus Q_i^*} \Phi\left(\mathcal{G}_N^0(b_i)(x)\right)\omega(x)\,dx, \tag{4.12}$$

where $Q_i^0 \equiv Q(x_i, 1)$. Notice that $s \geq \lfloor n(q_\omega/p_\Phi - 1)\rfloor$ implies $(s+n+1)p_\Phi > nq_\omega$. Thus, we take $q_0 \in (q_\omega, \infty)$ such that $(s+n+1)p_\Phi > nq_0$ and $\omega \in A_{q_0}^{\mathrm{loc}}(\mathbb{R}^n)$. By Lemma 2.3(i), we know that there exists an $\widetilde{\omega} \in A_{q_0}(\mathbb{R}^n)$ such that $\widetilde{\omega} = \omega$ on $2C_3Q_i^0$ and $A_{q_0}(\widetilde{\omega}) \lesssim A_{q_0}^{\mathrm{loc}}(\omega)$. Using Lemma 4.3, the lower p_Φ property of Φ, Lemma 2.3(viii) and the fact that $\mathcal{G}_N(f) > \lambda$ for all $x \in Q_i^*$, we conclude that

$$\begin{aligned}
&\int_{(2C_3Q_i^0)\setminus Q_i^*} \Phi\left(\mathcal{G}_N^0(b_i)(x)\right)\omega(x)\,dx \\
&\quad\leq \sum_{k=1}^{k_0} \int_{2^kQ_i^*\setminus 2^{k-1}Q_i^*} \Phi\left(\mathcal{G}_N^0(b_i)(x)\right)\widetilde{\omega}(x)\,dx \lesssim \sum_{k=1}^{k_0} \Phi\left(\frac{\lambda}{2^{k(n+s+1)}}\right)\int_{2^kQ_i^*} \widetilde{\omega}(x)\,dx \\
&\quad\lesssim \sum_{k=1}^{k_0} \Phi(\lambda)\frac{1}{2^{k(n+s+1)p_\Phi}}\int_{2^kQ_i^*} \widetilde{\omega}(x)\,dx \lesssim \sum_{k=1}^{k_0} \Phi(\lambda)2^{-k[(n+s+1)p_\Phi - nq_0]}\widetilde{\omega}(Q_i^*) \\
&\quad\lesssim \int_{Q_i^*} \Phi\left(\mathcal{G}_N(f)(x)\right)\widetilde{\omega}(x)\,dx \sim \int_{Q_i^*} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx,
\end{aligned} \tag{4.13}$$

where $k_0 \in \mathbb{N}$ satisfies $2^{k_0-2} \leq C_3 < 2^{k_0-1}$. From (4.12) and (4.13), we deduce that (4.10) holds. Then, by (4.10), we see that

$$\begin{aligned}
\sum_i \int_{\mathbb{R}^n} \Phi\left(\mathcal{G}_N^0(b_i)(x)\right)\omega(x)\,dx &\lesssim \sum_i \int_{Q_i^*} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx \\
&\lesssim \int_{\Omega_\lambda} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx.
\end{aligned}$$

Combining the above inequality with the completeness of $h_{\omega,N}^{\Phi}(\mathbb{R}^n)$, we infer that $\sum_i b_i$ converges in $h_{\omega,N}^{\Phi}(\mathbb{R}^n)$. So by Proposition 3.16, the series $\sum_i b_i$ converges in $\mathcal{D}'(\mathbb{R}^n)$ and hence

$$\mathcal{G}_N^0\left(\sum_i b_i\right)(x) \leq \sum_i \mathcal{G}_N^0(b_i)(x)$$

for all $x \in \mathbb{R}^n$, which gives (4.11). This finishes the proof of Lemma 4.4. ■

LEMMA 4.5. *Let $\omega \in A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$ and q_ω be as in* (2.4)*, $s \in \mathbb{Z}_+$ and integer $N \geq 2$. If $q \in (q_\omega, \infty]$ and $f \in L_\omega^q(\mathbb{R}^n)$, then the series $\sum_i b_i$ converges in $L_\omega^q(\mathbb{R}^n)$ and there exists a positive constant C_6, independent of f, such that*

$$\left\|\sum_i |b_i|\right\|_{L_\omega^q(\mathbb{R}^n)} \leq C_6\|f\|_{L_\omega^q(\mathbb{R}^n)}.$$

LEMMA 4.6. *Let integers s and N satisfy $0 \leq s < N$ and $N \geq 2$, $f \in \mathcal{D}'(\mathbb{R}^n)$ and $\lambda > \inf_{x\in\mathbb{R}^n} \mathcal{G}_N(f)(x)$. If $\sum_i b_i$ converges in $\mathcal{D}'(\mathbb{R}^n)$, then there exists a positive constant*

C_7, independent of f and λ, such that for all $x \in \mathbb{R}^n$,

$$\mathcal{G}_N^0(g)(x) \le \mathcal{G}_N^0(f)(x)\chi_{\Omega_\lambda^\complement}(x) + C_7\lambda \sum_i \frac{l_i^{n+s+1}}{(l_i + |x-x_i|)^{n+s+1}}\chi_{B(x_i,C_3)}(x),$$

where x_i is the center of Q_i and C_3 is as in Lemma 4.3.

Lemma 4.7. *Let Φ satisfy Assumption* (A), *$\omega \in A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$, q_ω and p_Φ be respectively as in* (2.4) *and* (2.6), *integer $N \ge N_{\Phi,\,\omega}$, where $N_{\Phi,\,\omega}$ is as in* (3.25), *and $q \in (q_\omega, \infty)$.*

(i) *If integers s and N satisfy $N > s \ge \lfloor n(q_\omega/p_\Phi - 1)\rfloor$ and $f \in h_{\omega,\,N}^\Phi(\mathbb{R}^n)$, then $\mathcal{G}_N^0(g) \in L_\omega^q(\mathbb{R}^n)$ and there exists a positive constant C_8, independent of f and λ, such that*

$$\int_{\mathbb{R}^n} \left[\mathcal{G}_N^0(g)(x)\right]^q \omega(x)\,dx \le C_8 \begin{cases} \lambda^{q-1}\int_{\mathbb{R}^n} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx, & \lambda \in (0,1), \\ \lambda^{q-p_\Phi}\int_{\mathbb{R}^n} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx, & \lambda \in [1,\infty). \end{cases} \tag{4.14}$$

(ii) *If $f \in L_\omega^q(\mathbb{R}^n)$, then $g \in L_\omega^\infty(\mathbb{R}^n)$ and there exists a positive constant C_9, independent of f and λ, such that $\|g\|_{L_\omega^\infty(\mathbb{R}^n)} \le C_9\lambda$.*

Proof. We first prove (i). Let $f \in h_{\omega,\,N}^\Phi(\mathbb{R}^n)$. By Lemma 4.4 and Proposition 3.16, $\sum_i b_i$ converges in both $h_{\omega,\,N}^\Phi(\mathbb{R}^n)$ and $\mathcal{D}'(\mathbb{R}^n)$. By $s \ge \lfloor n(q_\omega/p_\Phi - 1)\rfloor$, we know that there exists $q_0 \in (q_\omega, \infty)$ such that $(s+n+1)p_\Phi > nq_0$ and $\omega \in A_{q_0}^{\mathrm{loc}}(\mathbb{R}^n)$. Let

$$\mathrm{J} \equiv \int_{\Omega_\lambda^\complement} [\mathcal{G}_N(f)(x)]^q \omega(x)\,dx.$$

From Lemmas 4.6 and 3.10, we infer that

$$\begin{aligned}
\int_{\mathbb{R}^n} \left[\mathcal{G}_N^0(g)(x)\right]^q \omega(x)\,dx &\lesssim \lambda^q \int_{\mathbb{R}^n} \left[\sum_i \frac{l_i^{n+s+1}}{(l_i+|x-x_i|)^{n+s+1}}\chi_{B(x_i,C_3)}(x)\right]^q \omega(x)\,dx + \mathrm{J} \\
&\lesssim \lambda^q \int_{\mathbb{R}^n} \left(\sum_i \left[M_{2C_3}^{\mathrm{loc}}(\chi_{Q_i})(x)\right]^{(n+s+1)/n}\right)^q \omega(x)\,dx + \mathrm{J} \\
&\lesssim \lambda^q \int_{\mathbb{R}^n} \left(\sum_i [\chi_{Q_i}(x)]^{(n+s+1)/n}\right)^q \omega(x)\,dx + \mathrm{J} \\
&\lesssim \lambda^q \int_{\Omega_\lambda} \omega(x)\,dx + \mathrm{J} \sim \lambda^q \omega(\Omega_\lambda) + \mathrm{J}.
\end{aligned}$$

Now, we consider the following two cases for λ.

Case 1) $\lambda \ge 1$. In this case, since Φ has lower type p_Φ, we have

$$\begin{aligned}
\lambda^q\omega(\Omega_\lambda) &\le \lambda^{q-p_\Phi}\omega(\Omega_\lambda)\left[\inf_{x\in\Omega_\lambda} \mathcal{G}_N(f)(x)\right]^{p_\Phi} \le \lambda^{q-p_\Phi}\omega(\Omega_\lambda)\Phi\left(\inf_{x\in\Omega_\lambda} \mathcal{G}_N(f)(x)\right) \\
&\le \lambda^{q-p_\Phi}\int_{\Omega_\lambda} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx.
\end{aligned}$$

Recall that

$$\Omega_1 \equiv \{x \in \mathbb{R}^n : \mathcal{G}_N(f)(x) > 1\}.$$

From the fact that Φ has lower type p_Φ and upper type 1, it follows that

$$\begin{aligned}\mathrm{J} &= \int_{\Omega_\lambda^\complement\cap\Omega_1} [\mathcal{G}_N(f)(x)]^q\,\omega(x)\,dx + \int_{\Omega_\lambda^\complement\cap\Omega_1^\complement} \cdots\\ &\lesssim \int_{\Omega_\lambda^\complement} [\mathcal{G}_N(f)(x)]^{q-p_\Phi}\,\Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx + \int_{\Omega_\lambda^\complement} [\mathcal{G}_N(f)(x)]^{q-1}\,\Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx\\ &\lesssim \left(\lambda^{q-p_\Phi}+\lambda^{q-1}\right)\int_{\Omega_\lambda^\complement} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx \lesssim \lambda^{q-p_\Phi}\int_{\Omega_\lambda^\complement} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx,\end{aligned}$$

which together with the estimate of $\lambda^q\omega(\Omega_\lambda)$ implies (4.10) in Case 1).

Case 2) $\lambda\in(0,1)$. In this case, for any $x\in\Omega_\lambda$, if $\mathcal{G}_N(f)(x)\ge 1>\lambda$, using the fact that Φ has lower type p_Φ, we conclude that

$$\lambda^q \le \lambda^{q-p_\Phi}[\mathcal{G}_N(f)(x)]^{p_\Phi} \lesssim \lambda^{q-p_\Phi}\Phi\left(\mathcal{G}_N(f)(x)\right) \lesssim \lambda^{q-1}\Phi\left(\mathcal{G}_N(f)(x)\right).$$

If $\mathcal{G}_N(f)(x)<1$ and $\mathcal{G}_N(f)(x)>\lambda$, by the fact that Φ has upper type 1, we see that

$$\lambda^q \le \lambda^{q-1}\mathcal{G}_N(f)(x) \lesssim \lambda^{q-1}\Phi\left(\mathcal{G}_N(f)(x)\right).$$

From these estimates, we deduce that

$$\lambda^q\omega(\Omega_\lambda) \lesssim \lambda^{q-1}\int_{\Omega_\lambda} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx.$$

For J, by $\lambda\in(0,1)$, $\mathcal{G}_N(f)(x)\le\lambda$ for all $x\in\Omega_\lambda^\complement$ and the fact that Φ has upper type 1, we know that

$$\mathrm{J} \le \lambda^{q-1}\int_{\Omega_\lambda^\complement} \mathcal{G}_N(f)(x)\omega(x)\,dx \lesssim \lambda^{q-1}\int_{\Omega_\lambda^\complement} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx,$$

which together with the estimate of $\lambda^q\omega(\Omega_\lambda)$ implies (4.14) in Case 2). Thus, (i) holds.

Now we prove (ii). If $f\in L^q_\omega(\mathbb{R}^n)$, then g and $\{b_i\}_i$ are functions. By Lemma 4.5, we know that $\sum_i b_i$ converges in $L^q_\omega(\mathbb{R}^n)$ and hence in $\mathcal{D}'(\mathbb{R}^n)$ by Lemma 2.6(ii). Write

$$g = f - \sum_i b_i = f\left(1-\sum_i \zeta_i\right) + \sum_{i\in F} P_i\zeta_i = f\chi_{\Omega_\lambda^\complement} + \sum_{i\in F} P_i\zeta_i,$$

where $F\equiv\{i\in\mathbb{N}:\ l_i\in(0,1)\}$. By Lemma 4.1, we have that $|g(x)|\lesssim\lambda$ for all $x\in\Omega_\lambda$, which combined with Proposition 3.2(i) yields that

$$|g(x)| = |f(x)| \le \mathcal{G}_N(f)(x) \le \lambda$$

for almost every $x\in\Omega_\lambda^\complement$, Thus, $\|g\|_{L^\infty_\omega(\mathbb{R}^n)}\lesssim\lambda$. This shows (ii) and hence finishes the proof of Lemma 4.7. ■

Corollary 4.8. *Let Φ satisfy Assumption* (A), $\omega\in A^{\mathrm{loc}}_\infty(\mathbb{R}^n)$, q_ω *be as in* (2.4),

$$q\in(q_\omega,\infty)$$

and integer $N\ge N_{\Phi,\,\omega}$, where $N_{\Phi,\,\omega}$ is as in (3.25). *Then $h^\Phi_{\omega,\,N}(\mathbb{R}^n)\cap L^q_\omega(\mathbb{R}^n)$ is dense in $h^\Phi_{\omega,\,N}(\mathbb{R}^n)$.*

Proof. Let $f\in h^\Phi_{\omega,\,N}(\mathbb{R}^n)$. For any $\lambda>\inf_{x\in\mathbb{R}^n}\mathcal{G}_N(f)(x)$, let

$$f = g^\lambda + \sum_i b_i^\lambda$$

be the Calderón-Zygmund decomposition of f of degree s with $\lfloor n(q_\omega/p_\Phi-1)\rfloor \le s < N$ and height λ associated to $\mathcal{G}_N(f)$. By Lemma 4.4,

$$\int_{\mathbb{R}^n} \Phi\left(\mathcal{G}_N^0\left(\sum_i b_i^\lambda\right)(x)\right)\omega(x)\,dx \lesssim \int_{\{x\in\mathbb{R}^n:\,\mathcal{G}_N(f)(x)>\lambda\}} \Phi\left(\mathcal{G}_N(f)(x)\right)\omega(x)\,dx.$$

Therefore, $g^\lambda \to f$ in $h^\Phi_{\omega,\,N}(\mathbb{R}^n)$ as $\lambda\to\infty$. Moreover, by Lemma 4.7(i), we have

$$\mathcal{G}_N^0(g^\lambda) \in L^q_\omega(\mathbb{R}^n),$$

which together with Proposition 3.2(ii) implies $g^\lambda \in L^q_\omega(\mathbb{R}^n)$. This finishes the proof of Corollary 4.8. ■

5. Weighted atomic decompositions of $h^\Phi_{\omega,\,N}(\mathbb{R}^n)$

In this section, we establish the equivalence between $h^\Phi_{\omega,\,N}(\mathbb{R}^n)$ and $h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)$ by using the Calderón-Zygmund decomposition associated to the local grand maximal function stated in Section 4.

Let Φ satisfy Assumption (A), $\omega \in A^{\mathrm{loc}}_\infty(\mathbb{R}^n)$, q_ω, p_Φ and $N_{\Phi,\,\omega}$ be respectively as in (2.4), (2.6) and (3.25), integer $N \ge N_{\Phi,\,\omega}$ and $s_0 \equiv \lfloor n(q_\omega/p_\Phi-1)\rfloor$. *Throughout this section, let*

$$f \in h^\Phi_{\omega,\,N}(\mathbb{R}^n).$$

We take $k_0\in\mathbb{Z}$ such that $2^{k_0-1} \le \inf_{x\in\mathbb{R}^n}\mathcal{G}_N(f)(x) < 2^{k_0}$ when

$$\inf_{x\in\mathbb{R}^n}\mathcal{G}_N(f)(x) > 0,$$

and when $\inf_{x\in\mathbb{R}^n}\mathcal{G}_N(f)(x)=0$, let $k_0\equiv-\infty$. *Throughout the whole section, we always assume that* $k\ge k_0$. For each integer $k\ge k_0$, consider the Calderón-Zygmund decomposition of f of degree s and height $\lambda=2^k$ associated to $\mathcal{G}_N(f)$. Namely, for any $k\ge k_0$, by taking $\lambda\equiv 2^k$ in (4.1), we now write the Calderón-Zygmund decomposition in (4.7) by

$$f = g^k + \sum_i b_i^k, \tag{5.1}$$

where and in what follows of this section, we write $\{Q_i\}_i$ in (4.2), $\{\zeta_i\}_i$ in (4.3), $\{P_i\}_i$ in (4.5) and $\{b_i\}_i$ in (4.6), respectively, as $\{Q_i^k\}_i$, $\{\zeta_i^k\}_i$, $\{P_i^k\}_i$ and $\{b_i^k\}_i$. Now, the *center* and the *sidelength* of Q_i^k is respectively denoted by x_i^k and l_i^k. Recall that for all i and k,

$$\sum_i \zeta_i^k = \chi_{\Omega_{2^k}},\ \ \mathrm{supp}\,(b_i^k)\subset \mathrm{supp}\,(\zeta_i^k)\subset Q_i^{k*}, \tag{5.2}$$

$\{Q_i^{k*}\}_i$ has the bounded interior property, and for all $P\in\mathcal{P}_s(\mathbb{R}^n)$,

$$\langle f, P\zeta_i^k\rangle = \langle P_i^k, P\zeta_i^k\rangle. \tag{5.3}$$

For each integer $k\ge k_0$ and $i,\,j\in\mathbb{N}$, let $P^{k+1}_{i,\,j}$ be the orthogonal projection of $(f-P_j^{k+1})\zeta_i^k$ on $\mathcal{P}_s(\mathbb{R}^n)$ with respect to the norm

$$\|P\|_j^2 \equiv \frac{1}{\int_{\mathbb{R}^n}\zeta_j^{k+1}(y)\,dy}\int_{\mathbb{R}^n}|P(x)|^2\zeta_j^{k+1}(x)\,dx,$$

namely, $P_{i,j}^{k+1}$ is the unique polynomial of $\mathcal{P}_s(\mathbb{R}^n)$ such that for any $P \in \mathcal{P}_s(\mathbb{R}^n)$,

$$\langle (f - P_j^{k+1})\zeta_i^k, P\zeta_j^{k+1}\rangle = \int_{\mathbb{R}^n} P_{i,j}^{k+1}(x)P(x)\zeta_j^{k+1}(x)\,dx. \tag{5.4}$$

Recall that $a \equiv 1 + 2^{-(11+n)}$. In what follows, let $Q_i^{k*} \equiv aQ_i^k$,

$$E_1^k \equiv \left\{i \in \mathbb{N}: \ |Q_i^k| \ge 1/(2^4 n)\right\},$$

$$E_2^k \equiv \left\{i \in \mathbb{N}: \ |Q_i^k| < 1/(2^4 n)\right\},$$

$$F_1^k \equiv \left\{i \in \mathbb{N}: \ |Q_i^k| \ge 1\right\}$$

and

$$F_2^k \equiv \left\{i \in \mathbb{N}: \ |Q_i^k| < 1\right\}.$$

Observe that

$$P_{i,j}^{k+1} \ne 0 \text{ if and only if } Q_i^{k*} \cap Q_j^{(k+1)*} \ne \emptyset. \tag{5.5}$$

Indeed, this follows directly from the definition of $P_{i,j}^{k+1}$. The following Lemmas 5.1 through 5.3 are just Lemmas 5.1 through 5.3 in [49].

Lemma 5.1. *Let Ω_{2^k} be as in* (4.1) *with $\lambda = 2^k$, Q_i^{k*} and l_i^k be as above.*

(i) *If $Q_i^{k*} \cap Q_j^{(k+1)*} \ne \emptyset$, then $l_j^{k+1} \le 2^4\sqrt{n}\, l_i^k$ and $Q_j^{(k+1)*} \subset 2^6 n Q_i^{k*} \subset \Omega_{2^k}$.*

(ii) *There exists a positive integer L such that for each $i \in \mathbb{N}$, the cardinality of $\{j \in \mathbb{N}: \ Q_i^{k*} \cap Q_j^{(k+1)*} \ne \emptyset\}$ is bounded by L.*

Lemma 5.2. *There exists a positive constant C such that for all $i, j \in \mathbb{N}$ and integer $k \ge k_0$ with $l_j^{k+1} \in (0,1)$,*

$$\sup_{y\in\mathbb{R}^n} \left|P_{i,j}^{k+1}(y)\zeta_j^{k+1}(y)\right| \le C2^{k+1}. \tag{5.6}$$

Lemma 5.3. *For any $k \in \mathbb{Z}$ with $k \ge k_0$,*

$$\sum_{i\in\mathbb{N}} \left(\sum_{j \in F_2^{k+1}} P_{i,j}^{k+1}\zeta_j^{k+1} \right) = 0,$$

where the series converges both in $\mathcal{D}'(\mathbb{R}^n)$ and pointwise.

The following lemma gives the weighted atomic decomposition for a dense subspace of $h_{\omega,N}^{\Phi}(\mathbb{R}^n)$.

Lemma 5.4. *Let Φ satisfy Assumption* (A), *$\omega \in A_\infty^{\mathrm{loc}}(\mathbb{R}^n)$, q_ω, p_Φ and $N_{\Phi,\omega}$ be respectively as in* (2.4), (2.6) *and* (3.25). *If $q \in (q_\omega, \infty)$, integers $N \ge N_{\Phi,\omega}$, $s \ge \lfloor n(q_\omega/p_\Phi - 1)\rfloor$ and $N > s$, then for any $f \in (L_\omega^q(\mathbb{R}^n) \cap h_{\omega,N}^{\Phi}(\mathbb{R}^n))$, there exist $\lambda_0 \in \mathbb{C}$, $\{\lambda_i^k\}_{k\ge k_0, i} \subset \mathbb{C}$, a $(\rho,\infty)_\omega$-single-atom a_0 and $(\rho,\infty,s)_\omega$-atoms $\{a_i^k\}_{k\ge k_0, i}$ such that*

$$f = \sum_{k\ge k_0}\sum_i \lambda_i^k a_i^k + \lambda_0 a_0, \tag{5.7}$$

where the series converges both in $\mathcal{D}'(\mathbb{R}^n)$ and almost everywhere. Moreover, there exists a positive constant C, independent of f, such that

$$\Lambda(\{\lambda_i^k a_i^k\}_{k\ge k_0, i} \cup \{\lambda_0 a_0\}) \le C\|f\|_{h_{\omega,N}^{\Phi}(\mathbb{R}^n)}. \tag{5.8}$$

Proof. Let $f \in (L^q_\omega(\mathbb{R}^n) \cap h^\Phi_{\omega,\,N}(\mathbb{R}^n))$. We first consider the case that $k_0 = -\infty$. As above, for each $k \in \mathbb{Z}$, f has a Calderón-Zygmund decomposition of degree s and height $\lambda = 2^k$ associated to $\mathcal{G}_N(f)$ as in (5.1), namely,

$$f = g^k + \sum_i b^k_i.$$

By Corollary 4.8 and Proposition 3.2, we have that $g^k \to f$ in both $h^\Phi_{\omega,\,N}(\mathbb{R}^n)$ and $\mathcal{D}'(\mathbb{R}^n)$ as $k \to \infty$. By Lemma 4.7(i), $\|g^k\|_{L^q_\omega(\mathbb{R}^n)} \to 0$ as $k \to -\infty$, and furthermore, by Lemma 2.6(ii), $g^k \to 0$ in $\mathcal{D}'(\mathbb{R}^n)$ as $k \to -\infty$. Therefore,

$$f = \sum_{k=-\infty}^{\infty} \left(g^{k+1} - g^k\right) \tag{5.9}$$

in $\mathcal{D}'(\mathbb{R}^n)$. Moreover, since $\operatorname{supp}\left(\sum_i b^k_i\right) \subset \Omega_{2^k}$ and $\omega(\Omega_{2^k}) \to 0$ as $k \to \infty$, then $g^k \to f$ almost everywhere as $k \to \infty$. Thus, (5.9) also holds almost everywhere. By Lemma 5.3 and (5.2) with $\Omega_{2^{k+1}} \subset \Omega_{2^k}$,

$$\begin{aligned} g^{k+1} - g^k &= \left(f - \sum_j b^{k+1}_j\right) - \left(f - \sum_i b^k_i\right) \\ &= \sum_i b^k_i - \sum_j b^{k+1}_j + \sum_i \left(\sum_{j \in F^{k+1}_2} P^{k+1}_{i,\,j} \zeta^{k+1}_j\right) \\ &= \sum_i \left[b^k_i - \sum_j b^{k+1}_j \zeta^k_i + \sum_{j \in F^{k+1}_2} P^{k+1}_{i,\,j} \zeta^{k+1}_j\right] \equiv \sum_i h^k_i, \end{aligned} \tag{5.10}$$

where all the series converge in both $\mathcal{D}'(\mathbb{R}^n)$ and almost everywhere. Furthermore, from the definitions of b^k_j and b^{k+1}_j as in (4.3), we infer that when $l^k_i \in (0, 1)$,

$$h^k_i = f\chi_{\Omega^{\complement}_{2^{k+1}}} \zeta^k_i - P^k_i \zeta^k_i + \sum_{j \in F^{k+1}_2} P^{k+1}_j \zeta^k_i \zeta^{k+1}_j + \sum_{j \in F^{k+1}_2} P^{k+1}_{i,\,j} \zeta^{k+1}_j, \tag{5.11}$$

and when $l^k_i \in [1, \infty)$,

$$h^k_i = f\chi_{\Omega^{\complement}_{2^{k+1}}} \zeta^k_i + \sum_{j \in F^{k+1}_2} P^{k+1}_j \zeta^k_i \zeta^{k+1}_j + \sum_{j \in F^{k+1}_2} P^{k+1}_{i,\,j} \zeta^{k+1}_j. \tag{5.12}$$

By Proposition 3.2(i), we know that for almost every $x \in \Omega^{\complement}_{2^{k+1}}$,

$$|f(x)| \le \mathcal{G}_N(f)(x) \le 2^{k+1},$$

which, together with Lemma 4.1, Lemma 5.1(ii), (5.5), Lemma 5.2, (5.11) and (5.12), implies that there exists a positive constant C_{10} such that for all $i \in \mathbb{N}$,

$$\|h^k_i\|_{L^\infty_\omega(\mathbb{R}^n)} \le C_{10} 2^k. \tag{5.13}$$

Next, we show that for each i and k, h^k_i is a multiple of a $(\rho,\, \infty,\, s)_\omega$-atom by considering the following two cases for i.

Case 1) $i \in E^k_1$. In this case, from the fact that $l^{k+1}_j < 1$ for $j \in F^{k+1}_2$, we deduce that $Q^{(k+1)*}_j \subset Q(x^k_i, a(l^k_i + 2))$ for j satisfying $Q^{k*}_i \cap Q^{(k+1)*}_j \ne \emptyset$. Let $\gamma \equiv 1 + 2^{-12-n}$.

Thus, when $l_i^k \ge 2/(\gamma-1)$, if letting $\widetilde{Q}_i^k \equiv Q(x_i^k, a(l_i^k+2))$, then,

$$\operatorname{supp}(h_i^k) \subset \widetilde{Q}_i^k \subset \gamma Q_i^{k*} \subset \Omega_{2^k}.$$

When $l_i^k < 2/(\gamma-1)$, if letting $\widetilde{Q}_i^k \equiv 2^6 n Q_i^{k*}$, then by Lemma 5.1(i), we have

$$\operatorname{supp}(h_i^k) \subset \widetilde{Q}_i^k \subset \Omega_{2^k}.$$

From the definition of $\widetilde{Q}_i^k$, Lemma 2.3(v) and Remark 2.4 with $\widetilde{C} \equiv 2/(\gamma-1)$, we infer that there exists a positive constant C_{11} such that

$$\omega(\widetilde{Q}_i^k) \le C_{11}\omega(Q_i^{k*}). \tag{5.14}$$

Let $\widetilde{A}_1 \equiv \max\{C_{10}, C_{11}\}$,

$$\lambda_i^k \equiv \widetilde{A}_1 2^k \omega(\widetilde{Q}_i^k)\rho(\omega(\widetilde{Q}_i^k)) \tag{5.15}$$

and $a_i^k \equiv (\lambda_i^k)^{-1} h_i^k$. From (5.13) and $\operatorname{supp}(h_i^k) \subset \widetilde{Q}_i^k$ with $l(\widetilde{Q}_i^k) \ge 2a > 1$, it follows that a_i^k is a $(\rho,\, \infty,\, s)_\omega$-atom.

Case 2) $i \in E_2^k$. In this case, if $j \in F_1^{k+1}$, then $l_i^k < l_j^{k+1}/(2^4 n)$. By Lemma 5.1(i), we know that $Q_i^{k*} \cap Q_j^{(k+1)*} = \emptyset$ for $j \in F_1^{k+1}$. From this, (5.2) and (5.10), we conclude that

$$\begin{aligned} h_i^k &= \left(f - P_i^k\right)\zeta_i^k - \sum_{j \in F_1^{k+1}} f\zeta_j^{k+1}\zeta_i^k - \sum_{j\in F_2^{k+1}} \left(f - P_j^{k+1}\right)\zeta_j^{k+1}\zeta_i^k \\ &\quad + \sum_{j\in F_2^{k+1}} P_{i,j}^{k+1}\zeta_j^{k+1} \\ &= \left(f - P_i^k\right)\zeta_i^k - \sum_{j\in F_2^{k+1}} \left\{\left(f - P_j^{k+1}\right)\zeta_j^{k+1}\zeta_i^k - P_{i,j}^{k+1}\zeta_j^{k+1}\right\}. \end{aligned} \tag{5.16}$$

Let $\widetilde{Q}_i^k \equiv 2^6 n Q_i^{k*}$. Then $\operatorname{supp}(h_i^k) \subset \widetilde{Q}_i^k$. By $l_i^k < 1/(2^4 n)$, Lemma 2.3(v) and Remark 2.4 with $\widetilde{C} \equiv 4a$, we know that there exists a positive constant C_{12} such that

$$\omega(\widetilde{Q}_i^k) \le C_{12}\omega(Q_i^{k*}). \tag{5.17}$$

Moveover, h_i^k satisfies the desired moment conditions, which are deduced from the moment conditions of $(f - P_i^k)\zeta_i^k$ (see (5.3)) and $(f - P_j^{k+1})\zeta_j^{k+1}\zeta_i^k - P_{i,j}^{k+1}\zeta_j^{k+1}$ (see (5.4)). Let $\widetilde{A}_2 \equiv \max\{C_{10}, C_{12}\}$,

$$\lambda_i^k \equiv \widetilde{A}_2 2^k \omega(\widetilde{Q}_i^k)\rho(\omega(\widetilde{Q}_i^k)) \tag{5.18}$$

and $a_i^k \equiv (\lambda_i^k)^{-1} h_i^k$. By this, (5.13), $\operatorname{supp}(h_i^k) \subset \widetilde{Q}_i^k$ and the moment conditions of h_i^k, we know that a_i^k is a $(\rho,\, \infty,\, s)_\omega$-atom.

Thus, from (5.9), (5.10), Case 1) and Case 2), we infer that

$$f = \sum_{k\in\mathbb{Z}} \sum_{i\in\mathbb{N}} \lambda_i^k a_i^k$$

holds in both $\mathcal{D}'(\mathbb{R}^n)$ and almost everywhere, where for every k and i, $\lambda_i^k \in \mathbb{C}$ and a_i^k is a $(\rho,\, \infty,\, s)_\omega$-atom, which shows (5.7) in the case that $k_0 = -\infty$ by letting $\lambda_0 = 0$. Furthermore, by the fact that $\Phi(t) \sim \int_0^t \frac{\Phi(s)}{s}\,ds$ for all $t \in (0,\infty)$, (5.15), (5.18), (5.14), (5.17), the upper type 1 property of Φ, Fubini's theorem and the bounded interior property of $\{Q_i^{k*}\}$, we know that for any $\lambda \in (0,\infty)$,

$$\begin{aligned}
&\sum_{k,\,i} \omega(\widetilde{Q}_i^k)\Phi\left(\frac{|\lambda_i^k|}{\lambda\rho(\omega(\widetilde{Q}_i^k))\omega(\widetilde{Q}_i^k)}\right)\\
&\quad\lesssim \sum_{k,\,i} \omega(\widetilde{Q}_i^k)\Phi\left(\frac{2^k}{\lambda}\right) \lesssim \sum_{k,i} \omega(Q_i^{k*})\Phi\left(\frac{2^k}{\lambda}\right)\\
&\quad\lesssim \sum_{k} \omega(\Omega_{2^k})\Phi\left(\frac{2^k}{\lambda}\right) \sim \sum_{k} \int_{\Omega_{2^k}} \Phi\left(\frac{2^k}{\lambda}\right)\omega(x)\,dx\\
&\quad\lesssim \int_{\mathbb{R}^n} \sum_{k<\log[\mathcal{G}_N(f)(x)]} \Phi\left(\frac{2^k}{\lambda}\right)\omega(x)\,dx \lesssim \int_{\mathbb{R}^n} \sum_{k<\log[\mathcal{G}_N(f)(x)]} \int_{2^k}^{2^{k+1}} \Phi\left(\frac{t}{\lambda}\right)\frac{dt}{t}\omega(x)\,dx\\
&\quad\lesssim \int_{\mathbb{R}^n} \int_0^{2\mathcal{G}_N(f)(x)/\lambda} \Phi(t)\frac{dt}{t}\omega(x)\,dx \lesssim \int_{\mathbb{R}^n} \Phi\left(\frac{\mathcal{G}_N(f)(x)}{\lambda}\right)\omega(x)\,dx,
\end{aligned}$$

which implies (5.8) in the case that $k_0 = -\infty$.

Finally, we consider the case that $k_0 > -\infty$. In this case, by $f \in h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$, we see that $\omega(\mathbb{R}^n) < \infty$. Adapting the previous arguments, we conclude that

$$f = \sum_{k=k_0}^{\infty} \left(g^{k+1} - g^k\right) + g^{k_0} \equiv \widetilde{f} + g^{k_0}, \tag{5.19}$$

and for the function $\widetilde{f}$, we have the same $(\rho,\,\infty,\,s)_\omega$-atomic decomposition as above that

$$\widetilde{f} = \sum_{k\geq k_0,\,i} \lambda_i^k a_i^k \tag{5.20}$$

and

$$\Lambda\left(\{\lambda_i^k a_i^k\}_{k\geq k_0,\,i}\right) \lesssim \|f\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}. \tag{5.21}$$

From Lemma 4.7(ii), it follows that

$$\|g^{k_0}\|_{L^\infty_\omega(\mathbb{R}^n)} \leq C_9 2^{k_0} \leq 2C_9 \inf_{x\in\mathbb{R}^n} \mathcal{G}_N(f)(x), \tag{5.22}$$

where C_9 is the same as in Lemma 4.7(ii). Let $\lambda_0 \equiv 2C_9 2^{k_0}\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))$ and

$$a_0 \equiv \lambda_0^{-1} g^{k_0}.$$

Then we have

$$\|a_0\|_{L^\infty_\omega(\mathbb{R}^n)} \leq [\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))]^{-1}.$$

Thus, we know that a_0 is a $(\rho,\,\infty)_\omega$-single-atom and $g^{k_0} = \lambda_0 a_0$, which together with (5.19) and (5.20) implies (5.7) in the case that $k_0 > -\infty$. Moreover, from (5.22), we deduce that for any $\lambda \in (0,\infty)$,

$$\omega(\mathbb{R}^n)\Phi\left(\frac{|\lambda_0|}{\lambda\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))}\right) = \omega(\mathbb{R}^n)\Phi\left(\frac{C_9 2^{k_0}}{\lambda}\right) \lesssim \int_{\mathbb{R}^n} \Phi\left(\frac{\mathcal{G}_N(f)(x)}{\lambda}\right)\omega(x)\,dx,$$

which together with (5.21) implies (5.8) in the case that $k_0 > -\infty$. This finishes the proof of Lemma 5.4. ■

REMARK 5.5. By its proof, all $(\rho,\,\infty,\,s)_\omega$-atoms in Lemma 5.4 can be taken to have supports Q satisfying $l(Q)\in(0,2]$. Indeed, for any $(\rho,\,\infty,\,s)_\omega$-atom a supported in a cube Q_0 with $l(Q_0)>2$, we know that there exist $N_0\in\mathbb{N}$, depending on $l(Q_0)$ and n, and cubes $\{Q_i\}_{i=1}^{N_0}$ satisfying $l(Q_i)\in[1,2]$ with $i\in\{1,\,\cdots,\,N_0\}$ such that $\cup_{i=1}^{N_0}Q_i=Q_0$, for any $x\in Q_0$, $1\le\sum_{i=1}^{N_0}\chi_{Q_i}(x)\le C(n)$, and

$$a=\frac{1}{\sum_{j=1}^{N_0}\chi_{Q_j}}\sum_{i=1}^{N_0}a\chi_{Q_i},$$

where $C(n)$ is a positive integer, only depending on n. For any given $\lambda_0\in\mathbb{C}$ and $i\in\{1,\,\cdots,\,N_0\}$, let

$$\gamma_i\equiv\frac{\lambda_0\omega(Q_i)\rho(\omega(Q_i))}{\omega(Q_0)\rho(\omega(Q_0))}$$

and

$$b_i\equiv\frac{\omega(Q_0)\rho(\omega(Q_0))a\chi_{Q_i}}{\omega(Q_i)\rho(\omega(Q_i))\sum_{i=1}^{N_0}\chi_{Q_i}}.$$

Then for any $i\in\{1,\,2,\,\cdots,\,N_0\}$, b_i is a $(\rho,\,\infty,\,s)_\omega$-atom supported in the cube Q_i and

$$\lambda_0a=\sum_{i=1}^{N_0}\gamma_ib_i. \tag{5.23}$$

From the definitions of γ_i and b_i, $\cup_{i=1}^{N_0}Q_i=Q_0$, and for any $x\in Q_0$,

$$1\le\sum_{i=1}^{N_0}\chi_{Q_i}(x)\le C(n),$$

we also conclude that for all $\lambda\in(0,\infty)$,

$$\sum_{i=1}^{N_0}\omega(Q_i)\Phi\left(\frac{|\gamma_i|}{\lambda\omega(Q_i)\rho(\omega(Q_i))}\right)\le C(n)\omega(Q_0)\Phi\left(\frac{|\lambda_0|}{\lambda\omega(Q_0)\rho(\omega(Q_0))}\right). \tag{5.24}$$

Thus, by the proof of Lemma 5.4, (5.23) and (5.24), we see that the claim holds.

Now we state the weighted atomic decompositions of $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$ as follows.

THEOREM 5.6. *Let Φ satisfy Assumption* (A), *$\omega\in A^{\rm loc}_\infty(\mathbb{R}^n)$, and q_ω and $N_{\Phi,\,\omega}$ be respectively as in* (2.4) *and* (3.25). *If $q\in(q_\omega,\infty]$, integers s and N satisfy $N\ge N_{\Phi,\,\omega}$ and $N>s\ge\lfloor n(\frac{q_\omega}{p_\Phi}-1)\rfloor$, then*

$$h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)=h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)=h^{\Phi}_{\omega,\,N_{\Phi,\,\omega}}(\mathbb{R}^n)$$

with equivalent norms.

Proof. It is easy to see that

$$h^{\rho,\,\infty,\,s_1}_\omega(\mathbb{R}^n)\subset h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)\subset h^{\Phi}_{\omega,\,N_{\Phi,\,\omega}}(\mathbb{R}^n)\subset h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)\subset h^{\Phi}_{\omega,\,N_1}(\mathbb{R}^n),$$

where the integers s_1 and N_1 are respectively no less than s and N, and the inclusions are continuous. Thus, to prove Theorem 5.6, it suffices to prove that for any integers N, s satisfying $N>s\ge\lfloor n(\frac{q_\omega}{p_\Phi}-1)\rfloor$, $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)\subset h^{\rho,\,\infty,\,s}_\omega(\mathbb{R}^n)$ and for all $f\in h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$,

$$\|f\|_{h^{\rho,\,\infty,\,s}_\omega(\mathbb{R}^n)}\lesssim\|f\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}.$$

Let $f \in h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$. By Corollary 4.8, there exists a sequence of functions,

$$\{f_m\}_{m\in\mathbb{N}} \subset (h^{\Phi}_{\omega,\,N}(\mathbb{R}^n) \cap L^q_{\omega}(\mathbb{R}^n)),$$

such that for all $m \in \mathbb{N}$,

$$\|f_m\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)} \le 2^{-m}\|f\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)} \tag{5.25}$$

and $f = \sum_{m\in\mathbb{N}} f_m$ in $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$. By Lemma 5.4, we know that for each $m \in \mathbb{N}$, f_m has an atomic decomposition

$$f = \sum_{i\in\mathbb{Z}_+} \lambda_i^m a_i^m$$

in $\mathcal{D}'(\mathbb{R}^n)$ with

$$\Lambda(\{\lambda_i^m a_i^m\}_i) \lesssim \|f_m\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)},$$

where $\{\lambda_i^m\}_{i\in\mathbb{Z}_+} \subset \mathbb{C}$, $\{a_i^m\}_{i\in\mathbb{N}}$ are $(\rho,\,\infty,\,s)_\omega$-atoms and a_0^m is a $(\rho,\,\infty)_\omega$-single-atom.

Let

$$\widetilde{\lambda}_0 \equiv \omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n)) \sum_{m=1}^{\infty} |\lambda_0^m| \|a_0^m\|_{L^\infty_\omega(\mathbb{R}^n)}$$

and

$$\widetilde{a}_0 \equiv \left(\widetilde{\lambda}_0\right)^{-1} \sum_{m=1}^{\infty} \lambda_0^m a_0^m.$$

Then

$$\widetilde{\lambda}_0 \widetilde{a}_0 = \sum_{m=1}^{\infty} \lambda_0^m a_0^m.$$

It is easy to see that

$$\|\widetilde{a}_0\|_{L^\infty_\omega(\mathbb{R}^n)} \le [\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))]^{-1},$$

which implies that $\widetilde{a}_0$ is a $(\rho,\infty)_\omega$-single-atom. Since Φ is increasing, by (5.8), we know that for any $m \in \mathbb{N}$,

$$\omega(\mathbb{R}^n)\Phi\left(\frac{|\lambda_0^m|}{C\|f_m\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))}\right) \le 1, \tag{5.26}$$

where C is as in (5.8). Let

$$\widetilde{\gamma} \equiv C\left(\sum_{i=m}^{\infty} \|f_m\|^{p_\Phi}_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}\right)^{1/p_\Phi},$$

where C is as in (5.8). Then, from the continuity and subadditivity of Φ, the strictly lower type p_Φ property of Φ and (5.26), it follows that

$$\begin{aligned}
&\omega(\mathbb{R}^n)\Phi\left(\frac{|\widetilde{\lambda}_0|}{\widetilde{\gamma}\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))}\right)\\
&\quad = \omega(\mathbb{R}^n)\Phi\left(\frac{\sum_{m=1}^{\infty} |\lambda_0^m| \|a_0^m\|_{L^\infty_\omega(\mathbb{R}^n)}}{\widetilde{\gamma}}\right)\\
&\quad \le \omega(\mathbb{R}^n)\sum_{m=1}^{\infty} \frac{\|f_m\|^{p_\Phi}_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}}{\widetilde{\gamma}^{p_\Phi}}\Phi\left(\frac{|\lambda_0^m|}{C\|f_m\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))}\right) \le 1,
\end{aligned}$$

which together with (5.25) implies that

$$\Lambda(\{\widetilde{\lambda}_0\widetilde{a}_0\}) \le \widetilde{\gamma} \lesssim \|f\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}.$$

Thus, we see that

$$f = \sum_{m\in\mathbb{N}}\sum_{i\in\mathbb{N}} \lambda_i^m a_i^m + \widetilde{\lambda}_0\widetilde{a}_0 \in h^{\rho,\,\infty,\,s}_{\omega}(\mathbb{R}^n)$$

and

$$\|f\|_{h^{\rho,\,\infty,\,s}_{\omega}(\mathbb{R}^n)} \lesssim \|f\|_{h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)}.$$

This finishes the proof of Theorem 5.6. ∎

Remark 5.7. Let $p \in (0,1]$. Theorem 5.1 when $\Phi(t) \equiv t^p$ for all $t \in (0,\infty)$ was obtained by Tang [49, Theorem 5.1].

For simplicity, from now on, we denote by $h^{\Phi}_{\omega}(\mathbb{R}^n)$ the *weighted local Orlicz-Hardy space* $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$ when $N \ge N_{\Phi,\,\omega}$.

6. Finite atomic decompositions

In this section, we prove that for any given finite linear combination of weighted atoms when $q < \infty$ (or continuous $(\rho,\, q,\, s)_\omega$-atoms when $q = \infty$), its norm in $h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)$ can be achieved via all its finite weighted atomic decompositions. This extends the main results in [35, 57] to the setting of weighted local Orlicz-Hardy spaces. As applications, we see that for a given admissible triplet $(\rho,\, q,\, s)_\omega$ and a β-quasi-Banach space $\mathcal{B}_\beta$ with $\beta \in (0,1]$, if T is a $\mathcal{B}_\beta$-sublinear operator, and maps all $(\rho,\, q,\, s)_\omega$-atoms and $(\rho,\, q)_\omega$-single-atoms with $q < \infty$ (or all continuous $(\rho,\, q,\, s)_\omega$-atoms with $q = \infty$) into uniformly bounded elements of $\mathcal{B}_\beta$, then T uniquely extends to a bounded $\mathcal{B}_\beta$-sublinear operator from $h^{\Phi}_{\omega}(\mathbb{R}^n)$ to $\mathcal{B}_\beta$.

Definition 6.1. Let Φ satisfy Assumption (A), $\omega \in A^{\rm loc}_{\infty}(\mathbb{R}^n)$ and $(\rho,\, q,\, s)_\omega$ be admissible as in Definition 3.4. The *space* $h^{\rho,\,q,\,s}_{\omega,\,\rm fin}(\mathbb{R}^n)$ is defined to be the vector space of all finite linear combinations of $(\rho,\, q,\, s)_\omega$-atoms and a $(\rho,\, q)_\omega$-single-atom, and the *norm* of f in $h^{\rho,\,q,\,s}_{\omega,\,\rm fin}(\mathbb{R}^n)$ is defined by

$$\|f\|_{h^{\rho,\,q,\,s}_{\omega,\,\rm fin}(\mathbb{R}^n)} \equiv \inf\left\{\Lambda(\{\lambda_i a_i\}_i) : f = \sum_{i=0}^{k} \lambda_i a_i,\ k \in \mathbb{Z}_+,\ \{\lambda_i\}_{i=0}^k \subset \mathbb{C},\ \{a_i\}_{i=1}^k \text{ are } (\rho,\, q,\, s)_\omega\text{-atoms and } a_0 \text{ is a } (\rho,\, q)_\omega\text{-single-atom}\right\}.$$

Obviously, for any admissible triplet $(\rho,\, q,\, s)_\omega$, $h^{\rho,\,q,\,s}_{\omega,\,\rm fin}(\mathbb{R}^n)$ is dense in $h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$ with respect to the quasi-norm $\|\cdot\|_{h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)}$.

Theorem 6.2. *Let* Φ *satisfy Assumption* (A)*,* $\omega \in A^{\rm loc}_{\infty}(\mathbb{R}^n)$*,* q_ω *be as in* (2.4) *and* $(\rho,\, q,\, s)_\omega$ *be admissible as in Definition 3.4.*

(i) *If* $q \in (q_\omega,\infty)$*, then* $\|\cdot\|_{h^{\rho,\,q,\,s}_{\omega,\,\rm fin}(\mathbb{R}^n)}$ *and* $\|\cdot\|_{h^{\Phi}_{\omega}(\mathbb{R}^n)}$ *are equivalent quasi-norms on* $h^{\rho,\,q,\,s}_{\omega,\,\rm fin}(\mathbb{R}^n)$.

(ii) *Let* $h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin},\,c}(\mathbb{R}^n)$ *denote the set of all* $f\in h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$ *with compact support. Then* $\|\cdot\|_{h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)}$ *and* $\|\cdot\|_{h^{\Phi}_{\omega}(\mathbb{R}^n)}$ *are equivalent quasi-norms on* $h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin},\,c}(\mathbb{R}^n)\cap C(\mathbb{R}^n)$.

Proof. We first show (i). Let $q\in(q_\omega,\infty)$ and $(\rho,\,q,\,s)_\omega$ be admissible. Obviously, from Theorem 5.6, we infer that $h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)\subset h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)=h^{\Phi}_{\omega}(\mathbb{R}^n)$ and for all $f\in h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$,

$$\|f\|_{h^{\Phi}_{\omega}(\mathbb{R}^n)}\lesssim\|f\|_{h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)}.$$

Thus, we only need show that for all $f\in h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$,

$$\|f\|_{h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)}\lesssim\|f\|_{h^{\Phi}_{\omega}(\mathbb{R}^n)}. \tag{6.1}$$

By homogeneity, without loss of generality, we may assume that $f\in h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$ with $\|f\|_{h^{\Phi}_{\omega}(\mathbb{R}^n)}=1$. In the rest of this section, for any $f\in h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$, let k_0 be as in Section 5 and Ω_{2^k} with $k\ge k_0$ as in (4.1) with $\lambda=2^k$. Since $f\in(h^{\Phi}_{\omega,\,N}(\mathbb{R}^n)\cap L^q_\omega(\mathbb{R}^n))$, by Lemma 5.4, there exist $\lambda_0\in\mathbb{C}$, $\{\lambda^k_i\}_{k\ge k_0,\,i}\subset\mathbb{C}$, a $(\rho,\,\infty)_\omega$-single-atom a_0 and $(\rho,\,\infty,\,s)_\omega$-atoms $\{a^k_i\}_{k\ge k_0,\,i}$, such that

$$f=\sum_{k\ge k_0}\sum_i\lambda^k_i a^k_i+\lambda_0a_0 \tag{6.2}$$

holds both in $\mathcal{D}'(\mathbb{R}^n)$ and almost everywhere. First, we claim that (6.2) also holds in $L^q_\omega(\mathbb{R}^n)$. For any $x\in\mathbb{R}^n$, by $\mathbb{R}^n=\cup_{k\ge k_0}(\Omega_{2^k}\setminus\Omega_{2^{k+1}})$, we see that there exists $j\in\mathbb{Z}$ such that $x\in(\Omega_{2^j}\setminus\Omega_{2^{j+1}})$. By the proof of Lemma 5.4, we know that for all $k>j$, $\operatorname{supp}(a^k_i)\subset\widetilde{Q}^k_i\subset\Omega_{2^k}\subset\Omega_{2^{j+1}}$; then from (5.13) and (5.22), we conclude that

$$\left|\sum_{k\ge k_0}\sum_i\lambda^k_i a^k_i(x)\right|+|\lambda_0a_0(x)|\lesssim\sum_{k_0\le k\le j}2^k+2^{k_0}\lesssim 2^j\lesssim\mathcal{G}_N(f)(x).$$

Since $f\in L^q_\omega(\mathbb{R}^n)$, from Proposition 3.2(ii), we infer that $\mathcal{G}_N(f)(x)\in L^q_\omega(\mathbb{R}^n)$. This combined with the Lebesgue dominated convergence theorem implies that

$$\sum_{k\ge k_0}\sum_i\lambda^k_i a^k_i+\lambda_0a_0$$

converges to f in $L^q_\omega(\mathbb{R}^n)$, which completes the proof of the claim.

Next, we show (6.1) by considering the following two cases for ω.

Case 1) $\omega(\mathbb{R}^n)=\infty$. In this case, by $f\in L^q_\omega(\mathbb{R}^n)$, we know that $k_0=-\infty$ and $a_0(x)=0$ for almost every $x\in\mathbb{R}^n$ in (6.2). Thus, in this case, (6.2) has the version

$$f=\sum_{k\in\mathbb{Z}}\sum_i\lambda^k_i a^k_i.$$

Since, when $\omega(\mathbb{R}^n)=\infty$, all $(\rho,\,q)_\omega$-single-atoms are 0, if $f\in h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$, then f has compact support. Assume that $\operatorname{supp}(f)\subset Q_0\equiv Q(x_0,r_0)$ and

$$\widetilde{Q}_0\equiv Q(x_0,\sqrt{n}r_0+2^{3(10+n)+1}).$$

Then for any $\psi\in\mathcal{D}_N(\mathbb{R}^n)$, $x\in\mathbb{R}^n\setminus\widetilde{Q}_0$ and $t\in(0,1)$, we have

$$\psi_t*f(x)=\int_{Q(x_0,r_0)}\psi_t(x-y)f(y)\,dy=\int_{B(x,2^{3(10+n)})\cap Q(x_0,r_0)}\psi_t(x-y)f(y)\,dy=0.$$

Thus, for any $k \in \mathbb{Z}$, $\Omega_{2^k} \subset \widetilde{Q}_0$, which implies that $\operatorname{supp}(\sum_{k\in\mathbb{Z}} \sum_i \lambda_i^k a_i^k) \subset \widetilde{Q}_0$. For each positive integer K, let

$$F_K \equiv \{(i,k) :\, k \in \mathbb{Z},\, k \geq k_0, i \in \mathbb{N},\, |k| + i \leq K\}$$

and

$$f_K \equiv \sum_{(k,i)\in F_K} \lambda_i^k a_i^k.$$

Then, by the above claim, we know that f_K converges to f in $L^q_\omega(\mathbb{R}^n)$. Thus, for any given $\epsilon \in (0,1)$, there exists $K_0 \in \mathbb{N}$ large enough such that

$$\|(f - f_{K_0})/\epsilon\|_{L^q_\omega(\mathbb{R}^n)} \leq [\rho(\omega(\widetilde{Q}_0))]^{-1}[\omega(\widetilde{Q}_0)]^{1/q-1},$$

which together with $\operatorname{supp}(f - f_{K_0})/\epsilon \subset \widetilde{Q}_0$ implies that $(f - f_{K_0})/\epsilon$ is a $(\rho,\, q,\, s)_\omega$-atom. Moreover, we equivalently divide $\widetilde{Q}_0$ into the union of some cubes $\{Q_i\}_{i=1}^{N_0}$ with disjoint interior and their sidelengths satisfying $l_i \in (1,2]$, where N_0 depends only on r_0 and n. It is clear that

$$\|(f - f_{K_0})\chi_{Q_i}/\epsilon\|_{L^q_\omega(\mathbb{R}^n)} \leq [\rho(\omega(\widetilde{Q}_0))]^{-1}[\omega(\widetilde{Q}_0)]^{1/q-1} \leq [\rho(\omega(Q_i))]^{-1}[\omega(Q_i)]^{1/q-1},$$

which together with $\operatorname{supp}((f - f_{K_0})\chi_{Q_i}/\epsilon) \subset Q_i$ implies that $(f - f_{K_0})\chi_{Q_i}/\epsilon$ is a $(\rho,\, q,\, s)_\omega$-atom for $i = 1,\, 2, \cdots,\, N_0$. Thus,

$$f = f_{K_0} + \sum_{i=1}^{N_0} (f - f_{K_0})\chi_{Q_i}$$

almost everywhere is a finite linear weighted atom combination of f. Let

$$b_i \equiv (f - f_{K_0})\chi_{Q_i}/\epsilon$$

and take $\epsilon \equiv N_0^{-1/p_\Phi}$. Then, by (2.8) with $t \equiv \omega(Q_i)$, Remark 3.6(ii) and the lower type p_Φ property of Φ,

$$\begin{aligned}\|f\|_{h^{\rho,\, q,\, s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)} &\lesssim \Lambda\left(\{\lambda_i^k a_i^k\}_{(i,k)\in F_{K_0}}\right) + \Lambda\left(\{\epsilon b_i\}_{i=1}^{N_0}\right)\\ &\lesssim \|f\|_{h^{\rho,\, q,\, s}_\omega(\mathbb{R}^n)} + \inf\left\{\lambda > 0 : \sum_{i=1}^{N_0} \omega(Q_i)\Phi\left(\frac{\epsilon}{\lambda\omega(Q_i)\rho(\omega(Q_i))}\right) \leq 1\right\} \lesssim 1,\end{aligned}$$

which implies (6.1) in Case 1).

Case 2) $\omega(\mathbb{R}^n) < \infty$. In this case, f may not have compact support. Similarly to Case 1), for any positive integer K, let

$$f_K \equiv \sum_{(k,i)\in F_K} \lambda_i^k a_i^k + \lambda_0 a_0$$

and $b_K \equiv f - f_K$, where F_K is as in Case 1). From the above claim, we deduce that f_K converges to f in $L^q_\omega(\mathbb{R}^n)$. Thus, there exists a positive integer $K_1 \in \mathbb{N}$ large enough such that

$$\|b_{K_1}\|_{L^q_\omega(\mathbb{R}^n)} \leq [\rho(\omega(\mathbb{R}^n))]^{-1}[\omega(\mathbb{R}^n)]^{1/q-1}.$$

Thus, b_{K_1} is a $(\rho,\, q)_\omega$-single-atom and $f = f_{K_1} + b_{K_1}$ is a finite linear weighted atom combination of f. Moreover, by Remark 3.6(ii) and (2.8) with $t \equiv \omega(\mathbb{R}^n)$,

$$\begin{aligned}\|f\|_{h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)} &\lesssim \Lambda\left(\{\lambda_i^k a_i^k\}_{(i,k)\in F_{K_1}}\right) + \Lambda\left(\{b_{K_1}\}\right)\\ &\lesssim \|f\|_{h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)} + \inf\left\{\lambda>0:\ \omega(\mathbb{R}^n)\Phi\left(\frac{1}{\lambda\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))}\right)\le 1\right\} \lesssim 1,\end{aligned}$$

which implies (6.1) in Case 2). This finishes the proof of (i).

We now prove (ii). In this case, similarly to the proof of (i), we only need prove that for all $f \in h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin},\,c}(\mathbb{R}^n)$,

$$\|f\|_{h^{\rho,\,\infty,\,s}_{\omega}(\mathbb{R}^n)} \lesssim \|f\|_{h^{\Phi}_{\omega}(\mathbb{R}^n)}.$$

Again, by homogeneity, without loss of generality, we may assume that $\|f\|_{h^{\Phi}_{\omega}(\mathbb{R}^n)} = 1$. Since f has compact support, by the definition of $\mathcal{G}_N(f)$, it is easy to know that $\mathcal{G}_N(f)$ also has compact support. Assume that $\operatorname{supp}(\mathcal{G}_N(f)) \subset B(0, R_0)$ for some $R_0 \in (0,\infty)$. By $f \in L^\infty_\omega(\mathbb{R}^n)$, we have that $\mathcal{G}_N f \in L^\infty_\omega(\mathbb{R}^n)$. Thus, there exists $k_1 \in \mathbb{Z}$ such that $\Omega_{2^k} = \emptyset$ for any $k \in \mathbb{Z}$ with $k \ge k_1 + 1$. By Lemma 5.4, there exist $\lambda_0 \in \mathbb{C}$, $\{\lambda_i^k\}_{k_1\ge k\ge k_0,\,i} \subset \mathbb{C}$, a $(\rho,\,\infty)_\omega$-single-atom a_0 and $(\rho,\,\infty,\,s)_\omega$-atoms $\{a_i^k\}_{k_1\ge k\ge k_0,\,i}$ such that

$$f = \sum_{k=k_0}^{k_1}\sum_i \lambda_i^k a_i^k + \lambda_0 a_0$$

holds both in $\mathcal{D}'(\mathbb{R}^n)$ and almost everywhere. By the fact that f is uniformly continuous, we know that for any given $\varepsilon \in (0,\infty)$, there exists a $\delta \in (0,\infty)$ such that if

$$|x-y| < \sqrt{n}\delta/2,$$

then $|f(x) - f(y)| < \varepsilon$. Without loss of generality, we may assume that $\delta < 1$. Write $f = f_1^\varepsilon + f_2^\varepsilon$ with

$$f_1^\varepsilon \equiv \sum_{(i,k)\in G_1} \lambda_i^k a_i^k + \lambda_0 a_0$$

and

$$f_2^\varepsilon \equiv \sum_{(i,k)\in G_2} \lambda_i^k a_i^k,$$

where

$$G_1 \equiv \left\{(i,k):\ l(\widetilde{Q}_i^k) \ge \delta,\ k_0 \le k \le k_1\right\},$$

$$G_2 \equiv \left\{(i,k):\ l(\widetilde{Q}_i^k) < \delta,\ k_0 \le k \le k_1\right\},$$

and $\widetilde{Q}_i^k$ is the support of a_i^k (see the proof of Lemma 5.4). For any fixed integer $k \in [k_0, k_1]$, by Lemma 5.1(ii) and $\Omega_{2^k} \subset B(0, R_0)$, we see that G_1 is a finite set.

For any $(i,k) \in G_2$ and $x \in \widetilde{Q}_i^k$, $|f(x) - f(x_i^k)| < \varepsilon$. For all $x \in \mathbb{R}^n$, let

$$\widetilde{f}(x) \equiv [f(x) - f(x_i^k)]\chi_{\widetilde{Q}_i^k}(x)$$

and $\widetilde{P}_i^k(x) \equiv P_i^k(x) - f(x_i^k)$. By the definition of P_i^k, for all $P \in \mathcal{P}_s(\mathbb{R}^n)$,

$$\int_{\mathbb{R}^n}\left[\widetilde{f}(x) - \widetilde{P}_i^k(x)\right] P(x)\zeta_i^k(x)\,dx = 0.$$

Since $|\widetilde{f}(x)|<\varepsilon$ for all $x\in\mathbb{R}^n$ implies that $\mathcal{G}_N(\widetilde{f})(x)\lesssim\varepsilon$ for all $x\in\mathbb{R}^n$, then by Lemma 4.1, we see that

$$\sup_{y\in\mathbb{R}^n}\left|\widetilde{P}_i^k(y)\zeta_i^k(y)\right|\lesssim\sup_{y\in\mathbb{R}^n}\left|\mathcal{G}_N(\widetilde{f})(y)\right|\lesssim\varepsilon. \tag{6.3}$$

Let $\widetilde{P}_{i,\,j}^k\in\mathcal{P}_s(\mathbb{R}^n)$ be such that for any $P\in\mathcal{P}_s(\mathbb{R}^n)$,

$$\int_{\mathbb{R}^n}\left[\widetilde{f}(x)-\widetilde{P}_i^k(x)\right]\zeta_i^k(x)P(x)\zeta_i^{k+1}(x)\,dx=\int_{\mathbb{R}^n}\widetilde{P}_{i,\,j}^{k+1}(x)P(x)\zeta_j^{k+1}(x)\,dx.$$

Since $(\widetilde{f}-\widetilde{P}_i^k)\zeta_i^k=(f-P_i^k)\zeta_i^k$, by $\operatorname{supp}(\zeta_i^k)\subset\widetilde{Q}_i^k$, we have $\widetilde{P}_{i,\,j}^k=P_{i,\,j}^k$. Then from Lemma 5.2, we deduce that

$$\sup_{y\in\mathbb{R}^n}\left|\widetilde{P}_{i,\,j}^k(y)\zeta_i^{k+1}(y)\right|\lesssim\sup_{y\in\mathbb{R}^n}\left|\mathcal{G}_N(\widetilde{f})(y)\right|\lesssim\varepsilon. \tag{6.4}$$

Thus, by the definition of $\lambda_i^k a_i^k$, $\sum_j\zeta_j^{k+1}=\chi_{\Omega_{2^{k+1}}}$ and (5.11), we know that

$$\begin{aligned}\lambda_i^k a_i^k&=f\chi_{\Omega^\complement_{2^{k+1}}}\zeta_i^k-P_i^k\zeta_i^k+\sum_{j\in F_2^{k+1}}P_j^{k+1}\zeta_i^k\zeta_j^{k+1}+\sum_{j\in F_2^{k+1}}P_{i,\,j}^{k+1}\zeta_j^{k+1}\\&=\widetilde{f}\chi_{\Omega^\complement_{2^{k+1}}}\zeta_i^k-\widetilde{P}_i^k\zeta_i^k+\sum_{j\in F_2^{k+1}}\widetilde{P}_j^{k+1}\zeta_i^k\zeta_j^{k+1}+\sum_{j\in F_2^{k+1}}\widetilde{P}_{i,\,j}^{k+1}\zeta_j^{k+1}.\end{aligned}$$

From this together with (6.3), (6.4) and Lemma 5.1(ii), it follows that $|\lambda_i^k a_i^k|\lesssim\varepsilon$ for all $x\in\widetilde{Q}_i^k$ with $(i,k)\in G_2$. Moreover, using Lemma 5.1(ii) again, we conclude that

$$|f_2^\varepsilon|\lesssim\sum_{k=k_0}^{k_1}\varepsilon\lesssim(k_1-k_0)\varepsilon.$$

By the arbitrariness of ε, $\operatorname{supp}(f_2^\varepsilon)\subset B(0,R_0)$ and $|f_2^\varepsilon|\lesssim(k_1-k_0)\varepsilon$, we choose ε small enough such that f_2^ε is an arbitrarily small multiple of a $(\rho,\,\infty,\,s)_\omega$-atom. In particular, we choose $\varepsilon_0\in(0,\infty)$ such that $f_2^{\varepsilon_0}=\widetilde{\lambda}\widetilde{a}$ with $|\widetilde{\lambda}|\le1$ and $\widetilde{a}$ is a $(\rho,\,\infty,\,s)_\omega$-atom, then

$$f=\sum_{(i,k)\in G_1}\lambda_i^k a_i^k+\lambda_0a_0+\widetilde{\lambda}\widetilde{a}$$

is a finite weighted atomic decomposition of f, and

$$\|f\|_{h_\omega^{\rho,\,\infty,\,s}(\mathbb{R}^n)}\lesssim\|f\|_{h_\omega^\Phi(\mathbb{R}^n)}+1\lesssim1,$$

which completes the proof of Theorem 6.2. ■

REMARK 6.3. (i) From the proof of Theorem 6.2, it follows that for any $f\in h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$ with $q\in(q_\omega,\infty)$, there exist $\{\lambda_j\}_{j=0}^k\subset\mathbb{C}$, a $(\rho,\,q)_\omega$-single-atom a_0 and $(\rho,\,q,\,s)_\omega$-atoms $\{a_j\}_{j=1}^k$ satisfying $\operatorname{supp}(a_j)\subset Q_j$ with $l(Q_j)\in(0,2]$ such that $f=\sum_{j=0}^k\lambda_ja_j$ in both $L^q_\omega(\mathbb{R}^n)$ and $\mathcal{D}'(\mathbb{R}^n)$. Moreover, for all $f\in h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$,

$$\begin{aligned}&\|f\|_{h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)}\sim\|f\|_{h_\omega^\Phi(\mathbb{R}^n)}\\&\qquad\sim\inf\left\{\Lambda\{\lambda_ia_i\}_i:\ f=\sum_{i=0}^k\lambda_ia_i,\ k\in\mathbb{Z}_+,\ \{a_i\}_{i=1}^k\ \text{are}\,(\rho,\,q,\,s)_\omega\text{-atoms}\right.\\&\qquad\qquad\left.\text{satisfying }\operatorname{supp}(a_j)\subset Q_j,\ l(Q_j)\in(0,2]\right.\end{aligned}$$

$$\text{and } a_0 \text{ is a } (\rho,\, q)_\omega\text{-single-atom}\Bigg\}.$$

(ii) Obviously, when $\omega(\mathbb{R}^n) = \infty$,

$$h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin},\,c}(\mathbb{R}^n) \cap C(\mathbb{R}^n) = h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n) \cap C(\mathbb{R}^n).$$

As an application of Theorem 6.2, we establish the boundedness on $h^\Phi_\omega(\mathbb{R}^n)$ of quasi-Banach-valued sublinear operators.

Recall that a *quasi-Banach space* $\mathcal{B}$ is a vector space endowed with a quasi-norm $\|\cdot\|_{\mathcal{B}}$ which is nonnegative, non-degenerate (i. e., $\|f\|_{\mathcal{B}} = 0$ if and only if $f = 0$), homogeneous, and obeys the quasi-triangle inequality, i. e., there exists a positive constant K no less than 1 such that for all $f,\, g \in \mathcal{B}$,

$$\|f + g\|_{\mathcal{B}} \le K(\|f\|_{\mathcal{B}} + \|g\|_{\mathcal{B}}).$$

Let $\beta \in (0, 1]$. As in [56, 57], a quasi-Banach space $\mathcal{B}_\beta$ with the quasi-norm $\|\cdot\|_{\mathcal{B}_\beta}$ is called a *β-quasi-Banach space* if

$$\|f + g\|^\beta_{\mathcal{B}_\beta} \le \|f\|^\beta_{\mathcal{B}_\beta} + \|g\|^\beta_{\mathcal{B}_\beta}$$

for all $f,\, g \in \mathcal{B}_\beta$.

Notice that any Banach space is a 1-quasi-Banach space, and the quasi-Banach spaces l^β and $L^\beta_\omega(\mathbb{R}^n)$ are typical β-quasi-Banach spaces. Let Φ satisfy Assumption (A). By the subadditivity of Φ and (2.6), we know that $h^\Phi_\omega(\mathbb{R}^n)$ is a p_Φ-quasi-Banach space.

For any given β-quasi-Banach space $\mathcal{B}_\beta$ with $\beta \in (0, 1]$ and a linear space $\mathcal{Y}$, an operator T from $\mathcal{Y}$ to $\mathcal{B}_\beta$ is called *$\mathcal{B}_\beta$-sublinear* if for any $f,\, g \in \mathcal{B}_\beta$ and $\lambda,\, \nu \in \mathbb{C}$,

$$\|T(\lambda f + \nu g)\|_{\mathcal{B}_\beta} \le \left(|\lambda|^\beta \|T(f)\|^\beta_{\mathcal{B}_\beta} + |\nu|^\beta \|T(g)\|^\beta_{\mathcal{B}_\beta}\right)^{1/\beta},$$

and

$$\|T(f) - T(g)\|_{\mathcal{B}_\beta} \le \|T(f - g)\|_{\mathcal{B}_\beta}$$

(see [56, 57]).

We remark that if T is linear, then T is $\mathcal{B}_\beta$-sublinear. Moreover, if $\mathcal{B}_\beta$ is a space of functions, and T is nonnegative and sublinear in the classical sense, then T is also $\mathcal{B}_\beta$-sublinear.

THEOREM 6.4. *Let Φ satisfy Assumption* (A)*, $\omega \in A^{\mathrm{loc}}_\infty(\mathbb{R}^n)$, q_ω be as in* (2.4) *and $(\rho,\, q,\, s)_\omega$ be admissible. Let $\mathcal{B}_\beta$ be a β-quasi-Banach space with $\beta \in (0, 1]$ and $\widetilde{p}$ be a upper type of Φ satisfying $\widetilde{p} \in (0, \beta]$. Suppose that one of the following holds:*

(i) *$q \in (q_\omega, \infty)$ and $T : h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n) \to \mathcal{B}_\beta$ is a $\mathcal{B}_\beta$-sublinear operator such that*

$$S \equiv \sup\big\{\|T(a)\|_{\mathcal{B}_\beta} :\ a \textit{ is any } (\rho,\, q,\, s)_\omega\textit{-atom with } \operatorname{supp}(a) \subset Q \textit{ and}$$
$$l(Q) \in (0, 2] \textit{ or } (\rho,\, q)_\omega \textit{ -single -atom}\big\} < \infty.$$

(ii) *T is a $\mathcal{B}_\beta$-sublinear operator defined on continuous $(\rho,\, \infty,\, s)_\omega$-atoms such that*

$$S \equiv \sup\{\|T(a)\|_{\mathcal{B}_\beta} :\ a \textit{ is any continuous } (\rho,\, \infty,\, s)_\omega\textit{-atom}\} < \infty.$$

Then there exists a unique bounded $\mathcal{B}_\beta$-sublinear operator $\widetilde{T}$ from $h^\Phi_\omega(\mathbb{R}^n)$ to $\mathcal{B}_\beta$ which extends T.

Proof. We first show Theorem 6.4 under its assumption (i). For any $f \in h_{\omega,\,\mathrm{fin}}^{\rho,\,q,\,s}(\mathbb{R}^n)$, by Theorem 6.2(i) and Remark 6.3(i), there exist a sequence $\{\lambda_j\}_{j=0}^l \subset \mathbb{C}$ with some $l \in \mathbb{N}$, a $(\rho,\,q)_\omega$-single-atom a_0 and $(\rho,\,q,\,s)_\omega$-atoms $\{a_j\}_{j=1}^l$ satisfying $\mathrm{supp}\,(a_j) \subset Q_j$ and $l(Q_j) \in (0,2]$ for $j \in \{1,\,2,\,\cdots,\,l\}$ such that $f = \sum_{j=0}^l \lambda_j a_j$ pointwise and

$$\Lambda\left(\{\lambda_j a_j\}_{j=0}^l\right) \lesssim \|f\|_{h_\omega^\Phi(\mathbb{R}^n)}. \tag{6.5}$$

Then by the assumptions, we see that

$$\begin{aligned}\|T(f)\|_{\mathcal{B}_\beta} &\le \left\{\sum_{i=0}^l |\lambda_i|^\beta \|T(a)\|_{\mathcal{B}_\beta}^\beta\right\}^{1/\beta} \le \left\{\sum_{i=0}^l |\lambda_i|^{\widetilde{p}} \|T(a)\|_{\mathcal{B}_\beta}^{\widetilde{p}}\right\}^{1/\widetilde{p}}\\ &\lesssim \left\{\sum_{i=0}^l |\lambda_i|^{\widetilde{p}}\right\}^{1/\widetilde{p}}.\end{aligned} \tag{6.6}$$

Since Φ is of upper type $\widetilde{p}$, then for any $t \in (0,1]$ and $s \in (0,\infty)$, we have $\Phi(st) \gtrsim t^{\widetilde{p}}\Phi(s)$. Let $\widetilde{\lambda}_0 \equiv \{\sum_{i=0}^l |\lambda_i|^{\widetilde{p}}\}^{1/\widetilde{p}}$. Then,

$$\sum_{i=0}^l \omega(Q_i)\Phi\left(\frac{|\lambda_i|}{\widetilde{\lambda}_0 \omega(Q_i)\rho(\omega(Q_i))}\right) \gtrsim \sum_{i=0}^l \omega(Q_i)\left(\frac{|\lambda_i|}{\widetilde{\lambda}_0}\right)^{\widetilde{p}} \frac{1}{\omega(Q_i)} \sim 1.$$

Thus, from this we deduce that $\widetilde{\lambda}_0 \lesssim \Lambda(\{\lambda_i a_i\}_{i=0}^l)$, which together with (6.5) and (6.6) implies that

$$\|T(f)\|_{\mathcal{B}_\beta} \lesssim \widetilde{\lambda}_0 \lesssim \Lambda\left(\{\lambda_i a_i\}_{i=0}^l\right) \lesssim \|f\|_{h_\omega^\Phi(\mathbb{R}^n)}.$$

Since $h_{\omega,\,\mathrm{fin}}^{\rho,\,q,\,s}(\mathbb{R}^n)$ is dense in $h_\omega^\Phi(\mathbb{R}^n)$, a density argument gives the desired conclusion in this case.

Now we prove Theorem 6.4 under its assumption (ii) by considering the following two cases for ω.

Case 1) $\omega(\mathbb{R}^n) = \infty$. In this case, similarly to the proof of (i), using Theorem 6.2(ii) and Remark 6.3(ii), we see that for all $f \in h_{\omega,\,\mathrm{fin}}^{\rho,\,\infty,\,s}(\mathbb{R}^n) \cap C(\mathbb{R}^n)$,

$$\|T(f)\|_{\mathcal{B}_\beta} \lesssim \|f\|_{h_\omega^\Phi(\mathbb{R}^n)}.$$

To extend T to the whole $h_\omega^\Phi(\mathbb{R}^n)$, we only need prove that $h_{\omega,\,\mathrm{fin}}^{\rho,\,\infty,\,s}(\mathbb{R}^n) \cap C(\mathbb{R}^n)$ is dense in $h_\omega^\Phi(\mathbb{R}^n)$. Since $h_{\omega,\,\mathrm{fin}}^{\rho,\,\infty,\,s}(\mathbb{R}^n)$ is dense in $h_\omega^\Phi(\mathbb{R}^n)$, it suffices to prove that $h_{\omega,\,\mathrm{fin}}^{\rho,\,\infty,\,s}(\mathbb{R}^n) \cap C(\mathbb{R}^n)$ is dense in $h_{\omega,\,\mathrm{fin}}^{\rho,\,\infty,\,s}(\mathbb{R}^n)$ with respect to the quasi-norm $\|\cdot\|_{h_\omega^\Phi(\mathbb{R}^n)}$.

To see this, let $f \in h_{\omega,\,\mathrm{fin}}^{\rho,\,\infty,\,s}(\mathbb{R}^n)$. In this case, for any $(\rho,\,\infty)_\omega$-single-atom b, $b(x) = 0$ for almost every $x \in \mathbb{R}^n$. Thus, f is a finite linear combination of $(\rho,\,\infty,\,s)_\omega$-atoms. Then there exists a cube $Q_0 \equiv Q(x_0, r_0)$ such that $\mathrm{supp}\,(f) \subset Q_0$. Take $\phi \in \mathcal{D}(\mathbb{R}^n)$ such that $\mathrm{supp}\,(\phi) \subset Q(0,1)$ and $\int_{\mathbb{R}^n} \phi(x)\,dx = 1$. Then it is easy to see that for any $k \in \mathbb{N}$, $\mathrm{supp}\,(\phi_k * f) \subset Q(x_0, r_0 + 1)$ and $\phi_k * f \in \mathcal{D}(\mathbb{R}^n)$. Assume that $f = \sum_{i=1}^N \lambda_i a_i$ with some $N \in \mathbb{N}$, $\{\lambda_i\}_{i=1}^N \subset \mathbb{C}$ and $\{a_i\}_{i=1}^N$ being $(\rho,\,\infty,\,s)_\omega$-atoms. Then for any $k \in \mathbb{N}$,

$$\phi_k * f = \sum_{i=1}^N \lambda_i \phi_k * a_i.$$

For any $k \in \mathbb{N}$ and $i \in \{1,\,2\,\cdots,\,N\}$, we now prove that $\phi_k * a_i$ is a multiple of some continuous $(\rho,\,\infty,\,s)_\omega$-atom, which implies that for any $k\in\mathbb{N}$,

$$\phi_k*f\in h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)\cap C(\mathbb{R}^n). \tag{6.7}$$

For $i\in\{1,\,2,\,\cdots,\,N\}$, assume that $\mathrm{supp}\,(a_i)\subset Q_i\equiv Q(x_i,r_i)$. Then

$$\mathrm{supp}\,(\phi_k*a_i)\subset \widetilde{Q}_{i,\,k}\equiv Q(x_i,r_i+1/2^k).$$

Moreover,

$$\|\phi_k*a_i\|_{L^\infty_\omega(\mathbb{R}^n)}\le\|a_i\|_{L^\infty_\omega(\mathbb{R}^n)}\le\frac{1}{\omega(Q_i)\rho(\omega(Q_i))}.$$

Furthermore, for any $\alpha\in\mathbb{Z}^n_+$, $\int_{\mathbb{R}^n}a_i(x)x^\alpha\,dx=0$ implies that

$$\int_{\mathbb{R}^n}\phi_k*a_i(x)x^\alpha\,dx=0.$$

Thus, $\frac{\omega(Q_i)\rho(\omega(Q_i))}{\omega(\widetilde{Q}_{i,\,k})\rho(\omega(\widetilde{Q}_{i,\,k}))}\phi_k*a_i$ is a $(\rho,\,\infty,\,s)_\omega$-atom.

Likewise, $\mathrm{supp}\,(f-\phi_k*f)\subset Q(x_0,r_0+1)$ and $f-\phi_k*f$ has the same vanishing moments as f. Take $q\in(q_\omega,\infty)$. By Lemma 2.6(iii),

$$\|f-\phi_k*f\|_{L^q_\omega(\mathbb{R}^n)}\to0\text{ as }k\to\infty. \tag{6.8}$$

Without loss of generality, we may assume that when k is large enough,

$$\|f-\phi_k*f\|_{L^q_\omega(\mathbb{R}^n)}>0.$$

Let

$$c_k\equiv\|f-\phi_k*f\|_{L^q_\omega(\mathbb{R}^n)}[\omega(Q(x_0,r_0+1))]^{1/q-1}\rho(\omega(Q(x_0,r_0+1)))$$

and $a_k\equiv\frac{f-\phi_k*f}{c_k}$. Then, a_k is a $(\rho,\,q,\,s)_\omega$-atom, $f-\phi_k*f=c_ka_k$, and $c_k\to0$ as $k\to\infty$. Thus, from (2.8) with $t\equiv\omega(Q(x_0,r_0+1))$, and Theorem 5.6, we infer that

$$\|f-\phi_k*f\|_{h^\Phi_\omega(\mathbb{R}^n)}\lesssim\Lambda(\{c_ka_k\})\lesssim|c_k|\to0 \tag{6.9}$$

as $k\to\infty$, which together with (6.7) shows the desired conclusion in this case.

Case 2) $\omega(\mathbb{R}^n)<\infty$. In this case, similarly to the proof of Case 1), by Theorem 6.2(ii), to finish the proof of (ii) in this case, it suffices to prove that $h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin},\,c}(\mathbb{R}^n)\cap C(\mathbb{R}^n)$ is dense in $h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$ in the quasi-norm $\|\cdot\|_{h^\Phi_\omega(\mathbb{R}^n)}$.

For any $f\in h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$, assume that

$$f\equiv\sum_{i=1}^{N_1}\lambda_ia_i+\lambda_0a_0,$$

where $N_1\in\mathbb{N}$, $\{\lambda_i\}_{i=1}^{N_1}\subset\mathbb{C}$ and a_0 is a $(\rho,\,\infty)_\omega$-single-atom and $\{a_i\}_{i=1}^{N_1}$ are $(\rho,\,\infty,\,s)_\omega$-atoms. Let $\{\psi_k\}_{k\in\mathbb{N}}\subset\mathcal{D}(\mathbb{R}^n)$ satisfy $0\le\psi_k\le1$, $\psi_k\equiv1$ on the cube $Q(0,2^k)$ and

$$\mathrm{supp}\,(\psi_k)\subset Q(0,2^{k+1}).$$

We assume that $\mathrm{supp}\,(\sum_{i=1}^{N_1}\lambda_ia_i)\subset Q(0,R_0)$ for some $R_0\in(0,\infty)$ and k_0 is the smallest integer such that $2^{k_0}\ge R_0$. For any integer $k\ge k_0$, let $f_k\equiv f\psi_k$. Then $f_k\in h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin},\,c}(\mathbb{R}^n)$. Indeed, by the choice of ψ_k, we know that

$$f_k=\sum_{i=1}^{N_1}\lambda_ia_i+\lambda_0a_0\psi_k$$

and $\operatorname{supp}(f_k) \subset Q(0, 2^{k+1})$. Furthermore, from $\operatorname{supp}(a_0\psi_k) \subset Q(0, 2^{k+1})$ and

$$\|a_0\psi_k\|_{L^\infty_\omega(\mathbb{R}^n)} \le \|a_0\|_{L^\infty_\omega(\mathbb{R}^n)} \le \frac{1}{\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))} \le \frac{1}{\omega(Q(0,2^{k+1}))\rho(\omega(Q(0,2^{k+1})))},$$

we deduce that $a_0\psi_k$ is a $(\rho,\,\infty,\,s)_\omega$-atom. Thus, $f_k \in h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin},\,c}(\mathbb{R}^n)$. For any fixed integer $k \ge k_0$ and any $i \in \mathbb{N}$, let $\widetilde{f}_{k,\,i} \equiv f_k * \phi_i$, where ϕ is as in Case 1). Similarly to the proof of (6.7), we have $\widetilde{f}_{k,\,i} \in h^{\rho,\,\infty,\,s}_{\omega,\,\mathrm{fin},\,c}(\mathbb{R}^n) \cap C(\mathbb{R}^n)$. For any $q \in (q_\omega, \infty)$, from the choice of f_k and $\omega(\mathbb{R}^n) < \infty$, we conclude that

$$\begin{aligned}\|f - f_k\|_{L^q_\omega(\mathbb{R}^n)} &\le \left\{\int_{Q(0,2^k)^\complement} |f(x)|^q \omega(x)\,dx\right\}^{1/q}\\ &\le \|\lambda_0 a_0\|_{L^\infty_\omega(\mathbb{R}^n)} \left\{\int_{Q(0,2^k)^\complement} \omega(x)\,dx\right\}^{1/q} \to 0, \qquad (6.10)\end{aligned}$$

as $k \to \infty$. Furthermore, for any fixed $k \in \mathbb{Z}$ with $k \ge k_0$, similarly to the proof of (6.8), we see that $\|f_k - \widetilde{f}_{k,\,i}\|_{L^q_\omega(\mathbb{R}^n)} \to 0$ as $i \to \infty$, which together with (6.10) implies that

$$\left\|f - \widetilde{f}_{k,\,i}\right\|_{L^q_\omega(\mathbb{R}^n)} \to 0$$

as $k,\, i \to \infty$. Without loss of generality, we may assume that when k and i are large enough, $\|f - \widetilde{f}_{k,\,i}\|_{L^q_\omega(\mathbb{R}^n)} > 0$. Let

$$c_{k,\,i} \equiv \|f - \widetilde{f}_{k,\,i}\|_{L^q_\omega(\mathbb{R}^n)} [\omega(\mathbb{R}^n)]^{1/q-1} \rho(\omega(\mathbb{R}^n))$$

and $a_{k,\,i} \equiv \frac{f - \widetilde{f}_{k,\,i}}{c_{k,\,i}}$. Then, $f - \widetilde{f}_{k,\,i} = c_{k,\,i} a_{k,\,i}$, $a_{k,\,i}$ is a $(\rho,\,q)_\omega$-single-atom and $c_{k,\,i} \to 0$ as $k,\, i \to \infty$. Then, similarly to the proof of (6.9), we know that $\|f - \widetilde{f}_{k,\,i}\|_{h^\Phi_\omega(\mathbb{R}^n)} \to 0$ as $k,\, i \to \infty$, which completes the proof of Case 2) and hence of Theorem 6.4. ∎

Remark 6.5. Let $p \in (0, 1]$. We point out that Theorems 6.2(i) and 6.4(i) when $\Phi(t) \equiv t^p$ for all $t \in (0, \infty)$ were obtained by Tang [49, Theorems 6.1 and 6.2]. Theorems 6.2(ii) and 6.4(ii) are new even when $\omega \equiv 1$ and $\Phi(t) \equiv t^p$ for all $t \in (0, \infty)$.

7. Dual spaces

In this section, we introduce the BMO-type space $\mathrm{bmo}^q_{\rho,\,\omega}(\mathbb{R}^n)$ and establish the duality between $h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)$ and $\mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$, where and in what follows, $\frac{1}{q} + \frac{1}{q'} = 1$. From this and Theorem 5.6, we deduce the duality between $h^\Phi_\omega(\mathbb{R}^n)$ and $\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$, and that for $q \in [1, \frac{q_\omega}{q_\omega - 1})$, $\mathrm{bmo}^q_{\rho,\,\omega}(\mathbb{R}^n) = \mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$ with equivalent norms, where the *symbol* $\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$ denotes $\mathrm{bmo}^1_{\rho,\,\omega}(\mathbb{R}^n)$. We begin with some notions.

For any locally integrable function f on $\mathbb{R}^n$, we denote the *minimizing polynomial* of f on the cube Q with degree at most s by $P^s_Q f$, namely, for all multi-indices $\theta \in \mathbb{Z}^n_+$ with $0 \le |\theta| \le s$,

$$\int_Q \left[f(x) - P^s_Q f(x)\right] x^\theta\,dx = 0. \qquad (7.1)$$

It is well known that if f is locally integrable, then $P^s_Q f$ uniquely exists; see, for example, [48]. Now, we introduce the BMO-type space $\mathrm{bmo}^q_{\rho,\,\omega}(\mathbb{R}^n)$ as follows.

Definition 7.1. Let Φ satisfy Assumption (A), $\omega \in A^{\mathrm{loc}}_{\infty}(\mathbb{R}^n)$, and q_ω, p_Φ and ρ be respectively as in (2.4), (2.6) and (2.7). Let $q \in [1, \frac{q_\omega}{q_\omega - 1})$ and $s \in \mathbb{Z}_+$ with $s \ge \lfloor n(\frac{q_\omega}{p_\Phi} - 1)\rfloor$. When $\omega(\mathbb{R}^n) = \infty$, a locally integrable function f on $\mathbb{R}^n$ is said to belong to the *space* $\mathrm{bmo}^q_{\rho,\,\omega}(\mathbb{R}^n)$, if

$$\begin{aligned}\|f\|_{\mathrm{bmo}^q_{\rho,\,\omega}(\mathbb{R}^n)} \equiv{} & \sup_{Q\subset\mathbb{R}^n,\,|Q|<1} \frac{1}{\rho(\omega(Q))}\left\{\frac{1}{\omega(Q)}\int_Q \left|f(x) - P^s_Q f(x)\right|^q [\omega(x)]^{1-q}\,dx\right\}^{1/q}\\ & + \sup_{Q\subset\mathbb{R}^n,\,|Q|\ge 1} \frac{1}{\rho(\omega(Q))}\left\{\frac{1}{\omega(Q)}\int_Q |f(x)|^q[\omega(x)]^{1-q}\,dx\right\}^{1/q} < \infty,\end{aligned}$$

where the supremum is taken over all the cubes $Q \subset \mathbb{R}^n$ and $P^s_Q f$ is as in (7.1). When $\omega(\mathbb{R}^n) < \infty$, a function f on $\mathbb{R}^n$ is said to belong to the *space* $\mathrm{bmo}^q_{\rho,\,\omega}(\mathbb{R}^n)$ if

$$\begin{aligned}\|f\|_{\mathrm{bmo}^q_{\rho,\,\omega}(\mathbb{R}^n)} \equiv{} & \sup_{Q\subset\mathbb{R}^n,\,|Q|<1} \frac{1}{\rho(\omega(Q))}\left\{\frac{1}{\omega(Q)}\int_Q \left|f(x) - P^s_Q f(x)\right|^q [\omega(x)]^{1-q}\,dx\right\}^{1/q}\\ & + \sup_{Q\subset\mathbb{R}^n,\,|Q|\ge 1} \frac{1}{\rho(\omega(Q))}\left\{\frac{1}{\omega(Q)}\int_Q |f(x)|^q[\omega(x)]^{1-q}\,dx\right\}^{1/q}\\ & + \frac{1}{\rho(\omega(\mathbb{R}^n))}\left\{\frac{1}{\omega(\mathbb{R}^n)}\int_{\mathbb{R}^n} |f(x)|^q[\omega(x)]^{1-q}\,dx\right\}^{1/q} < \infty,\end{aligned}$$

where the supremum is taken over all the cubes $Q \subset \mathbb{R}^n$ and $P^s_Q f$ is as in (7.1).

When $\omega \equiv 1$, $\Phi \equiv t$ for all $t \in (0, \infty)$ and $q = 1$, the space $\mathrm{bmo}^q_{\rho,\,\omega}(\mathbb{R}^n)$ is just the space $\mathrm{bmo}(\mathbb{R}^n)$ introduced in [18].

Now, we establish the duality between $h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$ and $\mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$. We begin with the notion of the weighted atomic Orlicz-Hardy space $H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$.

Definition 7.2. Let Φ satisfy Assumption (A), $\omega \in A^{\mathrm{loc}}_{\infty}(\mathbb{R}^n)$, ρ be as in (2.7) and $(\rho,\, q,\, s)_\omega$ be admissible. A function a on $\mathbb{R}^n$ is called an $H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$-*atom* if there exists a cube $Q \subset \mathbb{R}^n$ such that

(i) $\mathrm{supp}\,(a) \subset Q$;

(ii) $\|a\|_{L^q_\omega(\mathbb{R}^n)} \le [\omega(Q)]^{1/q-1}[\rho(\omega(Q))]^{-1}$;

(iii) $\int_{\mathbb{R}^n} a(x)x^\alpha\,dx = 0$ for all multi-indices $\alpha \in \mathbb{Z}^n_+$ with $|\alpha| \le s$.

The *weighted atomic Orlicz-Hardy space* $H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$ is defined to be the space of all $f \in \mathcal{D}'(\mathbb{R}^n)$ satisfying that $f = \sum_{i=1}^\infty \lambda_i a_i$ in $\mathcal{D}'(\mathbb{R}^n)$, where $\{a_i\}_{i\in\mathbb{N}}$ are $H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$-atoms with $\mathrm{supp}\,(a_i) \subset Q_i$, and $\{\lambda_i\}_{i\in\mathbb{N}} \subset \mathbb{C}$, satisfying

$$\sum_{i=1}^{\infty} \omega(Q_i)\Phi\left(\frac{|\lambda_i|}{\omega(Q_i)\rho(\omega(Q_i))}\right) < \infty.$$

Moreover, the *quasi-norm* of $f \in H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$ is defined by

$$\|f\|_{H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)} \equiv \inf\left\{\Lambda\left(\{\lambda_i a_i\}_{i=1}^\infty\right)\right\},$$

where the infimum is taken over all the decompositions of f as above and

$$\Lambda(\{\lambda_i a_i\}_{i=1}^{\infty}) \equiv \inf\left\{\lambda>0:\ \sum_{i=1}^{\infty}\omega(Q_i)\Phi\left(\frac{|\lambda_i|}{\lambda\omega(Q_i)\rho(\omega(Q_i))}\right)\le 1\right\}.$$

Furthermore, the *space* $H^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$ is defined to be the set of all finite linear combinations of $H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$-atoms.

Obviously, the space $H^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$ is dense in the space $H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$ with respect to the quasi-norm $\|\cdot\|_{H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)}$.

Definition 7.3. Let Φ satisfy Assumption (A), $\omega\in A^{\mathrm{loc}}_{\infty}(\mathbb{R}^n)$, q_ω, p_Φ and ρ be respectively as in (2.4), (2.6) and (2.7). Let $q\in[1,\frac{q_\omega}{q_\omega-1})$ and $s\in\mathbb{Z}_+$ with $s\ge\lfloor n(\frac{q_\omega}{p_\Phi}-1)\rfloor$. A locally integrable function f on $\mathbb{R}^n$ is said to belong to the *space* $\mathrm{BMO}^q_{\rho,\,\omega}(\mathbb{R}^n)$ if

$$\|f\|_{\mathrm{BMO}^q_{\rho,\,\omega}(\mathbb{R}^n)}\equiv\sup_{Q\subset\mathbb{R}^n}\frac{1}{\rho(\omega(Q))}\left\{\frac{1}{\omega(Q)}\int_Q\left|f(x)-P^s_Qf(x)\right|^q[\omega(x)]^{1-q}\,dx\right\}^{1/q}<\infty,$$

where the supremum is taken over all cubes $Q\subset\mathbb{R}^n$ and P^s_Qf is as in (7.1).

Now, we establish the duality between $H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$ and $\mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$.

Lemma 7.4. *Let Φ satisfy Assumption* (A), *$\omega\in A^{\mathrm{loc}}_{\infty}(\mathbb{R}^n)$, q_ω and ρ be respectively as in* (2.4) *and* (2.7), *and $(\rho,\,q,\,s)_\omega$ be admissible. Then $[H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*$, the dual space of $H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$, coincides with $\mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$ in the following sense.*

(i) *Let $g\in\mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$. Then the linear functional L, which is initially defined on $H^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$ by*

$$L(f)=\langle g,f\rangle,\tag{7.2}$$

has a unique extension to $H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$ with

$$\|L\|_{[H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*}\le C\|g\|_{\mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)},$$

where C is a positive constant independent of g.

(ii) *Conversely, for any $L\in[H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*$, there exists $g\in\mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$ such that* (7.2) *holds for all $f\in H^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$ and*

$$\|g\|_{\mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)}\le C\|L\|_{[H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*},$$

where C is a positive constant independent of L.

Proof. We borrow some ideas from [48] and [33, Theorm 4.1]. Let $(\rho,\,q,\,s)_\omega$ be an admissible triplet. First, we prove (i). Let a be an $H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$-atom with $\operatorname{supp}(a)\subset Q$ and $g\in\mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$. Then by the vanishing condition of a and Hölder's inequality, we have

$$\begin{aligned}\left|\int_{\mathbb{R}^n}a(x)g(x)\,dx\right|&=\left|\int_{\mathbb{R}^n}a(x)\left[g(x)-P^s_Qg(x)\right]\,dx\right|\\&\le\|a\|_{L^q_\omega(\mathbb{R}^n)}\left\{\int_Q\left|g(x)-P^s_Qg(x)\right|^{q'}[\omega(x)]^{1-q'}\,dx\right\}^{1/q'}\\&\le\|g\|_{\mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)},\end{aligned}\tag{7.3}$$

where $P_Q^s g$ is as in (7.1). Let

$$f=\sum_{i=1}^{k_0}\lambda_i a_i\in H_{\omega,\,\mathrm{fin}}^{\rho,\,q,\,s}(\mathbb{R}^n),$$

where $k_0\in\mathbb{N}$, $\{\lambda_i\}_{i=1}^{k_0}\subset\mathbb{C}$ and for $i\in\{1,\,2,\,\cdots,\,k_0\}$, a_i is an $H_\omega^{\rho,\,q,\,s}(\mathbb{R}^n)$-atom with $\mathrm{supp}\,(a_i)\subset Q_i$. Since Φ is concave and has upper type 1, by Remark 3.6(iii), we know that $\sum_i|\lambda_i|\lesssim\Lambda(\{\lambda_i a_i\}_{i=1}^{k_0})$, which together with (7.3) implies that

$$\begin{aligned}\left|\int_{\mathbb{R}^n}f(x)g(x)\,dx\right|&\le\sum_{i=1}^{k_0}|\lambda_i|\left|\int_{Q_i}a_i(x)g(x)\,dx\right|\\&\le\left\{\sum_{i=1}^{k_0}|\lambda_i|\right\}\|g\|_{\mathrm{BMO}^{q'}_{\rho,\omega}(\mathbb{R}^n)}\lesssim\Lambda\left(\{\lambda_i a_i\}_{i=1}^{k_0}\right)\|g\|_{\mathrm{BMO}^{q'}_{\rho,\omega}(\mathbb{R}^n)}.\end{aligned}$$

Thus, by the above estimate and the fact that $H_{\omega,\,\mathrm{fin}}^{\rho,\,q,\,s}(\mathbb{R}^n)$ is dense in $H_\omega^{\rho,\,q,\,s}(\mathbb{R}^n)$ with respect to the quasi-norm $\|\cdot\|_{H_\omega^{\rho,\,q,\,s}(\mathbb{R}^n)}$, we know that (i) holds.

To prove (ii), assume that $L\in[H_\omega^{\rho,\,q,\,s}(\mathbb{R}^n)]^*$. Let $Q\subset\mathbb{R}^n$ be a closed cube and

$$L^q_{\omega,\,s}(Q)\equiv\left\{f\in L^q_\omega(Q):\ \int_Q f(x)x^\alpha\,dx=0,\,\alpha\in\mathbb{Z}_+^n,\,|\alpha|\le s\right\},$$

where $f\in L^q_\omega(Q)$ means that $f\in L^q_\omega(\mathbb{R}^n)$ and $\mathrm{supp}\,(f)\in Q$. We first prove that

$$[H_\omega^{\rho,\,q,\,s}(\mathbb{R}^n)]^*\subset\left[L^q_{\omega,\,s}(Q)\right]^*. \tag{7.4}$$

Obviously, for any given $f\in L^q_{\omega,\,s}(Q)$,

$$a\equiv[\omega(Q)]^{1/q-1}[\rho(\omega(Q))]^{-1}\|f\|^{-1}_{L^q_\omega(Q)}f$$

is an $H_\omega^{\rho,\,q,\,s}(\mathbb{R}^n)$-atom. Thus, $f\in H_\omega^{\rho,\,q,\,s}(\mathbb{R}^n)$ and

$$\|f\|_{H_\omega^{\rho,\,q,\,s}(\mathbb{R}^n)}\le[\omega(Q)]^{1/q'}\rho(\omega(Q))\|f\|_{L^q_\omega(Q)},$$

which implies that for all $f\in L^q_{\omega,\,s}(Q)$,

$$|Lf|\le\|L\|\|f\|_{H_\omega^{\rho,\,q,\,s}(\mathbb{R}^n)}\le[\omega(Q)]^{1/q'}\rho(\omega(Q))\|L\|\|f\|_{L^q_\omega(Q)}.$$

That is, $L\in[L^q_{\omega,\,s}(Q)]^*$. Thus, (7.4) holds.

From (7.4), the Hahn-Banach theorem and the Riesz represent theorem, it follows that there exists a $\widetilde{g}\in L^{q'}_\omega(Q)$ such that for all $f\in L^q_{\omega,\,s}(Q)$,

$$Lf=\int_Q f(x)\widetilde{g}(x)\omega(x)\,dx, \tag{7.5}$$

where when $q=\infty$, we used the fact that $L^\infty_{\omega,\,s}(Q)\subset L^\gamma_{\omega,\,s}(Q)$ for any $\gamma\in[1,\infty)$ and $L^{\gamma'}_{\omega,\,s}(Q)\subset L^1_{\omega,\,s}(Q)$. Taking a sequence $\{Q_j\}_{j\in\mathbb{N}}$ of cubes such that for any $j\in\mathbb{N}$, $Q_j\subset Q_{j+1}$ and $\lim_{j\to\infty}Q_j=\mathbb{R}^n$. From the above result, it follows that for each Q_j, there exists a $\widetilde{g}_j\in L^{q'}_\omega(Q_j)$ such that for all $f\in L^q_{\omega,\,s}(Q_j)$,

$$Lf=\int_{Q_j}f(x)\widetilde{g}_j(x)\omega(x)\,dx. \tag{7.6}$$

Now, we construct a function g such that

$$Lf=\int_{Q_j} f(x)g(x)\,dx$$

for all $f\in L^q_{\omega,\,s}(Q_j)$ and all $j\in\mathbb{N}$. First, assume that $f\in L^q_{\omega,\,s}(Q_1)$. By (7.6), we know that there exists a $\widetilde{g}_1\in L^{q'}_\omega(Q_1)$ such that

$$Lf=\int_{Q_1} f(x)\widetilde{g}_1(x)\omega(x)\,dx.$$

Notice that $f\in L^q_{\omega,\,s}(Q_1)\subset L^q_{\omega,\,s}(Q_2)$. By (7.6) again, there exists a $\widetilde{g}_2\in L^{q'}_\omega(Q_2)$ such that

$$Lf=\int_{Q_1} f(x)\widetilde{g}_1(x)\omega(x)\,dx=\int_{Q_2} f(x)\widetilde{g}_2(x)\omega(x)\,dx,$$

which implies that for all $f\in L^q_{\omega,\,s}(Q_1)$,

$$\int_{Q_1} f(x)\left[\widetilde{g}_1(x)-\widetilde{g}_2(x)\right]\omega(x)\,dx=0. \tag{7.7}$$

For any given $h\in L^q_\omega(Q_1)$, let $f_1\equiv h-P^s_{Q_1}h$. Then by (7.1), we know that $f_1\in L^q_{\omega,\,s}(Q_1)$. For f_1, by (7.7), we have

$$\int_{Q_1}[h(x)-P^s_{Q_1}h(x)][\widetilde{g}_1(x)-\widetilde{g}_2(x)]\omega(x)\,dx=0,$$

which combined with the well-known fact that

$$\int_{Q_1} P^s_{Q_1}h(x)\left[\widetilde{g}_1(x)-\widetilde{g}_2(x)\right]\omega(x)\,dx=\int_{Q_1} h(x)P^s_{Q_1}((\widetilde{g}_1-\widetilde{g}_2)\omega)(x)\,dx$$

implies that

$$\int_{Q_1} h(x)\left\{[\widetilde{g}_1(x)-\widetilde{g}_2(x)]\omega(x)-P^s_{Q_1}((\widetilde{g}_1-\widetilde{g}_2)\omega)(x)\right\}\,dx=0. \tag{7.8}$$

For $j=1,\,2$, let $g_j\equiv\widetilde{g}_j\omega$. By (7.8), we know that for all $h\in L^q_\omega(Q_1)$,

$$\int_{Q_1} h(x)\left\{\frac{[g_1(x)-g_2(x)]-P^s_{Q_1}(g_1-g_2)(x)}{\omega(x)}\right\}\omega(x)\,dx=0,$$

which implies that for almost every $x\in Q_1$,

$$g_1(x)-g_2(x)=P^s_{Q_1}(g_1-g_2)(x).$$

Let

$$g(x)\equiv\begin{cases} g_1(x), & \text{when } x\in Q_1,\\ g_1(x)+P^s_{Q_1}(g_1-g_2)(x), & \text{when } x\in(Q_2\setminus Q_1).\end{cases}$$

It is easy to see that for any $f\in L^q_{\omega,\,s}(Q_j)$ with $j\in\{1,\,2\}$,

$$Lf=\int_{Q_j} f(x)g(x)\,dx. \tag{7.9}$$

In this way, we obtain a function g on $\mathbb{R}^n$ such that (7.9) holds for any $j\in\mathbb{N}$.

Finally, we show that $g\in \mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$ and for all $f\in H^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$,

$$Lf=\int_{\mathbb{R}^n} f(x)g(x)\,dx. \tag{7.10}$$

Indeed, for any $H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$-atom a, there exists a $j_0 \in \mathbb{N}$ such that $a \in L^q_{\omega,\,s}(Q_{j_0})$. By this and the fact that (7.9) holds for any $j \in \mathbb{N}$, we see that (7.10) holds for any $f \in H^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$. It remains to prove that $g \in \mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$. Take any cube $Q \subset \mathbb{R}^n$ as well as any $f \in L^q_{\omega}(Q)$ satisfying $\|f\|_{L^q_{\omega}(Q)} \le 1$ and $\mathrm{supp}\,(f) \subset Q$. Let

$$a \equiv \widetilde{C}^{-1}[\omega(Q)]^{1/q'}[\rho(\omega(Q))]^{-1}(f - P^s_Q f)\chi_Q, \tag{7.11}$$

where $\widetilde{C}$ is a positive constant. Obviously, $\mathrm{supp}\,(a) \subset Q$. We choose $\widetilde{C}$ such that a becomes an $H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$-atom. From the equality

$$La = \int_Q a(x)g(x)\,dx$$

and $L \in [H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*$, it follows that

$$|La| = \left|\int_Q a(x)[g(x) - P^s_Q g(x)]\,dx\right| \le \|L\|_{[H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*}. \tag{7.12}$$

By (7.11), (7.12) and (7.1), for all $f \in L^q_{\omega}(Q)$ with $\|f\|_{L^q_{\omega}(Q)} \le 1$, we see that

$$[\omega(Q)]^{1/q'}[\rho(\omega(Q))]^{-1}\left|\int_Q f(x)[g(x) - P^s_Q g(x)]\,dx\right| \lesssim \|L\|_{[H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*},$$

which implies that

$$[\omega(Q)]^{1/q'}[\rho(\omega(Q))]^{-1}\left\{\int_Q |g(x) - P^s_Q g(x)|^{q'}[\omega(x)]^{1-q'}\,dx\right\}^{1/q'} \lesssim \|L\|_{[H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*}.$$

Thus, $g \in \mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$ and $\|g\|_{\mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)} \lesssim \|L\|_{[H^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*}$. This finishes the proof of Lemma 7.4. ■

Now, we give the duality between $h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$ and $\mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$ by invoking Lemma 7.4.

THEOREM 7.5. *Let Φ satisfy Assumption* (A), *$\omega \in A^{\mathrm{loc}}_{\infty}(\mathbb{R}^n)$, q_ω and ρ be respectively as in* (2.4) *and* (2.7), *and $(\rho,\,q,\,s)_\omega$ be admissible. Then $[h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*$, the dual space of $h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$, coincides with $\mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$ in the following sense.*

(i) *Let $g \in \mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$. Then the linear functional L, which is initially defined on $h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$ by*

$$L(f) = \langle g, f\rangle, \tag{7.13}$$

has a unique extension to $h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$ with

$$\|L\|_{[h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*} \le C\|g\|_{\mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)},$$

where C is a positive constant independent of g.

(ii) *Conversely, for any $L \in [h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*$, there exists $g \in \mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$ such that* (7.13) *holds for all $f \in h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$ and*

$$\|g\|_{\mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)} \le C\|L\|_{[h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*},$$

where C is a positive constant independent of L.

Proof. Let $(\rho,\,q,\,s)_\omega$ be an admissible triplet. Obviously, the proof of (i) is similar to the proof of Lemma 7.4(i). We omit the details.

Now, we prove (ii) by considering the following two cases for ω.

Case i) $\omega(\mathbb{R}^n)=\infty$. In this case, let $Q\subset\mathbb{R}^n$ be a cube with $l(Q)\in[1,\infty)$. We first prove that

$$[h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*\subset[L^q_\omega(Q)]^*\,. \tag{7.14}$$

Obviously, for any given $f\in L^q_\omega(Q)$,

$$a\equiv[\omega(Q)]^{1/q-1}[\rho(\omega(Q))]^{-1}\|f\|^{-1}_{L^q_\omega(Q)}f\chi_Q$$

is a $(\rho,\,q,\,s)_\omega$-atom. Thus, $f\in h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)$ and

$$\|f\|_{h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)}\le[\omega(Q)]^{1/q'}\rho(\omega(Q))\|f\|_{L^q_\omega(Q)},$$

which implies that for any $L\in[h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*$,

$$|Lf|\le\|L\|_{[h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*}\|f\|_{h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)}\le[\omega(Q)]^{1/q'}\rho(\omega(Q))\|f\|_{L^q_\omega(Q)}\|L\|_{[h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*}.$$

That is, $L\in[L^q_\omega(Q)]^*$. Thus, (7.14) holds.

Now, assume that $L\in[h^{\rho,\,q,\,s}_{\omega}(\mathbb{R}^n)]^*$. Similarly to the proof of (7.5), we know that there exists a $\widetilde{g}\in L^{q'}_\omega(Q)$ such that for all $f\in L^q_\omega(Q)$,

$$Lf=\int_Q f(x)\widetilde{g}(x)\omega(x)\,dx.$$

Taking a sequence $\{Q_j\}_{j\in\mathbb{N}}$ of cubes such that for any $j\in\mathbb{N}$, $Q_j\subset Q_{j+1}$, $\lim_{j\to\infty}Q_j=\mathbb{R}^n$ and $l(Q_1)\in[1,\infty)$. From the above result, it follows that for each Q_j, there exists a $\widetilde{g}_j\in L^{q'}_\omega(Q_j)$ such that for all $f\in L^q_\omega(Q_j)$,

$$Lf=\int_{Q_j} f(x)\widetilde{g}_j(x)\omega(x)\,dx. \tag{7.15}$$

Now, we construct a function g on $\mathbb{R}^n$ such that

$$Lf=\int_{Q_j} f(x)g(x)\,dx$$

holds for all $f\in L^q_\omega(Q_j)$ and $j\in\mathbb{N}$. First, assume that $f\in L^q_\omega(Q_1)$. By (7.15), we know that there exists a $\widetilde{g}_1\in L^{q'}_\omega(Q_1)$ such that

$$Lf=\int_{Q_1} f(x)\widetilde{g}_1(x)\omega(x)\,dx.$$

Notice that $f\in L^q_\omega(Q_1)\subset L^q_\omega(Q_2)$. By (7.15) again, there exists a $\widetilde{g}_2\in L^{q'}_\omega(Q_2)$ such that

$$Lf=\int_{Q_1} f(x)\widetilde{g}_1(x)\omega(x)\,dx=\int_{Q_2} f(x)\widetilde{g}_2(x)\omega(x)\,dx,$$

which implies that for all $f\in L^q_\omega(Q_1)$,

$$\int_{Q_1} f(x)[\widetilde{g}_1(x)-\widetilde{g}_2(x)]\omega(x)\,dx=0.$$

Thus, we obtain that for almost every $x\in Q_1$, $\widetilde{g}_1(x)=\widetilde{g}_2(x)$. For $j=1,\,2$, let $g_j\equiv\widetilde{g}_j\omega$ and

$$g(x)\equiv\begin{cases}g_1(x), & \text{when } x\in Q_1,\\ g_2(x), & \text{when } x\in Q_2\setminus Q_1.\end{cases}$$

It is easy to see that for all $f \in L^q_\omega(Q_j)$ with $j \in \{1, 2\}$,

$$Lf = \int_{Q_j} f(x)g_j(x)\,dx. \tag{7.16}$$

Continuing in this way, we obtain a function g on $\mathbb{R}^n$ such that (7.16) holds for all $j \in \mathbb{N}$.

Finally, we show that $g \in \mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$ and for all $f \in h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$,

$$Lf = \int_{\mathbb{R}^n} f(x)g(x)\,dx. \tag{7.17}$$

Indeed, since $\omega(\mathbb{R}^n) = \infty$, all $(\rho,\,q)_\omega$-single-atoms are 0, and for any $(\rho,\,q,\,s)_\omega$-atom a, there exists a $j_0 \in \mathbb{N}$ such that $a \in L^q_\omega(Q_{j_0})$. By this and the fact that (7.16) holds for all $j \in \mathbb{N}$, we see that (7.17) holds.

Now, we prove that $g \in \mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$. Take any cube $Q \subset \mathbb{R}^n$ with $l(Q) \in [1, \infty)$ as well as any $f \in L^q_\omega(Q)$ with $\|f\|_{L^q_\omega(Q)} \le 1$. Let

$$a \equiv [\omega(Q)]^{-1/q'}[\rho(\omega(Q))]^{-1} f\chi_Q.$$

Then a is a $(\rho,\,q,\,s)_\omega$-atom and $\mathrm{supp}\,(a) \subset Q$. From the equality

$$La = \int_Q a(x)g(x)\,dx$$

and $L \in [h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*$, we deduce that

$$|La| = \left|\int_Q a(x)g(x)\,dx\right| \le \|L\|_{[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*}.$$

Thus, for any $f \in L^q_\omega(Q)$ with $\|f\|_{L^q_\omega(Q)} \le 1$, we have

$$[\omega(Q)]^{-1/q'}[\rho(\omega(Q))]^{-1}\left|\int_Q f(x)g(x)\,dx\right| \le \|L\|_{[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*},$$

which implies that

$$[\omega(Q)]^{-1/q'}[\rho(\omega(Q))]^{-1}\left\{\int_Q |g(x)|^{q'}[\omega(x)]^{1-q'}\,dx\right\}^{1/q'} \le \|L\|_{[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*}. \tag{7.18}$$

Furthermore, from $h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n) \supset H^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)$ and

$$\|f\|_{h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)} \le \|f\|_{H^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)}$$

for all $f \in H^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)$, we deduce that

$$[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^* \subset [H^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*$$

and $L|_{H^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)} \in [H^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*$. Since (7.17) holds for all $f \in H^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$, by Lemma 7.4(ii), we know that $g \in \mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$ and

$$\|g\|_{\mathrm{BMO}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)} \lesssim \|L\,|_{H^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)}\,\|_{[H^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*} \lesssim \|L\|_{[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*}.$$

Thus, this estimate together with (7.18) implies that $g \in \mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$ and

$$\|g\|_{\mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)} \lesssim \|L\|_{[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*},$$

which completes the proof of Theorem 7.5(ii) in Case i).

Case ii) $\omega(\mathbb{R}^n)<\infty$. In this case, let

$$\widetilde{h^{\rho,\,q,\,s}_\omega}(\mathbb{R}^n)\equiv\left\{f=\sum_{i=1}^\infty\lambda_i a_i \text{ in } \mathcal{D}'(\mathbb{R}^n):\ \text{For } i\in\mathbb{N},\ a_i \text{ is a } (\rho,\,q,\,s)_\omega\text{-atom},\right.$$
$$\left.\operatorname{supp}(a_i)\subset Q_i,\ \lambda_i\in\mathbb{C} \text{ and } \sum_{i=1}^\infty\omega(Q_i)\Phi\left(\frac{|\lambda_i|}{\omega(Q_i)\rho(\omega(Q_i))}\right)<\infty\right\}$$

and for all $f\in\widetilde{h^{\rho,\,q,\,s}_\omega}(\mathbb{R}^n)$,

$$\|f\|_{\widetilde{h^{\rho,\,q,\,s}_\omega}(\mathbb{R}^n)}\equiv\inf\left\{\Lambda\left(\{\lambda_i a_i\}_{i=1}^\infty\right)\right\},$$

where the infimum is taken over all the decompositions of f as above. For any $f\in L^1_{\mathrm{loc}}(\mathbb{R}^n)$, let

$$\|f\|_{\widetilde{\mathrm{bmo}^{q'}_{\rho,\,\omega}}(\mathbb{R}^n)}\equiv\sup_{Q\subset\mathbb{R}^n,\,|Q|<1}\frac{1}{\rho(\omega(Q))}\left\{\frac{1}{\omega(Q)}\int_Q|f(x)-P^s_Qf(x)|^{q'}[\omega(x)]^{1-q'}\,dx\right\}^{1/q'}$$
$$+\sup_{Q\subset\mathbb{R}^n,\,|Q|\ge1}\frac{1}{\rho(\omega(Q))}\left\{\frac{1}{\omega(Q)}\int_Q|f(x)|^{q'}[\omega(x)]^{1-q'}\,dx\right\}^{1/q'}$$

and

$$\widetilde{\mathrm{bmo}^{q'}_{\rho,\,\omega}}(\mathbb{R}^n)\equiv\left\{f\in L^1_{\mathrm{loc}}(\mathbb{R}^n):\ \|f\|_{\widetilde{\mathrm{bmo}^{q'}_{\rho,\,\omega}}(\mathbb{R}^n)}<\infty\right\}.$$

Similarly to the proofs of (i) and Case i), we conclude that

$$\left[\widetilde{h^{\rho,\,q,\,s}_\omega}(\mathbb{R}^n)\right]^*=\widetilde{\mathrm{bmo}^{q'}_{\rho,\,\omega}}(\mathbb{R}^n).\tag{7.19}$$

Now we claim that

$$[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*\subset[L^q_\omega(\mathbb{R}^n)]^*.\tag{7.20}$$

Indeed, for any $f\in L^q_\omega(\mathbb{R}^n)$, let

$$a\equiv[\omega(\mathbb{R}^n)]^{1/q-1}[\rho(\omega(\mathbb{R}^n))]^{-1}\|f\|^{-1}_{L^q_\omega(\mathbb{R}^n)}f.$$

Then a is a $(\rho,\,q)_\omega$-single-atom, which implies that $f\in h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)$ and

$$\|f\|_{h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)}\le[\omega(\mathbb{R}^n)]^{1/q'}\rho(\omega(\mathbb{R}^n))\|f\|_{L^q_\omega(\mathbb{R}^n)}.$$

Thus, for any given $L\in[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*$ and all $f\in L^q_\omega(\mathbb{R}^n)$, we have

$$|Lf|\le\|L\|_{[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*}\|f\|_{h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)}\le[\omega(\mathbb{R}^n)]^{1/q'}\rho(\omega(\mathbb{R}^n))\|f\|_{L^q_\omega(\mathbb{R}^n)}\|L\|_{[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*}.$$

That is, $L\in[L^q_\omega(\mathbb{R}^n)]^*$. Thus, (7.20) holds.

Now, assume that $L\in[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*$. From $\omega(\mathbb{R}^n)<\infty$, it follows that

$$L^\infty_\omega(\mathbb{R}^n)\subset L^\gamma_\omega(\mathbb{R}^n)$$

for any $\gamma\in[1,\infty)$ and $L^{\gamma'}_\omega(\mathbb{R}^n)\subset L^1_\omega(\mathbb{R}^n)$. By this, (7.20), the Hahn-Banach theorem and the Riesz represent theorem, we conclude that there exists a $\widetilde{g}\in L^{q'}_\omega(\mathbb{R}^n)$ such that for all $f\in L^q_\omega(\mathbb{R}^n)$ with $q\in(q_\omega,\infty]$,

$$Lf=\int_{\mathbb{R}^n}f(x)\widetilde{g}(x)\omega(x)\,dx.$$

Let $g \equiv \widetilde{g}\omega$. Then we know that for all $f \in L^q_\omega(\mathbb{R}^n)$,

$$Lf = \int_{\mathbb{R}^n} f(x)g(x)\,dx. \tag{7.21}$$

Finally, we prove that $g \in \mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$ and

$$\|g\|_{\mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)} \lesssim \|L\|_{[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*}.$$

Obviously, (7.21) holds for all $f \in h^{\rho,\,q,\,s}_{\omega,\,\mathrm{fin}}(\mathbb{R}^n)$. For any $f \in L^q_\omega(\mathbb{R}^n)$ with $\|f\|_{L^q_\omega(\mathbb{R}^n)} \le 1$, let

$$a \equiv [\omega(\mathbb{R}^n)]^{-1/q'}[\rho(\omega(\mathbb{R}^n))]^{-1}f.$$

Then a is a $(\rho,\,q)_\omega$-single-atom. From (7.21) with $f \equiv a$ and $L \in [h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*$, we deduce that

$$|La| = \left|\int_{\mathbb{R}^n} a(x)g(x)\,dx\right| \le \|L\|_{[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*}.$$

That is,

$$[\omega(\mathbb{R}^n)]^{-1/q'}[\rho(\omega(\mathbb{R}^n))]^{-1}\left|\int_{\mathbb{R}^n} f(x)g(x)\,dx\right| \le \|L\|_{[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*},$$

which together with $\|f\|_{L^q_\omega(\mathbb{R}^n)} \le 1$ implies that

$$[\omega(\mathbb{R}^n)]^{-1/q'}[\rho(\omega(\mathbb{R}^n))]^{-1}\left\{\int_{\mathbb{R}^n} |g(x)|^{q'}[\omega(x)]^{1-q'}\,dx\right\}^{1/q'} \le \|L\|_{[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*}. \tag{7.22}$$

Moveover, from $h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n) \supset \widetilde{h^{\rho,\,q,\,s}_\omega}(\mathbb{R}^n)$ and

$$\|f\|_{h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)} \le \|f\|_{\widetilde{h^{\rho,\,q,\,s}_\omega}(\mathbb{R}^n)}$$

for all $f \in \widetilde{h^{\rho,\,q,\,s}_\omega}(\mathbb{R}^n)$, we conclude that $[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^* \subset [\widetilde{h^{\rho,\,q,\,s}_\omega}(\mathbb{R}^n)]^*$ and

$$L\,|_{\widetilde{h^{\rho,\,q,\,s}_\omega}(\mathbb{R}^n)} \in \left[\widetilde{h^{\rho,\,q,\,s}_\omega}(\mathbb{R}^n)\right]^*.$$

Thus, by (7.19) and (7.21), we know that $g \in \widetilde{\mathrm{bmo}^{q'}_{\rho,\,\omega}}(\mathbb{R}^n)$ and

$$\|g\|_{\widetilde{\mathrm{bmo}^{q'}_{\rho,\,\omega}}(\mathbb{R}^n)} \lesssim \left\|L\,|_{\widetilde{h^{\rho,\,q,\,s}_\omega}(\mathbb{R}^n)}\right\|_{[\widetilde{h^{\rho,\,q,\,s}_\omega}(\mathbb{R}^n)]^*} \lesssim \|L\|_{[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*},$$

which together with (7.22) implies that $g \in \mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)$ and

$$\|g\|_{\mathrm{bmo}^{q'}_{\rho,\,\omega}(\mathbb{R}^n)} \lesssim \|L\|_{[h^{\rho,\,q,\,s}_\omega(\mathbb{R}^n)]^*}.$$

This finishes the proof of Theorem 7.5. ■

When $q = 1$, we denote $\mathrm{bmo}^q_{\rho,\,\omega}(\mathbb{R}^n)$ simply by $\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$. By Theorems 5.6 and 7.5, we have the following conclusions.

Corollary 7.6. *Let Φ satisfy Assumption* (A)*, $\omega \in A^{\mathrm{loc}}_\infty(\mathbb{R}^n)$, q_ω and ρ be respectively as in* (2.4) *and* (2.7)*. Then for $q \in [1, \frac{q_\omega}{q_\omega - 1})$, $\mathrm{bmo}^q_{\rho,\,\omega}(\mathbb{R}^n) = \mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$ with equivalent norms.*

Corollary 7.7. *Let Φ satisfy Assumption* (A)*, $\omega \in A^{\mathrm{loc}}_\infty(\mathbb{R}^n)$ and ρ be as in* (2.7)*. Then $[h^\Phi_\omega(\mathbb{R}^n)]^* = \mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$.*

8. Some applications

In this section, we first show that local Riesz transforms are bounded on $h^{\Phi}_{\omega}(\mathbb{R}^n)$. Moreover, we introduce local fractional integrals and show that they are bounded from $h^{p}_{\omega^p}(\mathbb{R}^n)$ to $L^{q}_{\omega^q}(\mathbb{R}^n)$ when $q \in [1,\infty)$, and from $h^{p}_{\omega^p}(\mathbb{R}^n)$ to $h^{q}_{\omega^q}(\mathbb{R}^n)$ when $q \in (0,1]$. Finally, we prove that some pseudo-differential operators are bounded on $h^{\Phi}_{\omega}(\mathbb{R}^n)$, where $\omega \in A_p(\phi)$ which was introduced by Tang [50] (see also Definition 8.13 below) and is contained in $A^{\rm loc}_p(\mathbb{R}^n)$ for $p \in [1,\infty)$.

Now, we recall the notion of local Riesz transforms introduced by Goldberg [18]. In what follows, $\mathcal{S}(\mathbb{R}^n)$ denotes the *space of all Schwartz functions on* $\mathbb{R}^n$.

Definition 8.1. Let $\phi_0 \in \mathcal{D}(\mathbb{R}^n)$ such that $\phi_0 \equiv 1$ on $Q(0,1)$ and $\mathrm{supp}\,(\phi_0) \subset Q(0,2)$. For $j \in \{1,\,2,\,\cdots,\,n\}$ and $x \in \mathbb{R}^n$, let

$$k_j(x) \equiv \frac{x_j}{|x|^{n+1}}\phi_0(x).$$

For $f \in \mathcal{S}(\mathbb{R}^n)$, the *local Riesz transform* $r_j(f)$ of f is defined by $r_j(f) \equiv k_j * f$.

We remark that ϕ_0 in [18] was assumed that $\phi_0 \equiv 1$ in a neighborhood of the origin and $\phi_0 \in \mathcal{D}(\mathbb{R}^n)$. In this paper, for convenience, we assume $\phi_0 \equiv 1$ on $Q(0,1)$ and $\mathrm{supp}\,(\phi_0) \subset Q(0,2)$. We have the boundedness on $h^{\Phi}_{\omega}(\mathbb{R}^n)$ of local Riesz transforms $\{r_j\}_j$ as follows.

Theorem 8.2. *Let Φ satisfy Assumption* (A), *$\omega \in A^{\rm loc}_{\infty}(\mathbb{R}^n)$ and p_Φ be as in* (2.6). *For $j \in \{1,\,2,\,\cdots,\,n\}$, let r_j be the local Riesz operator as in Definition 8.1. If $p_\Phi = p^+_\Phi$ and Φ is of upper type p^+_Φ, then there exists a positive constant $C_0 \equiv C_0(\Phi,\,\omega,\,n)$, depending only on Φ, q_ω, the weight constant of ω and n, such that for all $f \in h^{\Phi}_{\omega}(\mathbb{R}^n)$,*

$$\|r_j(f)\|_{h^{\Phi}_{\omega}(\mathbb{R}^n)} \le C_0\|f\|_{h^{\Phi}_{\omega}(\mathbb{R}^n)}.$$

To prove Theorem 8.2, we need the following lemma established in [49, Lemma 8.2].

Lemma 8.3. *For $j \in \{1,\,2,\,\cdots,\,n\}$, let r_j be the local Riesz operator as in Definition 8.1.*

(i) *For $\omega \in A^{\rm loc}_p(\mathbb{R}^n)$ with $p \in (1,\,\infty)$, then there exists a positive constant*

$$C_1 \equiv C_1(p,\,\omega,\,n),$$

depending only on p, the weight constant of ω, and n, such that for all $f \in L^p_\omega(\mathbb{R}^n)$,

$$\|r_j(f)\|_{L^p_\omega(\mathbb{R}^n)} \le C_1\|f\|_{L^p_\omega(\mathbb{R}^n)}.$$

(ii) *For $\omega \in A^{\rm loc}_1(\mathbb{R}^n)$, there exists a positive constant $C_2 \equiv C_2(\omega,\,n)$, depending only on the weight constant of ω, and n, such that for all $f \in L^1_\omega(\mathbb{R}^n)$,*

$$\|r_j(f)\|_{L^{1,\infty}_\omega(\mathbb{R}^n)} \le C_2\|f\|_{L^1_\omega(\mathbb{R}^n)}.$$

Now, we prove Theorem 8.2 by using Theorem 6.2 and Lemma 8.3.

Proof of Theorem 8.2. Let $s \equiv \lfloor n(q_\omega/p_\Phi - 1)\rfloor$, where q_ω and p_Φ are respectively as in (2.4) and (2.6). Then $(n+s+1)p_\Phi > nq_\omega$, which implies that there exists $q \in (q_\omega,\infty)$ such that $(n+s+1)p_\Phi > nq$ and $\omega \in A^{\rm loc}_q(\mathbb{R}^n)$. To show Theorem 8.2, by Theorem 6.4(i) and Theorem 3.2, it suffices to show that for any $(\rho,\,q)_\omega$-single-atom a or $(\rho,\,q,\,s)_\omega$-atom a supported in $Q(x_0,\,R_0)$ with $R_0\in(0,2]$,

$$\left\|\mathcal{G}_N^0\left(r_j(a)\right)\right\|_{L^\Phi_\omega(\mathbb{R}^n)}\lesssim 1. \tag{8.1}$$

First, we prove (8.1) for any $(\rho,\,q)_\omega$-single-atom $a\neq 0$. In this case, $\omega(\mathbb{R}^n)<\infty$. Since Φ is concave, by Jensen's inequality, Hölder's inequality, Proposition 3.2(ii), Lemma 8.3(i) and (2.8) with $t\equiv\omega(\mathbb{R}^n)$, we have

$$\begin{aligned}
&\int_{\mathbb{R}^n}\Phi\left(\mathcal{G}_N^0\left(r_j(a)\right)(x)\right)\omega(x)\,dx\\
&\quad\le\omega(\mathbb{R}^n)\Phi\left(\frac{1}{\omega(\mathbb{R}^n)}\int_{\mathbb{R}^n}\mathcal{G}_N^0(r_j(a))(x)\omega(x)\,dx\right)\\
&\quad\le\omega(\mathbb{R}^n)\Phi\left(\frac{1}{[\omega(\mathbb{R}^n)]^{1/q}}\left\{\int_{\mathbb{R}^n}\left[\mathcal{G}_N^0(r_j(a))(x)\right]^q\omega(x)\,dx\right\}^{1/q}\right)\\
&\quad\lesssim\omega(\mathbb{R}^n)\Phi\left(\frac{1}{[\omega(\mathbb{R}^n)]^{1/q}}\|r_j(a)\|_{L^q_\omega(\mathbb{R}^n)}\right)\lesssim\omega(\mathbb{R}^n)\Phi\left(\frac{1}{[\omega(\mathbb{R}^n)]^{1/q}}\|a\|_{L^q_\omega(\mathbb{R}^n)}\right)\\
&\quad\lesssim\omega(\mathbb{R}^n)\Phi\left(\frac{1}{\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))}\right)\sim 1,
\end{aligned}$$

which implies (8.1) in this case.

Now, let a be any $(\rho,\,q,\,s)_\omega$-atom supported in $Q_0\equiv Q(x_0,\,R_0)$ with $R_0\in(0,2]$. We prove (8.1) for a by considering the following two cases for R_0.

Case 1) $R_0\in[1,2]$. In this case, by the definitions of $r_j(a)$ and $\mathcal{G}_N^0(r_j(a))$, we see that

$$\operatorname{supp}\left(\mathcal{G}_N^0(r_j(a))\right)\subset Q_0^*\equiv Q(x_0,R_0+8).$$

From this, Jensen's inequality, Hölder's inequality, Proposition 3.2(ii), Lemmas 8.3 and 2.3(v), Remark 2.4 with $\widetilde{C}\equiv 2$ and (2.8) with $t\equiv\omega(Q_0)$, we infer that

$$\begin{aligned}
&\int_{\mathbb{R}^n}\Phi\left(\mathcal{G}_N^0(r_j(a))(x)\right)\omega(x)\,dx\\
&\quad\le\omega(Q_0^*)\Phi\left(\frac{1}{\omega(Q_0^*)}\int_{Q_0^*}\mathcal{G}_N^0(r_j(a))(x)\omega(x)\,dx\right)\\
&\quad\le\omega(Q_0^*)\Phi\left(\frac{1}{[\omega(Q_0^*)]^{1/q}}\left\{\int_{Q_0^*}\left[\mathcal{G}_N^0(r_j(a))(x)\right]^q\omega(x)\,dx\right\}^{1/q}\right)\\
&\quad\lesssim\omega(Q_0^*)\Phi\left(\frac{1}{[\omega(Q_0)]^{1/q}}\|r_j(a)\|_{L^q_\omega(\mathbb{R}^n)}\right)\lesssim\omega(Q_0)\Phi\left(\frac{1}{[\omega(Q_0)]^{1/q}}\|a\|_{L^q_\omega(\mathbb{R}^n)}\right)\\
&\quad\lesssim\omega(Q_0)\Phi\left(\frac{1}{\omega(Q_0)\rho(\omega(Q_0))}\right)\sim 1,
\end{aligned}$$

which implies (8.1) in Case 1).

Case 2) $R_0\in(0,1)$. In this case, let $\widetilde{Q}_0\equiv 8nQ_0$. Then

$$\begin{aligned}
\int_{\mathbb{R}^n}\Phi\left(\mathcal{G}_N^0(r_j(a))(x)\right)\omega(x)\,dx&=\int_{\widetilde{Q}_0}\Phi\left(\mathcal{G}_N^0(r_j(a))(x)\right)\omega(x)\,dx+\int_{(\widetilde{Q}_0)^\complement}\cdots\\
&\equiv\mathrm{I}_1+\mathrm{I}_2.
\end{aligned}\tag{8.2}$$

For I_1, similarly to the proof of Case 1), we have

$$\begin{aligned}\mathrm{I}_1 &\le \omega(\widetilde{Q}_0)\Phi\left(\frac{1}{[\omega(\widetilde{Q}_0)]^{1/q}}\left\{\int_{\widetilde{Q}_0}\left[\mathcal{G}_N^0(r_j(a))(x)\right]^q\omega(x)\,dx\right\}^{1/q}\right)\\ &\lesssim \omega(\widetilde{Q}_0)\Phi\left(\frac{1}{[\omega(Q_0)]^{1/q}}\|r_j(a)\|_{L^q_\omega(\mathbb{R}^n)}\right)\\ &\lesssim \omega(Q_0)\Phi\left(\frac{1}{\omega(Q_0)\rho(\omega(Q_0))}\right)\sim 1. \qquad (8.3)\end{aligned}$$

To estimate I_2, let $x \in (\widetilde{Q}_0)^\complement$, $t \in (0,1)$, $\psi \in \mathcal{D}_N^0(\mathbb{R}^n)$ and P_ψ^s be the Taylor expansion of ψ about $(x-x_0)/t$ with degree s. Then by the vanishing condition of a, we see that

$$\begin{aligned}|r_j(a)*\psi_t(x)| &= \frac{1}{t^n}\left|\int_{\mathbb{R}^n} r_j(a)(y)\psi\left(\frac{x-y}{t}\right)dy\right|\\ &= \frac{1}{t^n}\left|\int_{\mathbb{R}^n} r_j(a)(y)\left\{\psi\left(\frac{x-y}{t}\right)-P_\psi^s\left(\frac{x-y}{t}\right)\right\}dy\right|\\ &\le \frac{1}{t^n}\int_{2\sqrt{n}Q_0}|r_j(a)(y)|\left|\psi\left(\frac{x-y}{t}\right)-P_\psi^s\left(\frac{x-y}{t}\right)\right|dy\\ &\quad+\frac{1}{t^n}\int_{Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})\setminus(2\sqrt{n}Q_0)}\cdots+\frac{1}{t^n}\int_{Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})^\complement}\cdots\\ &\equiv \mathrm{G}_1+\mathrm{G}_2+\mathrm{G}_3. \qquad (8.4)\end{aligned}$$

To estimate G_1, by $t\in(0,1)$ and $x\in(\widetilde{Q}_0)^\complement$, we see that $\mathrm{G}_1 \neq 0$ implies that $t > \frac{3|x-x_0|}{4}$. From this, Taylor's remainder theorem, Hölder's inequality, Lemma 8.3(i), (2.1) and Remark 2.2(ii), we deduce that

$$\begin{aligned}\mathrm{G}_1 &\lesssim \frac{1}{t^{n+s+1}}\|r_j(a)\|_{L^q_\omega(\mathbb{R}^n)}\left\{\sum_{\alpha\in\mathbb{Z}^n_+,\,|\alpha|=s+1}\int_{2\sqrt{n}Q_0}\left|(\partial^\alpha\psi)\left(\frac{\xi}{t}\right)\right|^{q'}\right.\\ &\quad\left.\times|y-x_0|^{(s+1)q'}[\omega(y)]^{-q'/q}\,dy\right\}^{1/q'}\\ &\lesssim \frac{R_0^{s+1}}{|x-x_0|^{n+s+1}}\|a\|_{L^q_\omega(\mathbb{R}^n)}\left\{\int_{2\sqrt{n}Q_0}[\omega(y)]^{-q'/q}\,dy\right\}^{1/q'}\\ &\lesssim \frac{1}{\omega(Q_0)\rho(\omega(Q_0))}\frac{R_0^{n+s+1}}{|x-x_0|^{n+s+1}}, \qquad (8.5)\end{aligned}$$

where $\theta\in(0,1)$, $\xi\equiv\theta(x-y)+(1-\theta)(x-x_0)$ and $\frac{1}{q}+\frac{1}{q'}=1$.

To estimate G_2, by the definition of k_j with $j\in\{1,\,2,\,\cdots,\,n\}$, we have

$$\sum_{\alpha\in\mathbb{Z}^n_+,\,|\alpha|=s+1}|(\partial^\alpha k_j)(z)| \lesssim \frac{1}{|z|^{n+s+1}} \qquad (8.6)$$

for all $z\in\mathbb{R}^n\setminus\{0\}$. For any fixed $y\in(Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})\setminus 2\sqrt{n}Q_0)$, let K_j^s be the Taylor expansion of $k_j(\cdot)$ at the point $y-x_0$ with degree s. Moreover, it is easy to see that $\mathrm{G}_2\neq 0$ implies that $t>\frac{|x-x_0|}{2}$. From this, Taylor's remainder theorem, (8.6), Hölder's inequality, Lemma 8.3(i) and (2.1), we conclude that

$$\begin{aligned}
\mathrm{G}_2 &\le \frac{1}{t^n}\int_{Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})\setminus(2\sqrt{n}Q_0)}\left\{\int_{Q_0}|a(z)||k_j(y-z)-K_j^s(y-z)|\,dz\right\}\\
&\quad\times\left|\psi\left(\frac{x-y}{t}\right)-P_\psi^s\left(\frac{x-y}{t}\right)\right|\,dy\\
&\lesssim \frac{1}{t^{n+s+1}}\int_{Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})\setminus(2\sqrt{n}Q_0)}\left\{\int_{Q_0}|a(z)|\frac{|z-x_0|^{s+1}}{|\xi|^{n+s+1}}\,dz\right\}\\
&\quad\times|y-x_0|^{s+1}\,dy\\
&\lesssim \frac{1}{|x-x_0|^{n+s+1}}\int_{Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})\setminus(2\sqrt{n}Q_0)}\frac{1}{|y-x_0|^n}\left\{\int_{Q_0}|a(z)||z-x_0|^{s+1}\,dz\right\}dy\\
&\lesssim \frac{R_0^{s+1}}{|x-x_0|^{n+s+1}}\|a\|_{L^q_\omega(\mathbb{R}^n)}\frac{|Q_0|}{[\omega(Q_0)]^{1/q}}\int_{Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})\setminus(2\sqrt{n}Q_0)}\frac{1}{|y-x_0|^n}\,dy\\
&\lesssim \frac{1}{\omega(Q_0)\rho(\omega(Q_0))}\frac{R_0^{n+s+1}}{|x-x_0|^{n+s+1}}\int_{\sqrt{n}R_0}^{\frac{|x-x_0|}{2\sqrt{n}}}z^{-1}\,dz\\
&\lesssim \frac{1}{\omega(Q_0)\rho(\omega(Q_0))}\frac{R_0^{n+s+1-\delta}}{|x-x_0|^{n+s+1-\delta}},
\end{aligned}\tag{8.7}$$

where $\xi\equiv\gamma(y-z)+(1-\gamma)(y-x_0)$ for some $\gamma\in(0,1)$, δ is a small positive constant which is determined later, and in the third inequality we used the fact that for any $y\in Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})\setminus(2\sqrt{n}Q_0)$ and $z\in Q_0$,

$$|(y-x_0)-\gamma(z-x_0)|\ge|y-x_0|-|y-x_0|/2=|y-x_0|/2.$$

Finally, we estimate G_3. For any $y\in[Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})]^\complement$, by the definition of P_ψ^s and the support condition ψ, we have

$$\frac{1}{t^n}\left|P_\psi^s\left(\frac{x-y}{t}\right)\right|\lesssim\frac{|y-x_0|^s}{|x-x_0|^{n+s}}.\tag{8.8}$$

Thus, from the vanishing condition of a, Taylor's remainder theorem, (8.6), Hölder's inequality, (2.1) and (8.8), we deduce that

$$\begin{aligned}
\mathrm{G}_3 &\lesssim \frac{1}{t^n}\int_{Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})^\complement}\left\{\int_{Q_0}|a(z)||k_j(y-z)-K_j^s(y-z)|\,dz\right\}\\
&\quad\times\left\{\left|\psi\left(\frac{x-y}{t}\right)\right|+\left|P_\psi^s\left(\frac{x-y}{t}\right)\right|\right\}dy\\
&\lesssim \frac{1}{t^n}\int_{Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})^\complement}\left\{\int_{Q_0}|a(z)|\frac{|z-x_0|^{s+1}}{|\xi|^{n+s+1}}\,dz\right\}\\
&\quad\times\left\{\left|\psi\left(\frac{x-y}{t}\right)\right|+\left|P_\psi^s\left(\frac{x-y}{t}\right)\right|\right\}dy\\
&\lesssim \|a\|_{L^q_\omega(\mathbb{R}^n)}\frac{R_0^{s+n+1}}{[\omega(Q_0)]^{1/q}}\frac{1}{t^n}\int_{Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})^\complement}\frac{1}{|y-x_0|^{n+s+1}}
\end{aligned}$$

$$
\begin{aligned}
&\times\left\{\left|\psi\left(\frac{x-y}{t}\right)\right|+\left|P_\psi^s\left(\frac{x-y}{t}\right)\right|\right\}dy\\
&\lesssim \|a\|_{L^q_\omega(\mathbb{R}^n)}\frac{R_0^{s+n+1}}{[\omega(Q_0)]^{1/q}}\left\{\frac{1}{|x-x_0|^{n+s+1}}\frac{1}{t^n}\int_{Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})^{\complement}}\left|\psi\left(\frac{x-y}{t}\right)\right|dy\right.\\
&\quad\left.+\frac{1}{t^n}\int_{Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})^{\complement}}\frac{1}{|y-x_0|^{n+s+1}}\left|P_\psi^s\left(\frac{x-y}{t}\right)\right|dy\right\}\\
&\lesssim \frac{1}{\omega(Q_0)\rho(\omega(Q_0))}\left\{\frac{R_0^{n+s+1}}{|x-x_0|^{n+s+1}}\right.\\
&\quad\left.+\frac{R_0^{n+s+1}}{|x-x_0|^{n+s}}\int_{Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})^{\complement}}\frac{1}{|y-x_0|^{n+1}}\,dy\right\}\\
&\lesssim \frac{1}{\omega(Q_0)\rho(\omega(Q_0))}\frac{R_0^{n+s+1}}{|x-x_0|^{n+s+1}}, \qquad (8.9)
\end{aligned}
$$

where $\xi\equiv\gamma(y-z)+(1-\gamma)(y-x_0)$ for some $\gamma\in(0,1)$ and in the third inequality we used the fact that for any $y\in[Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})]^{\complement}$ and $z\in Q_0$,

$$|(y-x_0)-\gamma(z-x_0)|\gtrsim|y-x_0|.$$

Thus, by (8.4), (8.5), (8.7), (8.9) and $|x-x_0|\ge 4nR_0$, we know that

$$
\begin{aligned}
|r_j(a)*\psi_t(x)|&\lesssim\frac{1}{\omega(Q_0)\rho(\omega(Q_0))}\left\{\frac{R_0^{(n+s+1)}}{|x-x_0|^{n+s+1}}+\frac{R_0^{(n+s+1-\delta)}}{|x-x_0|^{n+s+1-\delta}}\right\}\\
&\lesssim\frac{1}{\omega(Q_0)\rho(\omega(Q_0))}\frac{R_0^{(n+s+1-\delta)}}{|x-x_0|^{n+s+1-\delta}},
\end{aligned}
$$

which together with the arbitrariness of $\psi\in\mathcal{D}_N^0(\mathbb{R}^n)$ implies that for all $x\in(\widetilde{Q}_0)^{\complement}$,

$$\mathcal{G}_N^0\left(r_j(a)\right)(x)\lesssim\frac{1}{\omega(Q_0)\rho(\omega(Q_0))}\frac{R_0^{(n+s+1-\delta)}}{|x-x_0|^{n+s+1-\delta}}. \qquad (8.10)$$

Take $\delta\in(0,\infty)$ small enough such that $p_\Phi(n+s+1-\delta)>nq$. By the fact that

$$\operatorname{supp}\left(\mathcal{G}_N^0(r_j(a))\right)\subset Q(x_0,R_0+8)\subset Q(x_0,9)$$

and Lemma 2.3(i), we know that there exists an $\widetilde{\omega}\in A_q(\mathbb{R}^n)$ such that $\widetilde{\omega}=\omega$ on $Q(x_0,9)$. Let m_0 be the integer such that $2^{m_0-1}nR_0\le 9<2^{m_0}nR_0$. By (8.10), the lower type p_Φ property of Φ, Lemma 2.3(viii) and $p_\Phi(n+s+1-\delta)>nq$, we conclude that

$$
\begin{aligned}
\mathrm{I}_2&\lesssim\int_{Q(x_0,9)\setminus\widetilde{Q}_0}\Phi\left(\mathcal{G}_N^0(r_j(a))(x)\right)\widetilde{\omega}(x)\,dx\\
&\lesssim\sum_{j=3}^{m_0}\int_{2^{j+1}nQ_0\setminus 2^jnQ_0}\Phi\left(\frac{1}{\omega(Q_0)\rho(\omega(Q_0))}\frac{R_0^{(n+s+1-\delta)}}{|x-x_0|^{n+s+1-\delta}}\right)\widetilde{\omega}(x)\,dx\\
&\lesssim\frac{1}{\omega(Q_0)}\sum_{j=3}^{m_0}\int_{2^{j+1}nQ_0\setminus 2^jnQ_0}\left(\frac{R_0^{n+s+1-\delta}}{|x-x_0|^{n+s+1-\delta}}\right)^{p_\Phi}\widetilde{\omega}(x)\,dx
\end{aligned}
$$

$$\lesssim \sum_{j=3}^{m_0} 2^{k[(n+s+1-\delta)p_\Phi - nq]} \lesssim 1,$$

which together with (8.2) and (8.3) implies (8.1) in Case 2). This finishes the proof of Theorem 8.2. ■

Remark 8.4. Theorem 8.2 when $\omega \in A_1^{\mathrm{loc}}(\mathbb{R}^n)$ and $\Phi(t) \equiv t$ for all $t \in (0,\infty)$ was obtained by Tang [49, Lemma 8.3].

Next, we introduce the local fractional integral and, using Theorem 6.4, prove that they are boundedness from $h^p_{\omega^p}(\mathbb{R}^n)$ to $L^q_{\omega^q}(\mathbb{R}^n)$ when $q \in [1,\infty)$, and from $h^p_{\omega^p}(\mathbb{R}^n)$ to $h^q_{\omega^q}(\mathbb{R}^n)$ when $q \in (0,1]$, provided that ω satisfies $\omega^{\frac{nr}{nr-n-r\alpha}} \in A_1^{\mathrm{loc}}(\mathbb{R}^n)$ for some $r \in (\frac{n}{n-\alpha},\infty)$ and $\int_{\mathbb{R}^n}[\omega(x)]^p\,dx = \infty$. We begin with some notions.

Definition 8.5. Let $\alpha \in [0,n)$ and ϕ_0 be as in Definition 8.1. For any $f \in \mathcal{S}(\mathbb{R}^n)$ and all $x \in \mathbb{R}^n$, the *local fractional integral* $I_\alpha^{\mathrm{loc}}(f)$ of f is defined by

$$I_\alpha^{\mathrm{loc}}(f)(x) \equiv \int_{\mathbb{R}^n} \frac{\phi_0(y)}{|y|^{n-\alpha}} f(x-y)\,dy.$$

Definition 8.6. (i) If there exist $r \in (1,\infty)$ and a positive constant C such that for all cubes $Q \subset \mathbb{R}^n$ with sidelength $l(Q) \in (0,1]$,

$$\left(\frac{1}{|Q|}\int_Q [\omega(x)]^r\,dx\right)^{1/r} \le C\left(\frac{1}{|Q|}\int_Q \omega(x)\,dx\right), \tag{8.11}$$

then ω is said to satisfy the *local reverse Hölder inequality of order r,* which is denoted by $\omega \in RH_r^{\mathrm{loc}}(\mathbb{R}^n)$. Furthermore, let $RH_r^{\mathrm{loc}}(\omega) \equiv \inf C$, where the infimum is taken over all the positive constants C satisfying (8.11).

(ii) Let $p,\ q \in (1,\infty)$. A locally integrable nonnegative function ω on $\mathbb{R}^n$ is said to belong to the *class* $A^{\mathrm{loc}}(p,\ q)$, if there exists a positive constant C such that for all cubes $Q \subset \mathbb{R}^n$ with sidelength $l(Q) \in (0,1]$,

$$\left(\frac{1}{|Q|}\int_Q [\omega(x)]^q\,dx\right)^{1/q}\left(\frac{1}{|Q|}\int_Q [\omega(x)]^{-p'}\,dx\right)^{1/p'} \le C, \tag{8.12}$$

where and in what follows, $\frac{1}{p}+\frac{1}{p'}=1$. Furthermore, let $A^{\mathrm{loc}}(p,\ q)(\omega) \equiv \inf\{C\}$, where the infimum is taken over all the positive constants C satisfying (8.12).

Remark 8.7. (i) Let r be as in Definition 8.6(i). If (8.11) holds for all cubes $Q \subset \mathbb{R}^n$, then ω is said to satisfy the *reverse Hölder inequality of order r,* which is denoted by $\omega \in RH_r(\mathbb{R}^n)$ (see, for example, [17]). Let p, q be as in Definition 8.6(ii). If (8.12) holds for all cubes $Q \subset \mathbb{R}^n$, then ω is said to belong to the *class* $A(p,\ q)$.

(ii) For any given positive constant A_1, let the cube Q satisfy $l(Q) = A_1$. Similarly to the proof of Lemma 2.3(i), we have that for any $\omega \in RH_r^{\mathrm{loc}}(\mathbb{R}^n)$, there exists an $\widetilde{\omega} \in RH_r(\mathbb{R}^n)$ such that $\omega = \widetilde{\omega}$ on Q and $RH_r(\widetilde{\omega}) \lesssim RH_r^{\mathrm{loc}}(\omega)$, where $RH_r(\widetilde{\omega})$ is defined similarly to $RH_r^{\mathrm{loc}}(\omega)$ and the implicit constant depends only on A_1 and n.

(iii) Similarly to Remark 2.2(ii), for any given constant $A_2 \in (0,\infty)$, the condition $l(Q) \in (0,1]$ in (8.11) can be replaced by $l(Q) \in (0,A_2]$ with the positive constant C in (8.11) depending on A_2.

About the relations of $A_\infty^{\rm loc}(\mathbb{R}^n)$, $RH_r^{\rm loc}(\mathbb{R}^n)$ and $A^{\rm loc}(p,\,q)$, we have the following conclusions.

Lemma 8.8. (i) *Let* $r\in(1,\,\infty)$. *Then* $\omega^r\in A_\infty^{\rm loc}(\mathbb{R}^n)$ *if and only if* $\omega\in RH_r^{\rm loc}(\mathbb{R}^n)$.

(ii) *Let* $\alpha\in(0,\,n)$, $p\in(1,\,n/\alpha)$ *and* $1/q=1/p-\alpha/n$. *Then* $\omega\in A^{\rm loc}(p,\,q)$ *if and only if* $\omega^{-p'}\in A^{\rm loc}_{1+p'/q}(\mathbb{R}^n)$.

Proof. We first prove (i). Let $\omega^r\in A_\infty^{\rm loc}(\mathbb{R}^n)$. Then by Lemma 2.3(i), we know that for any cube $Q\equiv Q(x_0,l(Q))$ with $l(Q)\in(0,1]$, there exists a function $\widetilde{\omega}$ on $\mathbb{R}^n$ such that

$$\widetilde{\omega}^r\in A_\infty(\mathbb{R}^n)\text{ and }\widetilde{\omega}=\omega\text{ on }Q(x_0,1). \tag{8.13}$$

Moreover, by [12, Lemma A], we know that

$$\widetilde{\omega}^r\in A_\infty(\mathbb{R}^n)\text{ if and only if }\widetilde{\omega}\in RH_r(\mathbb{R}^n). \tag{8.14}$$

Thus, for any cube $Q(x_0,l(Q))$ with $l(Q)\in(0,1]$, by (8.13) and (8.14), we have

$$\left(\frac{1}{|Q|}\int_Q[\omega(x)]^r\,dx\right)^{1/r}=\left(\frac{1}{|Q|}\int_Q[\widetilde{\omega}(x)]^r\,dx\right)^{1/r}\lesssim\frac{1}{|Q|}\int_Q\widetilde{\omega}(x)\,dx\sim\frac{1}{|Q|}\int_Q\omega(x)\,dx,$$

which together with the arbitrariness of the cube $Q(x_0,l(Q))$ implies that $\omega\in RH_r^{\rm loc}(\mathbb{R}^n)$.

Conversely, let $\omega\in RH_r^{\rm loc}(\mathbb{R}^n)$. Then by Remark 8.7(ii), we know that for any cube $Q(x_0,l(Q))$ with $l(Q)\in(0,1]$, there exists a function $\widetilde{\omega}$ on $\mathbb{R}^n$ such that $\widetilde{\omega}\in RH_r(\mathbb{R}^n)$ and $\widetilde{\omega}=\omega$ on $Q(x_0,1)$, which together with (8.14) and the arbitrariness of the cube $Q(x_0,l(Q))$ implies that $\omega\in A_\infty^{\rm loc}(\mathbb{R}^n)$. This finishes the proof of (i).

By the definitions of $A^{\rm loc}(p,\,q)$ and $A^{\rm loc}_{1+p'/q}(\mathbb{R}^n)$, we see that (ii) holds, which completes the proof of Lemma 8.8. ■

To establish the boundedness of local fractional integrals, we need the following technical lemma.

Lemma 8.9. *Let* $\alpha\in(0,n)$, $p\in(1,\frac{n}{\alpha})$ *and* $\frac{1}{q}=\frac{1}{p}-\frac{\alpha}{n}$. *For some* $r\in(q,\infty)$, *if*

$$\omega^{-r'}\in A^{\rm loc}(q'/r',\,p'/r'),$$

then there exists a positive constant C *such that for all* $f\in L^p_{\omega^p}(\mathbb{R}^n)$,

$$\left\|I_\alpha^{\rm loc}(f)\right\|_{L^q_{\omega^q}(\mathbb{R}^n)}\le C\|f\|_{L^p_{\omega^p}(\mathbb{R}^n)}, \tag{8.15}$$

where p', q' *and* r' *respectively denote the conjugate indices of* p, q *and* r.

Proof. Let $\omega^{-r'}\in A^{\rm loc}(q'/r',\,p'/r')$. For any unit cube $Q\subset\mathbb{R}^n$, from Lemmas 8.8(ii) and 2.3(i), and Remark 2.4, we deduce that there exists a function $\widetilde{\omega}$ on $\mathbb{R}^n$ such that $\widetilde{\omega}^{-r'}\in A(q'/r',\,p'/r')$ and $\widetilde{\omega}=\omega$ on $5Q$. For $\widetilde{\omega}^{-r'}\in A(q'/r',\,p'/r')$, similarly to the proof of [13, Theorem 2], we know that for all $f\in L^p_{\widetilde{\omega}^p}(\mathbb{R}^n)$,

$$\left\|I_\alpha^{\rm loc}(f)\right\|_{L^q_{\widetilde{\omega}^q}(\mathbb{R}^n)}\lesssim\|f\|_{L^p_{\widetilde{\omega}^p}(\mathbb{R}^n)},$$

which combined with the definition of $I_\alpha^{\rm loc}(f)$ implies that

$$\left\|I_\alpha^{\rm loc}(f)\right\|_{L^q_{\omega^q}(Q)}=\left\|I_\alpha^{\rm loc}(f\chi_{5Q})\right\|_{L^q_{\widetilde{\omega}^q}(Q)}\lesssim\|f\chi_{5Q}\|_{L^p_{\widetilde{\omega}^p}(\mathbb{R}^n)}\sim\|f\|_{L^p_{\omega^p}(5Q)}. \tag{8.16}$$

Take unit cubes $\{Q_i\}_{i=1}^{\infty}$ such that $\cup_{i=1}^{\infty} Q_i = \mathbb{R}^n$, their interiors are disjoint and

$$\sum_{i=1}^{\infty} \chi_{5Q_i} \le M,$$

where M is a positive integer depending only on n. From this and (8.16), we infer that

$$\left\|I_\alpha^{\mathrm{loc}}(f)\right\|_{L^q_{\omega^q}(\mathbb{R}^n)}^q = \sum_{i=1}^{\infty} \left\|I_\alpha^{\mathrm{loc}}(f)\right\|_{L^q_{\omega^q}(Q_i)}^q \lesssim \sum_{i=1}^{\infty} \|f\|_{L^p_{\omega^p}(5Q_i)}^q \lesssim \|f\|_{L^p_{\omega^p}(\mathbb{R}^n)}^q,$$

which implies (8.15). This finishes the proof of Lemma 8.9. ■

Theorem 8.10. *Let $\alpha \in (0,n)$, $p \in [\frac{n}{n+\alpha}, 1]$ and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$. For some $r \in (\frac{n}{n-\alpha}, \infty)$, if the weight ω satisfies $\omega^{\frac{nr}{nr-n-r\alpha}} \in A_1^{\mathrm{loc}}(\mathbb{R}^n)$ and $\int_{\mathbb{R}^n} [\omega(x)]^p\,dx = \infty$, then there exists a positive constant C such that for all $f \in h^p_{\omega^p}(\mathbb{R}^n)$,*

$$\left\|I_\alpha^{\mathrm{loc}}(f)\right\|_{L^q_{\omega^q}(\mathbb{R}^n)} \le C\|f\|_{h^p_{\omega^p}(\mathbb{R}^n)}.$$

Proof. Let $r \in (\frac{n}{n-\alpha}, \infty)$ and ω satisfy $\omega^{\frac{nr}{nr-n-r\alpha}} \in A_1^{\mathrm{loc}}(\mathbb{R}^n)$ and $\int_{\mathbb{R}^n} [\omega(x)]^p\,dx = \infty$. Then by

$$\omega^{\frac{nr}{nr-n-r\alpha}} \in A_1^{\mathrm{loc}}(\mathbb{R}^n)$$

and Lemma 2.3(ii), we know that there exists an $\eta_1 \in (0,\infty)$ such that

$$\omega^{\frac{nr(1+\eta_1)}{nr-n-r\alpha}} \in A_1^{\mathrm{loc}}(\mathbb{R}^n). \tag{8.17}$$

Let

$$\frac{1}{p_1} \equiv \frac{1}{r} + \frac{\alpha}{n} + \left(1 - \frac{1}{r} - \frac{\alpha}{n}\right)/(1+\eta_1) \text{ and } \frac{1}{q_1} \equiv \frac{1}{p_1} - \frac{\alpha}{n}. \tag{8.18}$$

Then by $r \in (\frac{n}{n-\alpha}, \infty)$, we know that

$$p_1 \in \left(1, \frac{n}{\alpha}\right),\ r > q_1 \text{ and } \omega^{-r'} \in A^{\mathrm{loc}}\left(\frac{q_1'}{r'}, \frac{p_1'}{r'}\right). \tag{8.19}$$

Furthermore, from (8.17), the fact that $p_1 < \frac{nr(1+\eta_1)}{nr-n-r\alpha}$ and Hölder's inequality, we infer that

$$\omega^{p_1} \in A_1^{\mathrm{loc}}(\mathbb{R}^n), \tag{8.20}$$

which together with Lemma 2.3(ii) implies that there exists an $\eta_2 \in (0,\infty)$ such that $\omega^{p_1(1+\eta_2)} \in A_1^{\mathrm{loc}}(\mathbb{R}^n)$. Let

$$\widetilde{q} \equiv p_1(1+\eta_2). \tag{8.21}$$

By $\frac{nr}{nr-n-r\alpha} > p$ and Hölder's inequality, we see that $\omega^p \in A_1^{\mathrm{loc}}(\mathbb{R}^n)$. Let $s \equiv \lfloor n(1/p-1) \rfloor$. To show Theorem 8.10, by the facts that $h^p_{\omega^p}(\mathbb{R}^n)$ and $L^q_{\omega^q}(\mathbb{R}^n)$ are respectively p-quasi-Banach space and 1-quasi-Banach space, and Theorem 6.4(i) with $\Phi(t) \equiv t^p$ for all $t \in (0,\infty)$, it suffices to show that for any $(p, \widetilde{q}, s)_{\omega^p}$-atom a supported in $Q_0 \equiv Q(x_0, R_0)$ with $R_0 \in (0,2]$,

$$\left\|I_\alpha^{\mathrm{loc}}(a)\right\|_{L^q_{\omega^q}(\mathbb{R}^n)} \lesssim 1. \tag{8.22}$$

By $\operatorname{supp}(a) \subset Q_0$ and the definition of $I_\alpha^{\mathrm{loc}}(a)$, we see that

$$\operatorname{supp}\left(I_\alpha^{\mathrm{loc}}(a)\right) \subset Q(x_0, R_0 + 4). \tag{8.23}$$

Now, we prove (8.22) by considering the following two cases for R_0.

Case 1) $R_0\in[1,2]$. In this case, from (8.23), Hölder's inequality, (8.19), Lemma 8.3, $R_0\in[1,2]$ and $\frac{1}{q}-\frac{1}{q_1}=\frac{1}{p}-\frac{1}{p_1}$, we deduce that

$$\begin{aligned}&\left\{\int_{\mathbb{R}^n}\left|I_\alpha^{\rm loc}(a)(x)\right|^q[\omega(x)]^q\,dx\right\}^{1/q}\\&\quad\lesssim\left\{\int_{Q(x_0,R_0+4)}\left|I_\alpha^{\rm loc}(a)(x)\right|^{q_1}[\omega(x)]^{q_1}\,dx\right\}^{1/q_1}|Q_0|^{\frac{1}{q}-\frac{1}{q_1}}\\&\quad\lesssim\|a\|_{L^{p_1}_{\omega^{p_1}}(\mathbb{R}^n)}|Q_0|^{\frac{1}{p}-\frac{1}{p_1}}.\end{aligned}\tag{8.24}$$

By (8.20) and the definition of $A_1^{\rm loc}(\mathbb{R}^n)$, we know that $\omega^{\frac{p_1(\widetilde{q}-p)}{(\widetilde{q}-p_1)}}\in A_1^{\rm loc}(\mathbb{R}^n)$. From this and Lemma 8.2(i), we infer that $\omega^p\in RH^{\rm loc}_{\frac{p_1(\widetilde{q}-p)}{p(\widetilde{q}-p_1)}}(\mathbb{R}^n)$, which implies that

$$\left\{\int_{Q_0}[\omega(x)]^p\,dx\right\}^{\frac{1}{\widetilde{q}}-\frac{1}{p}}\left\{\int_{Q_0}[\omega(x)]^{\frac{p_1(\widetilde{q}-p)}{(\widetilde{q}-p_1)}}\,dx\right\}^{\frac{1}{p_1}-\frac{1}{\widetilde{q}}}\lesssim|Q_0|^{\frac{1}{p_1}-\frac{1}{p}}.\tag{8.25}$$

This, combined with (8.24), Hölder's inequality and the fact that a is a $(p,\,\widetilde{q},\,s)_{\omega^p}$-atom, yields that

$$\begin{aligned}\left\|I_\alpha^{\rm loc}(a)\right\|_{L^q_{\omega^q}(\mathbb{R}^n)}&\lesssim\|a\|_{L^{p_1}_{\omega^{p_1}}(\mathbb{R}^n)}|Q_0|^{\frac{1}{p}-\frac{1}{p_1}}\\&\lesssim\left\{\int_{Q_0}|a(x)|^{\widetilde{q}}[\omega(x)]^p\,dx\right\}^{1/\widetilde{q}}\left\{\int_{Q_0}[\omega(x)]^{\frac{p_1(\widetilde{q}-p)}{(\widetilde{q}-p_1)}}\,dx\right\}^{\frac{1}{p_1}-\frac{1}{\widetilde{q}}}|Q_0|^{\frac{1}{p}-\frac{1}{p_1}}\\&\lesssim\left\{\int_{Q_0}[\omega(x)]^p\,dx\right\}^{\frac{1}{\widetilde{q}}-\frac{1}{p}}\left\{\int_{Q_0}[\omega(x)]^{\frac{p_1(\widetilde{q}-p)}{(\widetilde{q}-p_1)}}\,dx\right\}^{\frac{1}{p_1}-\frac{1}{\widetilde{q}}}|Q_0|^{\frac{1}{p}-\frac{1}{p_1}}\lesssim1.\end{aligned}$$

This shows (8.22) in Case 1).

Case 2) $R_0\in(0,1)$. In this case, let $\widetilde{Q}_0\equiv4nQ_0$. From (8.23), it follows that

$$\begin{aligned}\left\|I_\alpha^{\rm loc}(a)\right\|_{L^q_{\omega^q}(\mathbb{R}^n)}&\le\left\{\int_{\widetilde{Q}_0}\left|I_\alpha^{\rm loc}(a)(x)\right|^q[\omega(x)]^q\,dx\right\}^{1/q}\\&\quad+\left\{\int_{Q(x_0,R_0+4)\setminus\widetilde{Q}_0}\left|I_\alpha^{\rm loc}(a)(x)\right|^q[\omega(x)]^q\,dx\right\}^{1/q}\\&\equiv{\rm I}_1+{\rm I}_2.\end{aligned}\tag{8.26}$$

To estimate ${\rm I}_1$, by Hölder's inequality, (8.15) and (8.25), we conclude that

$$\begin{aligned}{\rm I}_1&\le\left(\int_{Q_1}\left|I_\alpha^{\rm loc}(a)(x)\right|^{q_1}[\omega(x)]^{q_1}\,dx\right)^{1/q_1}|Q_0|^{\frac{1}{p}-\frac{1}{p_1}}\\&\lesssim\|a\|_{L^{p_1}_{\omega^{p_1}}(\mathbb{R}^n)}|Q_0|^{\frac{1}{p}-\frac{1}{p_1}}\\&\lesssim\left\{\int_{Q_0}[\omega(x)]^p\,dx\right\}^{\frac{1}{\widetilde{q}}-\frac{1}{p}}\left\{\int_{Q_0}[\omega(x)]^{\frac{p_1(\widetilde{q}-p)}{(\widetilde{q}-p_1)}}\,dx\right\}^{\frac{1}{p_1}-\frac{1}{\widetilde{q}}}|Q_0|^{\frac{1}{p}-\frac{1}{p_1}}\lesssim1.\end{aligned}\tag{8.27}$$

To estimate ${\rm I}_2$, for any fixed $x\in Q(x_0,R_0+4)\setminus\widetilde{Q}_0$, let E^s be the Taylor expansion of $\frac{\phi_0(\cdot)}{|\cdot|^{n-\alpha}}$ about $x-x_0$ with degree s. Let m_0 be the integer such that

$$2^{m_0-1}nR_0 \le R_0+4 < 2^{m_0}nR_0.$$

Since $\omega^p \in A_1^{\rm loc}(\mathbb{R}^n) \subset A_{\widetilde{q}}^{\rm loc}(\mathbb{R}^n)$, by (2.1), we have

$$\left(\int_{Q_0}[\omega(x)]^p\,dx\right)^{\frac{1}{\widetilde{q}}-\frac{1}{p}}\left(\int_{Q_0}[\omega(x)]^{-\frac{p\widetilde{q}'}{\widetilde{q}}}\,dx\right)^{\frac{1}{\widetilde{q}'}} \lesssim \left(\int_{Q_0}[\omega(x)]^p\,dx\right)^{\frac{1}{p}}|Q_0|.$$

From this, the vanishing condition of a, Minkowski's inequality, Taylor's remainder theorem, the fact that

$$\sum_{\alpha\in\mathbb{Z}_+^n,\,|\alpha|=s+1}\left|\partial^\alpha\left(\frac{\phi_0(\cdot)}{|\cdot|^{n-\alpha}}\right)(z)\right| \lesssim \frac{1}{|z|^{n+s+1-\alpha}}$$

for all $z\in\mathbb{R}^n\setminus\{0\}$, and Hölder's inequality, we deduce that

$$\begin{aligned}
\mathrm{I}_2 &\le \left(\sum_{k=2}^{m_0}\int_{2^{k+1}nQ_0\setminus 2^knQ_0}\left\{\int_{Q_0}\left|\frac{\phi_0(x-y)}{|x-y|^{n-\alpha}}-E^s(x-y)\right|\right.\right.\\
&\qquad\left.\left.\times|a(y)|\,dy\right\}^q[\omega(x)]^q\,dx\right)^{1/q}\\
&\le \sum_{k=2}^{m_0}\int_{Q_0}|a(y)|\left\{\int_{2^{k+1}nQ_0\setminus 2^knQ_0}\left|\frac{\phi_0(x-y)}{|x-y|^{n-\alpha}}-E^s(x-y)\right|^q[\omega(x)]^q\,dx\right\}^{1/q}dy\\
&\lesssim \sum_{k=2}^{m_0}\int_{Q_0}|a(y)|\left\{\int_{2^{k+1}nQ_0\setminus 2^knQ_0}\left(\frac{|y-x_0|^{s+1}}{|\theta(x-y)-(1-\theta)(x-x_0)|^{n+s+1-\alpha}}\right)^q\right.\\
&\qquad\left.\times[\omega(x)]^q\,dx\right\}^{1/q}dy\\
&\lesssim \sum_{k=2}^{m_0}\int_{Q_0}|a(y)|\left\{\int_{2^{k+1}nQ_0\setminus 2^knQ_0}\left(\frac{|y-x_0|^{s+1}}{|x-x_0|^{n+s+1-\alpha}}\right)^q[\omega(x)]^q\,dx\right\}^{1/q}dy\\
&\lesssim \sum_{k=2}^{m_0}\frac{R_0^{\alpha-n}}{2^{k(n+s+1-\alpha)}}\left\{\int_{Q_0}|a(y)|\,dy\right\}\left\{\int_{2^{k+1}nQ_0}[\omega(x)]^q\,dx\right\}^{1/q}\\
&\lesssim \sum_{k=2}^{m_0}\frac{R_0^{\alpha-n}}{2^{k(n+s+1-\alpha)}}\left\{\int_{Q_0}|a(y)|^{\widetilde{q}}[\omega(y)]^p\,dy\right\}^{1/\widetilde{q}}\\
&\qquad\times\left\{\int_{Q_0}[\omega(y)]^{-p\widetilde{q}'/\widetilde{q}}\,dy\right\}^{1/\widetilde{q}'}\left\{\int_{2^{k+1}nQ_0}[\omega(x)]^q\,dx\right\}^{1/q}\\
&\lesssim \sum_{k=2}^{m_0}\frac{R_0^{\alpha}}{2^{k(n+s+1-\alpha)}}\left\{\int_{Q_0}[\omega(x)]^p\,dx\right\}^{-1/p}\left\{\int_{2^{k+1}nQ_0}[\omega(x)]^q\,dx\right\}^{1/q},
\end{aligned}\tag{8.28}$$

where $\theta\in(0,1)$ and in the fourth inequality we used the fact that for any $y\in Q_0$ and $x\in 2^{k+1}nQ_0\setminus 2^knQ_0$ with $k\in\{2,\cdots,m_0\}$, $|(x-x_0)-\theta(y-x_0)|\gtrsim|x-x_0|$.

From $\frac{nr(1+\eta_1)}{nr-n-r\alpha}=\frac{rq_1}{r-q_1}>\frac{rq}{r-q}$, (8.17) and Hölder's inequality, it follows that

$$\omega^{\frac{rq}{r-q}}\in A_1^{\mathrm{loc}}(\mathbb{R}^n). \tag{8.29}$$

By Lemma 2.3(i) and Remark 2.4 with $\widetilde{C}\equiv 20n$, we know that there exists a function $\widetilde{\omega}$ on $\mathbb{R}^n$ such that $\widetilde{\omega}^{\frac{rq}{r-q}}\in A_1(\mathbb{R}^n)$ such that $\widetilde{\omega}=\omega$ on $Q(x_0,20n)$, which together with $2^{m_0+1}nQ_0\subset Q(x_0,20n)$ and Lemma 2.3(viii) implies that for any $k\in\{1,\,2,\,\cdots,\,m_0\}$,

$$\int_{2^knQ_0}[\omega(x)]^{\frac{rq}{r-q}}\,dx=\int_{2^knQ_0}[\widetilde{\omega}(x)]^{\frac{rq}{r-q}}\,dx\lesssim 2^{kn}\int_{Q_0}[\widetilde{\omega}(x)]^{\frac{rq}{r-q}}\,dx\lesssim 2^{kn}\int_{Q_0}[\omega(x)]^{\frac{rq}{r-q}}\,dx.$$

By this estimate and Hölder's inequality, we have

$$\left\{\int_{2^{k+1}nQ_0}[\omega(x)]^q\,dx\right\}^{1/q}\lesssim R_0^{n/r}2^{\frac{kn}{q}}\left\{\int_{Q_0}[\omega(x)]^{\frac{rq}{r-q}}\,dx\right\}^{\frac{1}{q}-\frac{1}{r}}. \tag{8.30}$$

Moreover, by (8.29) and Lemma 8.8(i), we know that $\omega^p\in RH^{\mathrm{loc}}_{\frac{rq}{p(r-q)}}(\mathbb{R}^n)$. Thus, we have

$$\left\{\int_{Q_0}[\omega(x)]^p\,dx\right\}^{-1/p}\left\{\int_{Q_0}[\omega(x)]^{\frac{rq}{r-q}}\,dx\right\}^{\frac{1}{q}-\frac{1}{r}}\lesssim R_0^{-\frac{n}{r}-\alpha},$$

which together with (8.28) and (8.30) implies that

$$\mathrm{I}_2\lesssim\sum_{k=2}^{m_0}2^{-k(n+s+1-\alpha-n/q)}.$$

From $\frac{1}{q}=\frac{1}{p}-\frac{\alpha}{n}$ and $r>\frac{n}{n-\alpha}$, we deduce that $n+s+1-\alpha-n/q>n+s+1-n/p$, which together with $s=\lfloor n(1/p-1)\rfloor$ implies that $n+s+1-\alpha-n/q>0$. Thus,

$$\mathrm{I}_2\lesssim\sum_{k=2}^{m_0}2^{-k(n+s+1-\alpha-n/q)}\lesssim 1.$$

This combined with (8.26) and (8.27) proves (8.22) in Case 2), which completes the proof of Theorem 8.10. ■

THEOREM 8.11. *Let* $\alpha\in(0,1)$, $p\in(0,\frac{n}{n+\alpha}]$ *and* $\frac{1}{q}=\frac{1}{p}-\frac{\alpha}{n}$. *For some* $r\in(\frac{n}{n-\alpha},\infty)$, *if the weight* ω *satisfies* $\omega^{\frac{nr}{nr-n-r\alpha}}\in A_1^{\mathrm{loc}}(\mathbb{R}^n)$ *and* $\int_{\mathbb{R}^n}[\omega(x)]^p\,dx=\infty$, *then there exists a positive constant* C *such that for all* $f\in h^p_{\omega^p}(\mathbb{R}^n)$,

$$\left\|I_\alpha^{\mathrm{loc}}(f)\right\|_{h^q_{\omega^q}(\mathbb{R}^n)}\le C\|f\|_{h^p_{\omega^p}(\mathbb{R}^n)}.$$

Proof. Let p_1, q_1 and $\widetilde{q}$ be respectively as in (8.18) and (8.21). To show Theorem 8.11, by the facts that $h^p_{\omega^p}(\mathbb{R}^n)$ and $h^q_{\omega^q}(\mathbb{R}^n)$ are respectively the p-quasi-Banach space and the q-quasi-Banach space, Theorem 6.4(i) with $\Phi(t)\equiv t^p$ for all $t\in(0,\infty)$ and Theorem 3.14, it suffices to show that for any $(p,\,\widetilde{q},\,s)_{\omega^p}$-atom a supported in $Q_0\equiv Q(x_0,R_0)$ with $R_0\in(0,2]$,

$$\left\|\mathcal{G}_N^0\left(I_\alpha^{\mathrm{loc}}(a)\right)\right\|_{L^\Phi_{\omega^q}(\mathbb{R}^n)}\lesssim 1. \tag{8.31}$$

By $\mathrm{supp}\,(a)\subset Q_0$ and definitions of $I_\alpha^{\mathrm{loc}}(a)$ and $\mathcal{G}_N^0(I_\alpha^{\mathrm{loc}}(a))$, we know that

$$\mathrm{supp}\,\left(\mathcal{G}_N^0(I_\alpha^{\mathrm{loc}}(a))\right)\subset Q(x_0,R_0+8). \tag{8.32}$$

Now, we prove (8.31) by considering the following two cases for R_0.

Case 1) $R_0 \in [1,2]$. In this case, by (8.17), the fact that $\frac{nr(1+\eta_1)}{nr-n-r\alpha} > q_1$ and Hölder's inequality, we know that $\omega^{q_1} \in A_1^{\mathrm{loc}}(\mathbb{R}^n)$, where η_1 is as in (8.17). From this, (8.32), Hölder's inequality, Proposition 3.2(ii), Lemma 8.9 and (8.25), it follows that

$$\begin{aligned}
&\int_{\mathbb{R}^n} \left|\mathcal{G}_N^0\left(I_\alpha^{\mathrm{loc}}(a)\right)(x)\right|^q [\omega(x)]^q\,dx\\
&\quad\lesssim |Q_0|^{1-\frac{q}{q_1}}\left\{\int_{Q(x_0,R_0+8)} \left|\mathcal{G}_N^0\left(I_\alpha^{\mathrm{loc}}(a)\right)(x)\right|^{q_1}[\omega(x)]^{q_1}\,dx\right\}^{q/q_1}\\
&\quad\lesssim |Q_0|^{1-\frac{q}{q_1}}\left\{\int_{Q(x_0,R_0+8)} \left|I_\alpha^{\mathrm{loc}}(a)(x)\right|^{q_1}[\omega(x)]^{q_1}\,dx\right\}^{q/q_1}\\
&\quad\lesssim |Q_0|^{1-\frac{q}{q_1}}\left\{\int_{Q_0} |a(x)|^{p_1}[\omega(x)]^{p_1}\,dx\right\}^{q/p_1}\\
&\quad\lesssim |Q_0|^{1-\frac{q}{q_1}}\left\{\int_{Q_0} |a(x)|^{\widetilde{q}}[\omega(x)]^{p}\,dx\right\}^{q/\widetilde{q}}\left\{\int_{Q_0}[\omega(x)]^{\frac{p_1(\widetilde{q}-p)}{\widetilde{q}-p_1}}\,dx\right\}^{(\frac{1}{p_1}-\frac{1}{\widetilde{q}})q}\\
&\quad\lesssim \left\{|Q_0|^{\frac{1}{q}-\frac{1}{q_1}}\left(\int_{Q_0}[\omega(x)]^p\,dx\right)^{\frac{1}{\widetilde{q}}-\frac{1}{p}}\left(\int_{Q_0}[\omega(x)]^{\frac{p_1(\widetilde{q}-p)}{\widetilde{q}-p_1}}\,dx\right)^{\frac{1}{p_1}-\frac{1}{\widetilde{q}}}\right\}^q \lesssim 1,
\end{aligned}$$

which proves (8.31) in Case 1).

Case 2) $R_0 \in (0,1)$. In this case, let $\widetilde{Q}_0 \equiv 8nQ_0$. Then from (8.32), we conclude that

$$\begin{aligned}
\int_{\mathbb{R}^n}\left[\mathcal{G}_N^0\left(I_\alpha^{\mathrm{loc}}a\right)(x)\right]^q[\omega(x)]^q\,dx &= \int_{\widetilde{Q}_0}\left[\mathcal{G}_N^0\left(I_\alpha^{\mathrm{loc}}a\right)(x)\right]^q[\omega(x)]^q\,dx\\
&\quad+\int_{Q(x_0,R_0+8)\setminus\widetilde{Q}_0}\cdots \equiv \mathrm{I}_1+\mathrm{I}_2. \qquad (8.33)
\end{aligned}$$

For I_1, similarly to the proof of Case 1), we have

$$\begin{aligned}
\mathrm{I}_1 &\lesssim |Q_0|^{1-\frac{q}{q_1}}\left\{\int_{\widetilde{Q}_0}\left[\mathcal{G}_N^0\left(I_\alpha^{\mathrm{loc}}(a)\right)(x)\right]^{q_1}[\omega(x)]^{q_1}\,dx\right\}^{q/q_1}\\
&\lesssim |Q_0|^{1-\frac{q}{q_1}}\left\{\int_{\mathbb{R}^n}\left|I_\alpha^{\mathrm{loc}}(a)(x)\right|^{q_1}[\omega(x)]^{q_1}\,dx\right\}^{q/q_1}\\
&\lesssim |Q_0|^{1-\frac{q}{q_1}}\left\{\int_{Q_0}|a(x)|^{p_1}[\omega(x)]^{p_1}\,dx\right\}^{q/p_1}\\
&\lesssim |Q_0|^{1-\frac{q}{q_1}}\left\{\int_{Q_0}|a(x)|^{\widetilde{q}}[\omega(x)]^{p}\,dx\right\}^{q/\widetilde{q}}\left\{\int_{Q_0}[\omega(x)]^{\frac{p_1(\widetilde{q}-p)}{\widetilde{q}-p_1}}\,dx\right\}^{(\frac{1}{p_1}-\frac{1}{\widetilde{q}})q}\\
&\lesssim \left\{|Q_0|^{\frac{1}{q}-\frac{1}{q_1}}\left(\int_{Q_0}[\omega(x)]^p\,dx\right)^{\frac{1}{\widetilde{q}}-\frac{1}{p}}\left(\int_{Q_0}[\omega(x)]^{\frac{p_1(\widetilde{q}-p)}{\widetilde{q}-p_1}}\,dx\right)^{\frac{1}{p_1}-\frac{1}{\widetilde{q}}}\right\}^q \lesssim 1. \qquad (8.34)
\end{aligned}$$

To estimate I_2, let $x \in (\widetilde{Q}_0)^{\complement}$, $\psi \in \mathcal{D}_N^0(\mathbb{R}^n)$, $t \in (0,1)$ and P_ψ^s be the Taylor expansion of ψ about $(x-x_0)/t$ with degree s, where $s \equiv \lfloor n(1/p-1)\rfloor$. Then we have

$$\left|I_\alpha^{\mathrm{loc}}(a)*\psi_t(x)\right| = \frac{1}{t^n}\left|\int_{\mathbb{R}^n} I_\alpha^{\mathrm{loc}}(a)(y)\psi\left(\frac{x-y}{t}\right)\,dy\right|$$

$$\begin{aligned}
&= \frac{1}{t^n}\left|\int_{\mathbb{R}^n} I_\alpha^{\rm loc}(a)(y)\left[\psi\left(\frac{x-y}{t}\right)-P_\psi^s\left(\frac{x-y}{t}\right)\right]dy\right|\\
&\le \frac{1}{t^n}\int_{2\sqrt{n}Q_0}\left|I_\alpha^{\rm loc}(a)(y)\right|\left|\psi\left(\frac{x-y}{t}\right)-P_\psi^s\left(\frac{x-y}{t}\right)\right|dy\\
&\quad+\frac{1}{t^n}\int_{Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})\setminus 2\sqrt{n}Q_0}\cdots+\frac{1}{t^n}\int_{[Q(x_0,\frac{|x-x_0|}{2\sqrt{n}})]^\complement}\cdots\\
&\equiv \mathrm{E}_1+\mathrm{E}_2+\mathrm{E}_3. \qquad (8.35)
\end{aligned}$$

To estimate E_1, by $x\in(\widetilde{Q}_0)^\complement$ and $t\in(0,1)$, we see that $\mathrm{E}_1\neq 0$ implies that $t>\frac{|x-x_0|}{2}$. From

$$\omega^{\frac{nr}{nr-n-r\alpha}}\in A_1^{\rm loc}(\mathbb{R}^n)$$

and the definition of $A_1^{\rm loc}(\mathbb{R}^n)$, it follows that $\omega\in A_1^{\rm loc}(\mathbb{R}^n)$. Let $q_2\equiv\frac{2q_1-1}{q_1}$. Then by $\omega\in A_1^{\rm loc}(\mathbb{R}^n)\subset A_{q_2}^{\rm loc}(\mathbb{R}^n)$ and Lemma 2.3(iv), we know that $\omega^{-q_1'}=\omega^{1-q_2'}\in A_{q_2'}^{\rm loc}(\mathbb{R}^n)$. From these facts, Taylor's remainder theorem, Hölder's inequality, Lemmas 8.9 and 2.3(v), Remark 2.4 with $\widetilde{C}=2\sqrt{n}$, (8.25) and (2.2), we infer that

$$\begin{aligned}
\mathrm{E}_1&\lesssim\frac{1}{t^{n+s+1}}\left\|I_\alpha^{\rm loc}(a)\right\|_{L^{q_1}_{\omega^{q_1}}(\mathbb{R}^n)}\left\{\sum_{\alpha\in\mathbb{Z}_+^n,\,|\alpha|=s+1}\int_{2\sqrt{n}Q_0}\left|\partial^\alpha\psi\left(\frac{\xi}{t}\right)\right|^{q_1'}\right.\\
&\qquad\left.\times|y-x_0|^{(s+1)q_1'}[\omega(y)]^{-q_1'}\,dy\right\}^{1/q_1'}\\
&\lesssim\frac{R_0^{s+1}}{|x-x_0|^{n+s+1}}\|a\|_{L^{p_1}_{\omega^{p_1}}(\mathbb{R}^n)}\left\{\int_{2\sqrt{n}Q_0}[\omega(y)]^{-q_1'}\,dy\right\}^{1/q_1'}\\
&\lesssim\frac{R_0^{s+1}}{|x-x_0|^{n+s+1}}\left\{\int_{Q_0}|a(x)|^{\widetilde{q}}[\omega(x)]^p\,dx\right\}^{1/\widetilde{q}}\\
&\qquad\times\left\{\int_{Q_0}[\omega(x)]^{\frac{p_1(\widetilde{q}-p)}{\widetilde{q}-p_1}}\,dx\right\}^{\frac{1}{p_1}-\frac{1}{\widetilde{q}}}\left\{\int_{Q_0}[\omega(y)]^{-q_1'}\,dy\right\}^{1/q_1'}\\
&\lesssim\frac{R_0^{s+1}}{|x-x_0|^{n+s+1}}|Q_0|^{\frac{1}{q_1}-\frac{1}{q}}\left\{\int_{Q_0}[\omega(y)]^{-q_1'}\,dy\right\}^{1/q_1'}\\
&\lesssim\frac{R_0^{s+1}}{|x-x_0|^{n+s+1}}|Q_0|^{1-\frac{1}{q}}\left[\operatorname*{essinf}_{z\in Q_0}\omega(z)\right]^{-1}, \qquad (8.36)
\end{aligned}$$

where $\gamma\in(0,1)$ and $\xi\equiv\gamma(x-y)+(1-\gamma)(x-x_0)$. Similarly to the estimates of G_2 and G_3 in the proof of Theorem 8.2, we have

$$\max\{\mathrm{E}_2,\mathrm{E}_3\}\lesssim\frac{R_0^{n+s+1}}{|x-x_0|^{n+s+1-\alpha}}|Q_0|^{-1/p}\left[\operatorname*{essinf}_{z\in Q_0}\omega(z)\right]^{-1}. \qquad (8.37)$$

Thus, from (8.35), (8.36), (8.37) and the facts that $|x-x_0|\ge 4nR_0$ and $\frac{1}{q}=\frac{1}{p}-\frac{\alpha}{n}$, we deduce that

$$\left|\left[I_\alpha^{\rm loc}(a)\right]*\psi_t(x)\right|\lesssim\frac{R_0^{s+1}}{|x-x_0|^{n+s+1}}|Q_0|^{1-\frac{1}{q}}\left[\operatorname*{essinf}_{z\in Q_0}\omega(z)\right]^{-1}$$

$$
\begin{aligned}
&+\frac{R_0^{n+s+1}}{|x-x_0|^{n+s+1-\alpha}}|Q_0|^{-1/p}\left[\operatorname*{essinf}_{z\in Q_0}\omega(z)\right]^{-1}\\
&\lesssim \frac{R_0^{n+s+1}}{|x-x_0|^{n+s+1-\alpha}}|Q_0|^{-1/p}\left[\operatorname*{essinf}_{z\in Q_0}\omega(z)\right]^{-1},
\end{aligned}
$$

which together with the arbitrariness of $\psi\in\mathcal{D}_N^0(\mathbb{R}^n)$ implies that for any $x\in(\widetilde{Q}_0)^{\complement}$,

$$
\mathcal{G}_N^0\left(I_\alpha^{\mathrm{loc}}(a)\right)(x)\lesssim \frac{R_0^{n+s+1}}{|x-x_0|^{n+s+1-\alpha}}|Q_0|^{-1/p}\left[\operatorname*{essinf}_{z\in Q_0}\omega(z)\right]^{-1}. \tag{8.38}
$$

By $s=\lfloor n(1/p-1)\rfloor$ and $\frac{1}{q}=\frac{1}{p}-\frac{\alpha}{n}$, we know that $(n+s+1-\alpha)q-n>0$. Let m_0 be the integer such that $2^{m_0}nR_0\le R_0+8<2^{m_0+1}nR_0$. By

$$
\omega^{\frac{nr}{nr-n-r\alpha}}\in A_1^{\mathrm{loc}}(\mathbb{R}^n)
$$

and the definition of $A_1^{\mathrm{loc}}(\mathbb{R}^n)$, we see that $\omega^q\in A_1^{\mathrm{loc}}(\mathbb{R}^n)$, which together Lemma 2.3(i) implies that there exists a function $\widetilde{\omega}$ on $\mathbb{R}^n$ such that $\widetilde{\omega}=\omega$ on $Q(x_0,R_0+8)$ and $\widetilde{\omega}^q\in A_1(\mathbb{R}^n)$. From this, $\frac{1}{q}=\frac{1}{p}-\frac{\alpha}{n}$, (8.38), (i) and (viii) of Lemma 2.3, the definition of $A_1^{\mathrm{loc}}(\mathbb{R}^n)$ and $(n+s+1-\alpha)q-n>0$, we infer that

$$
\begin{aligned}
\mathrm{I}_2&\lesssim\int_{Q(x_0,R_0+8)\setminus\widetilde{Q}_0}\left\{\frac{R_0^{n+s+1}}{|x-x_0|^{n+s+1-\alpha}}|Q_0|^{-1/p}\left[\operatorname*{essinf}_{z\in Q_0}\omega(z)\right]^{-1}\right\}^q[\widetilde{\omega}(x)]^q\,dx\\
&\lesssim\sum_{k=3}^{m_0}\int_{2^{k+1}nQ_0\setminus 2^knQ_0}\left\{\frac{R_0^{n+s+1-n/q-\alpha}}{(2^kR_0)^{n+s+1-\alpha}}\left[\operatorname*{essinf}_{z\in Q_0}\omega(z)\right]^{-1}\right\}^q[\widetilde{\omega}(x)]^q\,dx\\
&\lesssim\sum_{k=3}^{m_0}\frac{R_0^{-n}}{2^{k[(n+s+1-\alpha)q-n]}}\left[\operatorname*{essinf}_{z\in Q_0}\omega(z)\right]^{-q}\int_{Q_0}[\omega(x)]^q\,dx\\
&\lesssim\sum_{k=3}^{m_0}\frac{R_0^{-n}}{2^{k[(n+s+1-\alpha)q-n]}}|Q_0|\left[\operatorname*{essinf}_{z\in Q_0}\omega(z)\right]^{-q}\operatorname*{essinf}_{z\in Q_0}[\omega(z)]^q\\
&\lesssim\sum_{k=3}^{m_0}2^{-k[(n+s+1-\alpha)q-n]}\lesssim 1,
\end{aligned}
$$

which together with (8.33) and (8.34) implies (8.31) in Case 2). This finishes the proof of Theorem 8.11. ■

The pseudo-differential operators have been extensively studied in the literature, and they are important in the study of partial differential equations and harmonic analysis; see, for example, [46, 51, 45, 50]. Now, we recall the notion of the pseudo-differential operators of order zero.

Definition 8.12. Let δ be a real number. A *symbol* in $S_{1,\delta}^0(\mathbb{R}^n)$ is a smooth function $\sigma(x,\xi)$ defined on $\mathbb{R}^n\times\mathbb{R}^n$ such that for all multi-indices α and β, the following estimate holds:

$$
\left|\partial_x^\alpha\partial_\xi^\beta\sigma(x,\xi)\right|\le C(\alpha,\,\beta)(1+|\xi|)^{-|\beta|+\delta|\alpha|},
$$

where $C(\alpha,\,\beta)$ is a positive constant independent of x and ξ. Let f be a Schwartz function and $\hat{f}$ denote its Fourier transform. The operator T given by setting, for all $x \in \mathbb{R}^n$,

$$Tf(x) \equiv \int_{\mathbb{R}^n} \sigma(x,\xi) e^{2\pi i x\xi} \hat{f}(\xi)\,d\xi$$

is called a *pseudo-differential operator* with symbol $\sigma(x,\xi) \in S^0_{1,\delta}(\mathbb{R}^n)$.

In the rest of this section, let

$$\phi(t) \equiv (1+t)^\alpha \tag{8.39}$$

for all $\alpha \in (0,\infty)$ and $t \in (0,\infty)$. Recall that a *weight* always means a locally integrable function which is positive almost everywhere. The following notion of the weight class $A_p(\phi)$ was introduced by Tang [50].

Definition 8.13. A weight ω is said to belong to the *class* $A_p(\phi)$ for $p \in (1,\infty)$, if there exists a positive constant C such that for all cubes $Q \equiv Q(x,r)$,

$$\left(\frac{1}{\phi(|Q|)|Q|}\int_Q \omega(y)\,dy\right)\left(\frac{1}{\phi(|Q|)|Q|}\int_Q [\omega(y)]^{-\frac{1}{p-1}}\,dy\right)^{p-1} \le C.$$

A weight ω is said to belong to the *class* $A_1(\phi)$, if there exists a positive constant C such that for all cubes $Q \subset \mathbb{R}^n$ and almost every $x \in \mathbb{R}^n$, $M_\phi(\omega)(x) \le C\omega(x)$, where for all $x \in \mathbb{R}^n$,

$$M_\phi(\omega)(x) \equiv \sup_{Q \ni x} \frac{1}{\phi(|Q|)|Q|}\int_Q |f(y)|\,dy,$$

and the supremum is taken over all cubes $Q \subset \mathbb{R}^n$ and $Q \ni x$.

Remark 8.14. By the definition of $A_p(\phi)$, we see that $A_p(\phi) \subset A_p^{\rm loc}(\mathbb{R}^n)$, and that $\phi(t) \ge 1$ for all $t \in (0,\infty)$ implies that $A_p(\mathbb{R}^n) \subset A_p(\phi)$ for all $p \in [1,\infty)$. Moreover, if $\omega \in A_p(\phi)$, $\omega(x)\,dx$ may not be a doubling measure; see the remark of Section 7 in [49] for the details.

Similarly to the classical Muckenhoupt weights, we recall some properties of weights $\omega \in A_\infty(\phi) \equiv \cup_{1 \le p < \infty} A_p(\phi)$. The following Lemmas 8.15 and 8.16 are respectively Lemmas 7.3 and 7.4 in [49].

Lemma 8.15. (i) *If $1 \le p_1 < p_2 < \infty$, then $A_{p_1}(\phi) \subset A_{p_2}(\phi)$.*

(ii) *For $p \in (1,\infty)$, $\omega \in A_p(\phi)$ if and only if $\omega^{-\frac{1}{p-1}} \in A_{p'}(\phi)$, where $1/p + 1/p' = 1$.*

(iii) *If $\omega \in A_p(\phi)$ for $p \in [1,\infty)$, then there exists a positive constant C such that for any cube $Q \subset \mathbb{R}^n$ and measurable set $E \subset Q$,*

$$\frac{|E|}{\phi(|Q|)|Q|} \le C\left(\frac{\omega(E)}{\omega(Q)}\right)^{1/p}.$$

Lemma 8.16. *Let T be an $S^0_{1,0}(\mathbb{R}^n)$ pseudo-differential operator. Then for $\omega \in A_p(\phi)$ with $p \in (1,\infty)$, there exists a positive constant $C(p,\omega)$ such that for all $f \in L^p_\omega(\mathbb{R}^n)$,*

$$\|Tf\|_{L^p_\omega(\mathbb{R}^n)} \le C(p,\omega)\|f\|_{L^p_\omega(\mathbb{R}^n)}.$$

The following Lemma 8.17 is just [18, Lemma 6].

Lemma 8.17. *Let T be an $S^0_{1,0}(\mathbb{R}^n)$ pseudo-differential operator. If $\psi \in \mathcal{D}(\mathbb{R}^n)$, then $T_t f = \psi_t * Tf$ has a symbol σ_t which satisfies that*

$$\left|\partial_x^\beta \partial_\xi^\alpha \sigma(x,\xi)\right| \le C(\alpha,\,\beta)(1+|\xi|)^{-|\alpha|},$$

and a kernel $K_t(x,\,z)$ which satisfies that

$$\left|\partial_x^\beta \partial_z^\alpha K_t(x,z)\right| \le C(\alpha,\,\beta)|z|^{-n-|\alpha|},$$

where $C(\alpha,\,\beta)$ is independent of t when $t \in (0,1)$.

Now, we establish the boundedness on $h^\Phi_\omega(\mathbb{R}^n)$ of the $S^0_{1,0}(\mathbb{R}^n)$ pseudo-differential operators as follows.

Theorem 8.18. *Let T be an $S^0_{1,0}(\mathbb{R}^n)$ pseudo-differential operator, Φ satisfy Assumption* (A), *$\omega \in A_\infty(\phi)$ and p_Φ be as in* (2.6). *If $p_\Phi = p_\Phi^+$ and Φ is of upper type p_Φ^+, then there exists a positive constant $C(\Phi,\,\omega)$, depending only on Φ, q_ω and the weight constant of $A_\infty(\phi)$, such that for all $f \in h^\Phi_\omega(\mathbb{R}^n)$,*

$$\|Tf\|_{h^\Phi_\omega(\mathbb{R}^n)} \le C(\Phi,\,\omega)\|f\|_{h^\Phi_\omega(\mathbb{R}^n)}.$$

Proof. Since $\omega \in A_\infty(\phi)$, then $\omega \in A_q(\phi)$ for some $q \in (1,\infty)$. To prove Theorem 8.18, by the fact that $\omega \in A_\infty(\phi) \subset A^{\rm loc}_\infty(\mathbb{R}^n)$, Theorem 6.4(i) and Theorem 3.14, it suffices to show that for all $(\rho,\,q)_\omega$-single-atoms and $(\rho,\,q,\,s)_\omega$-atoms a supported in $Q_0 \equiv Q(x_0,\,R_0)$ with $R_0 \in (0,2]$,

$$\left\|\mathcal{G}^0_N\,(Ta)\right\|_{L^\Phi_\omega(\mathbb{R}^n)} \lesssim 1. \tag{8.40}$$

By Theorem 5.6, we may assume that s satisfies $(n+s+1)p_\Phi > nq_\omega(1+\alpha)$, where p_Φ, q_ω and α are respectively as in (2.6), (2.4) and (8.39).

First, we prove (8.40) for any $(\rho,\,q)_\omega$-single-atom $a \ne 0$. In this case, $\omega(\mathbb{R}^n) < \infty$. Since Φ is concave, by Jensen's inequality, Hölder's inequality, Proposition 3.2(ii) and Lemma 8.16 and (2.8) with $t \equiv \omega(\mathbb{R}^n)$, we have

$$\begin{aligned}
&\int_{\mathbb{R}^n} \Phi\left(\mathcal{G}^0_N(Ta)(x)\right)\omega(x)\,dx\\
&\quad\le \omega(\mathbb{R}^n)\Phi\left(\frac{1}{\omega(\mathbb{R}^n)}\int_{\mathbb{R}^n} \mathcal{G}^0_N(Ta)(x)\omega(x)\,dx\right)\\
&\quad\le \omega(\mathbb{R}^n)\Phi\left(\frac{1}{[\omega(\mathbb{R}^n)]^{1/q}}\left\{\int_{\mathbb{R}^n}\left[\mathcal{G}^0_N(Ta)(x)\right]^q\omega(x)\,dx\right\}^{1/q}\right)\\
&\quad\lesssim \omega(\mathbb{R}^n)\Phi\left(\frac{1}{[\omega(\mathbb{R}^n)]^{1/q}}\|Ta\|_{L^q_\omega(\mathbb{R}^n)}\right) \lesssim \omega(\mathbb{R}^n)\Phi\left(\frac{1}{[\omega(\mathbb{R}^n)]^{1/q}}\|a\|_{L^q_\omega(\mathbb{R}^n)}\right)\\
&\quad\lesssim \omega(\mathbb{R}^n)\Phi\left(\frac{1}{\omega(\mathbb{R}^n)\rho(\omega(\mathbb{R}^n))}\right) \sim 1,
\end{aligned}$$

which shows (8.40) in this case.

Now, let a be any $(\rho,\,q,\,s)_\omega$-atom supported in $Q_0 \equiv Q(x_0,\,R_0)$ with $R_0 \in (0,2]$. Let $\widetilde{Q}_0 \equiv 2Q_0$. Then from Jensen's inequality, Hölder's inequality, Proposition 3.2(ii), Lemma 8.16, Lemma 8.15(iii) and (2.8) with $t \equiv \omega(\widetilde{Q}_0)$, it follows that

$$\int_{\widetilde{Q}_0} \Phi\left(\mathcal{G}^0_N(Ta)(x)\right)\omega(x)\,dx$$

$$
\begin{aligned}
&\le \omega(\widetilde{Q}_0)\Phi\left(\frac{1}{\omega(\widetilde{Q}_0)}\int_{\widetilde{Q}_0}\mathcal{G}_N^0(Ta)(x)\omega(x)\,dx\right)\\
&\le \omega(\widetilde{Q}_0)\Phi\left(\frac{1}{[\omega(\widetilde{Q}_0)]^{1/q}}\left\{\int_{\widetilde{Q}_0}\left[\mathcal{G}_N^0(Ta)(x)\right]^q\omega(x)\,dx\right\}^{1/q}\right)\\
&\lesssim \omega(\widetilde{Q}_0)\Phi\left(\frac{1}{[\omega(\widetilde{Q}_0)]^{1/q}}\|Ta\|_{L^q_\omega(\mathbb{R}^n)}\right)\lesssim \omega(\widetilde{Q}_0)\Phi\left(\frac{1}{[\omega(\widetilde{Q}_0)]^{1/q}}\|a\|_{L^q_\omega(\mathbb{R}^n)}\right)\\
&\lesssim \omega(\widetilde{Q}_0)\Phi\left(\frac{1}{\omega(\widetilde{Q}_0)\rho(\omega(\widetilde{Q}_0))}\right)\sim 1. \qquad (8.41)
\end{aligned}
$$

For any $\psi\in\mathcal{D}_N^0(\mathbb{R}^n)$ and $t\in(0,1)$, let $K_t(x,x-z)$ be the kernel of $T_ta(x)\equiv\psi_t*Ta(x)$. To estimate $\int_{\mathbb{R}^n\setminus\widetilde{Q}_0}\Phi[\mathcal{G}_N^0(Ta)(x)]\omega(x)\,dx$, we consider the following two cases for R_0.

Case 1) $R_0\in(0,1)$. In this case, we expand $K_t(x,x-z)$ into a Taylor series about $z=x_0$ such that for any $x\in(\mathbb{R}^n\setminus\widetilde{Q}_0)$,

$$
\psi_t*Ta(x)=\int_{\mathbb{R}^n}K_t(x,x-z)a(z)\,dz=\int_{Q_0}\sum_{\substack{\alpha\in\mathbb{Z}_+^n,\\ |\alpha|=s+1}}(\partial_z^\alpha K_t)(x,x-\xi)(z-x_0)^\alpha a(z)\,dz,
$$

where $\xi\equiv\theta z+(1-\theta)x_0$ for some $\theta\in(0,1)$. By z, $x_0\in Q_0$, we know that $\xi\in Q_0$. Thus, for any $x\in(\mathbb{R}^n\setminus\widetilde{Q}_0)$, $|x-\xi|\sim|x-x_0|$. From the above facts and Lemma 8.17, we deduce that

$$
|\psi_t*Ta(x)|\lesssim|x-\xi|^{-(n+s+1)}R_0^{s+1}\|a\|_{L^1(\mathbb{R}^n)}\lesssim|x-x_0|^{-(n+s+1)}|Q_0|^{\frac{s+1}{n}}\|a\|_{L^1(\mathbb{R}^n)},
$$

which together with the arbitrariness of $\psi\in\mathcal{D}_N^0(\mathbb{R}^n)$ implies that for all $x\in(\mathbb{R}^n\setminus\widetilde{Q}_0)$,

$$
\mathcal{G}_N^0(Ta)(x)\lesssim|x-x_0|^{-(n+s+1)}|Q_0|^{\frac{s+1}{n}}\|a\|_{L^1(\mathbb{R}^n)}.
$$

This, combined with Hölder's inequality, Lemma 8.15(iii) and the definition of $A_p(\phi)$, yields that

$$
\begin{aligned}
&\int_{\mathbb{R}^n\setminus\widetilde{Q}_0}\Phi\left(\mathcal{G}_N^0(Ta)(x)\right)\omega(x)\,dx\\
&\quad\le C\int_{\mathbb{R}^n\setminus\widetilde{Q}_0}\Phi\left(|x-x_0|^{-(n+s+1)}|Q_0|^{\frac{s+1}{n}}\|a\|_{L^1(\mathbb{R}^n)}\right)\omega(x)\,dx\\
&\quad\le C\int_{\mathbb{R}^n\setminus\widetilde{Q}_0}\Phi\Big(|Q_0|^{\frac{s+1}{n}}|x-x_0|^{-(n+s+1)}\|a\|_{L^q_\omega(\mathbb{R}^n)}\\
&\qquad\times\phi(|Q_0|)|Q_0|[\omega(Q_0)]^{-1/q}\Big)\omega(x)\,dx\\
&\quad\le C\int_{\mathbb{R}^n\setminus\widetilde{Q}_0}\Phi\left(|Q_0|^{\frac{s+1}{n}}|x-x_0|^{-(n+s+1)}\frac{\phi(|Q_0|)|Q_0|}{\omega(Q_0)\rho(\omega(Q_0))}\right)\omega(x)\,dx\\
&\quad\le C\sum_{k=1}^{\infty}\int_{2^kQ_0}\Phi\left(|Q_0|^{\frac{s+1}{n}}(2^kR_0)^{-(n+s+1)}\frac{\phi(|Q_0|)|Q_0|}{\omega(Q_0)\rho(\omega(Q_0))}\right)\omega(x)\,dx\\
&\quad\le C\Bigg\{\sum_{k=1}^{m_0}\int_{2^kQ_0}\Phi\left(|Q_0|^{\frac{s+1}{n}}(2^kR_0)^{-(n+s+1)}\frac{|Q_0|}{\omega(Q_0)\rho(\omega(Q_0))}\right)\omega(x)\,dx
\end{aligned}
$$

$$+\sum_{k=m_0+1}^{\infty}\int_{2^kQ_0}\cdots\Bigg\}\equiv C(\mathrm{I}_1+\mathrm{I}_2), \tag{8.42}$$

where the integer m_0 satisfies $2^{m_0-1}\le\frac{1}{R_0}<2^{m_0}$.

To estimate I_1, for any $k\in\{1,\,2,\,\cdots,\,m_0\}$, by the choice of m_0 and $R_0\in(0,1)$, we know that $2^kR_0^n\le 1$, which, together with Jensen's inequality, the lower type p_Φ property of Φ, Lemma 8.15(iii) and the fact that $(n+s+1)p_\Phi>nq(1+\alpha)$, implies that

$$\begin{aligned}\mathrm{I}_1&\lesssim\sum_{k=1}^{m_0}\omega(2^kQ_0)\Phi\left(\frac{1}{\omega(2^kQ_0)}\int_{2^kQ_0}2^{-k(n+s+1)}\{\omega(Q_0)\rho(\omega(Q_0))\}^{-1}\omega(x)\,dx\right)\\&\lesssim\sum_{k=1}^{m_0}2^{-k(n+s+1)p_\Phi}\frac{\omega(2^kQ_0)}{\omega(Q_0)}\lesssim\sum_{k=1}^{m_0}2^{knq}[\phi(|2^kQ_0|)]^q2^{-k(n+s+1)p_\Phi}\\&\lesssim\sum_{k=1}^{m_0}2^{-k[(n+s+1)p_\Phi-nq]}\lesssim 1.\end{aligned} \tag{8.43}$$

For I_2, similarly to the estimate of I_1, we have

$$\begin{aligned}\mathrm{I}_2&\lesssim\sum_{k=m_0+1}^{\infty}\omega(2^kQ_0)\Phi\left(\frac{1}{\omega(2^kQ_0)}\int_{2^kQ_0}2^{-k(n+s+1)}\{\omega(Q_0)\rho(\omega(Q_0))\}^{-1}\omega(x)\,dx\right)\\&\lesssim\sum_{k=m_0+1}^{\infty}2^{-k(n+s+1)p_\Phi}\frac{\omega(2^kQ_0)}{\omega(Q_0)}\lesssim\sum_{k=m_0+1}^{\infty}2^{knq}[\phi(|2^kQ_0|)]^q2^{-k(n+s+1)p_\Phi}\\&\lesssim\sum_{k=m_0+1}^{\infty}2^{-k[(n+s+1)p_\Phi-q(\alpha+1)]}\lesssim 1,\end{aligned}$$

which together with (8.43), (8.42) and (8.41) implies (8.40) in Case 1).

Case 2) $R_0\in[1,2]$. In this case, for any $x\in(\mathbb{R}^n\setminus\widetilde{Q}_0)$ and $z\in Q_0$, we have

$$|x-z|\sim|x-x_0|$$

and $|x-x_0|>1$. From this and [46, p. 235, (9)], we infer that for any positive integer M, there exists a positive constant $C(M)$ such that

$$|K_t(x,x-z)|\le C(M)|x-x_0|^{-M},$$

which implies that for any $x\in(\mathbb{R}^n\setminus\widetilde{Q}_0)$,

$$|\psi_t*Ta(x)|=\int_{\mathbb{R}^n}|K_t(x,x-z)a(z)|\,dz\lesssim|x-x_0|^{-M}\|a\|_{L^1(\mathbb{R}^n)}.$$

This, combined with the arbitrariness of $\psi\in\mathcal{D}_N^0(\mathbb{R}^n)$, yields that for any $x\in(\mathbb{R}^n\setminus\widetilde{Q}_0)$,

$$\mathcal{G}_N^0(Ta)(x)\lesssim|x-x_0|^{-M}\|a\|_{L^1(\mathbb{R}^n)}. \tag{8.44}$$

Take $M>\frac{nq(1+\alpha)}{p_\Phi}$. By Jensen's inequality, (8.44), Hölder's inequality and Lemma 8.15(iii), we have

$$\begin{aligned}&\int_{\mathbb{R}^n\setminus\widetilde{Q}_0}\Phi\left(\mathcal{G}_N^0(Ta)(x)\right)\omega(x)\,dx\\&\quad\lesssim\int_{\mathbb{R}^n\setminus\widetilde{Q}_0}\Phi\left(|x-x_0|^{-M}\|a\|_{L^1(\mathbb{R}^n)}\right)\omega(x)\,dx\end{aligned}$$

$$
\begin{aligned}
&\lesssim \sum_{k=1}^{\infty} \int_{2^k Q_0} \Phi\left((2^k R_0)^{-M} \frac{\phi(|Q_0|)|Q_0|}{\omega(Q_0)\rho(\omega(Q_0))}\right) \omega(x)\, dx \\
&\lesssim \sum_{k=1}^{\infty} \omega(2^k Q_0) \Phi\left((2^k R_0)^{-M} \frac{1}{\omega(Q_0)\rho(\omega(Q_0))}\right) \\
&\lesssim \sum_{k=1}^{\infty} 2^{-kMp_\Phi} R_0^{-Mp_\Phi} \frac{\omega(2^k Q_0)}{\omega(Q_0)} \lesssim \sum_{k=1}^{\infty} 2^{-k(Mp_\Phi - nq)} \phi(|2^k Q_0|) R_0^{-Mp_\Phi} \\
&\lesssim \sum_{k=1}^{\infty} 2^{-k[Mp_\Phi - nq(1+\alpha)]} R_0^{-(Mp_\Phi - nq\alpha)} \lesssim \sum_{k=1}^{\infty} 2^{-k[Mp_\Phi - nq(1+\alpha)]} \lesssim 1,
\end{aligned}
$$

which together with (8.41) implies (8.40) in Case 2). This finishes the proof of Theorem 8.18. ■

REMARK 8.19. Let $p \in (0,1]$. Theorem 8.18 when $\omega \equiv 1$ and $\Phi(t) \equiv t^p$ for all $t \in (0,\infty)$ was obtained by Goldberg [18, Theorem 4]; moreover, Theorem 8.18 when $\Phi(t) \equiv t^p$ for all $t \in (0,\infty)$ was obtained by Tang [49, Theorem 7.3]. Also, Theorem 8.18 when $\omega \in A_1(\mathbb{R}^n)$ and $\Phi(t) \equiv t$ for all $t \in (0,\infty)$ was obtained by Lee, C-C. Lin and Y.-C. Lin [32, Theorem 2].

By Theorems 8.18 and 7.5, [46, p. 233, (4)] and the proposition in [46, p. 259], we have the following result.

COROLLARY 8.20. *Let T be an $S^0_{1,\,0}(\mathbb{R}^n)$ pseudo-differential operator, Φ satisfy Assumption* (A), *$\omega \in A_\infty(\phi)$ and p_Φ be as in* (2.6). *If $p_\Phi = p_\Phi^+$ and Φ is of upper type p_Φ^+, then there exists a positive constant $C(\Phi,\,\omega)$ such that for all $f \in \mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)$,*

$$\|Tf\|_{\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)} \le C(\Phi,\,\omega)\|f\|_{\mathrm{bmo}_{\rho,\,\omega}(\mathbb{R}^n)}.$$

References

[1] P. Auscher and E. Russ, *Hardy spaces and divergence operators on strongly Lipschitz domains of* $\mathbb{R}^n$, J. Funct. Anal. 201 (2003), 148-184.

[2] Z. Birnbaum and W. Orlicz, *Über die verallgemeinerung des begriffes der zueinander konjugierten potenzen*, Studia Math. 3 (1931), 1-67.

[3] M. Bownik, *Anisotropic Hardy spaces and wavelets*, Mem. Amer. Math. Soc. 164 (2003), no. 781, 1-122.

[4] —, *Boundedness of operators on Hardy spaces via atomic decompositions*, Proc. Amer. Math. Soc. 133 (2005), 3535-3542.

[5] M. Bownik, B. Li, D. Yang and Y. Zhou, *Weighted anisotropic Hardy spaces and their applications in boundedness of sublinear operators*, Indiana Univ. Math. J. 57 (2008), 3065-3100.

[6] H. Bui, *Weighted Hardy spaces*, Math. Nachr. 103 (1981), 45-62.

[7] S.-S. Byun, F. Yao and S. Zhou, *Gradient estimates in Orlicz space for nonlinear elliptic equations*, J. Funct. Anal. 255 (2008), 1851-1873.

[8] D.-C. Chang, G. Dafni and E. M. Stein, *Hardy spaces,* BMO*, and boundary value problems for the Laplacian on a smooth domain in* $\mathbb{R}^n$, Trans. Amer. Math. Soc. 351 (1999), 1605-1661.

[9] D.-C. Chang, S. G. Krantz and E. M. Stein, H^p *theory on a smooth domain in* $\mathbb{R}^N$ *and elliptic boundary value problems*, J. Funct. Anal. 114 (1993), 286-347.

[10] R. R. Coifman and G. Weiss, *Analyse harmonique non-commutative sur certains espaces homogènes*, Lecture Notes in Math. 242, Springer, Berlin, 1971.

[11] D. Deng, X. T. Duong and L. Yan, *A characterization of the Morrey-Campanato spaces*, Math. Z. 250 (2005), 641-655.

[12] Y. Ding, M. Lee and C. Lin, *Fractional integrals on weighted Hardy spaces*, J. Math. Anal. Appl. 282 (2003), 356-368.

[13] Y. Ding and S. Lu, *Weighted norm inequalities for fractional integral operators with rough kernel*, Canad. J. Math. 50 (1998), 29-39.

[14] X. T. Duong, J. Xiao and L. Yan, *Old and new Morrey spaces with heat kernel bounds*, J. Fourier Anal. Appl. 13 (2007), 87-111.

[15] C. Fefferman and E. M. Stein, H^p *spaces of several variables*, Acta. Math. 129 (1972), 137-195.

[16] J. García-Cuerva, *Weighted* H^p *spaces*, Dissertationes Math. (Rozprawy Mat.) 162 (1979), 1-63.

[17] J. García-Cuerva and J. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, Amsterdam, North-Holland, 1985.

[18] D. Goldberg, *A local version of real Hardy spaces*, Duke Math. J. 46 (1979), 27-42.

[19] L. Grafakos, *Classical Fourier Analysis, Second Edition*, Graduate Texts in Math., No. 249, Springer, New York, 2008.

[20] L. Grafakos, L. Liu and D. Yang, *Maximal function characterizations of Hardy spaces on RD-spaces and their applications*, Sci. China Ser. A 51 (2008), 2253-2284.

[21] D. D. Haroske and L. Skrzypczak, *Entropy and approximation numbers of embeddings of function spaces with Muckenhoupt weights, I*, Rev. Mat. Complut. 21 (2008), 135-177.

[22] —, —, *Entropy and approximation numbers of embeddings of function spaces with muckenhoupt weights, II. general weights*, Ann. Acad. Sci. Fenn. Math. 36 (2011), 111-138.

[23] —, —, *Entropy numbers of embeddings of function spaces with Muckenhoupt weights, III. Some limiting cases*, J. Funct. Spaces Appl. (to appear).

[24] —, —, *Spectral theory of some degenerate elliptic operators with local singularities*, J. Math. Anal. Appl. 371 (2010), 282-299.

[25] T. Iwaniec and J. Onninen, $\mathcal{H}^1$*-estimates of Jacobians by subdeterminants*, Math. Ann. 324 (2002), 341-358.

[26] S. Janson, *Generalizations of Lipschitz spaces and an application to Hardy spaces and bounded mean oscillation*, Duke Math. J. 47 (1980), 959-982.

[27] R. Jiang and D. Yang, *New Orlicz-Hardy spaces associated with divergence form elliptic operators*, J. Funct. Anal. 258 (2010), 1167-1224.

[28] R. Jiang, D. Yang and Y. Zhou, *Orlicz-Hardy spaces associated with operators*, Sci. China Ser. A 52 (2009), 1042-1080.

[29] F. John and L. Nirenberg, *On functions of bounded mean oscillation*, Comm. Pure. Appl. Math. 14 (1961), 415-426.

[30] R. Johnson and C. J. Neugebauer, *Homeomorphisms preserving* A_p, Rev. Mat. Iberoamericana 3 (1987), 249-273.

[31] A. Jonsson, P. Sjögren and H. Wallin, *Hardy and Lipschitz spaces on subsets of* R^n, Studia

Math. 80 (1984), 141-166.

[32] M.-Y. Lee, C.-C. Lin and Y.-C. Lin, *The continuity of pseudo-differential operators on weighted local Hardy spaces*, Studia Math. 198 (2010), 69-77.

[33] S. Lu, *Four Lectures on Real H^p Spaces*, World Scientific Publishing, Singapore, 1995.

[34] S. Martínez and N. Wolanski, *A minimum problem with free boundary in Orlicz spaces*, Adv. Math. 218 (2008), 1914-1971.

[35] S. Meda, P. Sjögren and M. Vallarino, *On the H^1-L^1 boundedness of operators*, Proc. Amer. Math. Soc. 136 (2008), 2921-2931.

[36] A. Miyachi, *H^p spaces over open subsets of* $\mathbb{R}^n$, Studia Math. 95 (1990), 205-228.

[37] C. B. Morrey, *Partial regularity results for non-linear elliptic systems*, J. Math. Mech. 17 (1967/1968), 649-670.

[38] E. Nakai, *The Campanato, Morrey and Hölder spaces on spaces of homogeneous type*, Studia Math. 176 (2006), 1-19.

[39] W. Orlicz, *Über eine gewisse Klasse von Räumen vom Typus B*, Bull. Int. Acad. Pol. Ser. A 8 (1932), 207-220.

[40] M. Rao and Z. Ren, *Theory of Orlicz Spaces*, Marcel Dekker, Inc., New York, 1991.

[41] —, —, *Applications of Orlicz Spaces*, Marcel Dekker, Inc., New York, 2002.

[42] F. Ricci and J. Verdera, *Duality in spaces of finite linear combinations of atoms*, Trans. Amer. Math. Soc. 363 (2011), 1311-1323.

[43] V. S. Rychkov, *Littlewood-Paley theory and function spaces with $A_p^{\rm loc}$ weights*, Math. Nachr. 224 (2001), 145-180.

[44] T. Schott, *Function spaces with exponential weights I*, Math. Nachr. 189 (1998), 221-242.

[45] —, *Pseudodifferential operators in function spaces with exponential weights*, Math. Nachr. 200 (1999), 119-149.

[46] E. M. Stein, *Harmonic Analysis: Real-variable Methods, Orthogonality, and Oscillatory integrals*, Princeton University Press, Princeton, N. J., 1993.

[47] J.-O. Strömberg, *Bounded mean oscillation with Orlicz norms and duality of Hardy spaces*, Indiana Univ. Math. J 28 (1979), 511-544.

[48] M. H. Taibleson and G. Weiss, *The molecular characterization of certain Hardy spaces. Representation theorems for Hardy spaces*, pp. 67-149, Astérisque, 77, Soc. Math. France, Paris, 1980.

[49] L. Tang, *Weighted local Hardy spaces and their applications*, arXiv: 1004.5294.

[50] —, *Weighted norm inequalities for pseudo-differential operators with smooth symbols and their commutators*, arXiv: 1006.4685.

[51] M. E. Taylor, *Pseudodifferential operators and nonlinear PDE*, Progress in Mathematics, 100, Birkhäuser, Boston, 1991.

[52] H. Triebel, *Theory of Function Spaces*, Birkhäuser Verlag, Basel, 1983.

[53] —, *Theory of Function Spaces II*, Birkhäuser Verlag, Basel, 1992.

[54] H. Triebel and H. Winkelvoß, *Intrinsic atomic characterizations of function spaces on domains*, Math. Z. 221 (1996), 647-673.

[55] B. E. Viviani, *An atomic decomposition of the predual of* BMO(ρ), Rev. Mat. Iberoamericana 3 (1987), 401-425.

[56] D. Yang and Y. Zhou, *Boundedness of sublinear operators in Hardy spaces on RD-spaces via atoms*, J. Math. Anal. Appl. 339 (2008), 622-635.

[57] —, —, *A boundedness criterion via atoms for linear operators in Hardy spaces*, Constr. Approx. 29 (2009), 207-218.